

表 2119 身長(cm) 原疾患別（透析患者全体）

原疾患	130未満	130～	135～	140～	145～	150～	155～	160～	165～	170～	175～	180～	185～	190～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	61	204	793	3,060	7,507	12,683	14,141	15,936	14,308	9,422	3,356	822	123	13	82,429	10,918	93,347	159.86	9.46
(%)	(0.1)	(0.2)	(1.0)	(3.7)	(9.1)	(15.4)	(17.2)	(19.3)	(17.4)	(11.4)	(4.1)	(1.0)	(0.1)	(0.0)	(100.0)				
慢性腎盂腎炎	3	11	39	154	340	486	480	457	351	199	82	28	1	1	2,632	303	2,935	157.59	9.64
(%)	(0.1)	(0.4)	(1.5)	(5.9)	(12.9)	(18.5)	(18.2)	(17.4)	(13.3)	(7.6)	(3.1)	(1.1)	(0.0)	(0.0)	(100.0)				
急速進行性糸球体腎炎	1	5	36	131	229	365	393	390	307	220	62	24	4		2,167	311	2,478	158.36	9.74
(%)	(0.0)	(0.2)	(1.7)	(6.0)	(10.6)	(16.8)	(18.1)	(18.0)	(14.2)	(10.2)	(2.9)	(1.1)	(0.2)		(100.0)				
妊娠腎/妊娠中毒症	1	1	14	106	283	454	370	146	44	8	1	1			1,429	116	1,545	152.92	6.27
(%)	(0.1)	(0.1)	(1.0)	(7.4)	(19.8)	(31.8)	(25.9)	(10.2)	(3.1)	(0.6)	(0.1)	(0.1)			(100.0)				
その他の分類不能の腎炎	6	1	25	53	116	187	206	218	226	152	56	18	1		1,265	140	1,405	159.62	10.78
(%)	(0.5)	(0.1)	(2.0)	(4.2)	(9.2)	(14.8)	(16.3)	(17.2)	(17.9)	(12.0)	(4.4)	(1.4)	(0.1)		(100.0)				
多発性嚢胞腎	11	8	50	220	704	1,419	1,756	1,777	1,666	1,471	710	211	49	4	10,056	1,200	11,256	161.81	9.82
(%)	(0.1)	(0.1)	(0.5)	(2.2)	(7.0)	(14.1)	(17.5)	(17.7)	(16.6)	(14.6)	(7.1)	(2.1)	(0.5)	(0.0)	(100.0)				
腎硬化症	27	115	492	1,520	2,734	3,851	4,496	5,357	4,139	2,328	693	171	21	1	25,945	3,860	29,805	158.34	9.65
(%)	(0.1)	(0.4)	(1.9)	(5.9)	(10.5)	(14.8)	(17.3)	(20.6)	(16.0)	(9.0)	(2.7)	(0.7)	(0.1)	(0.0)	(100.0)				
悪性高血圧	1	7	27	66	165	291	342	442	451	332	154	35	5	1	2,319	299	2,618	161.62	9.71
(%)	(0.0)	(0.3)	(1.2)	(2.8)	(7.1)	(12.5)	(14.7)	(19.1)	(19.4)	(14.3)	(6.6)	(1.5)	(0.2)	(0.0)	(100.0)				
糖尿病性腎症	122	153	741	3,140	7,459	12,739	15,930	21,966	21,057	14,095	5,422	1,483	253	54	104,614	15,664	120,278	161.40	9.45
(%)	(0.1)	(0.1)	(0.7)	(3.0)	(7.1)	(12.2)	(15.2)	(21.0)	(20.1)	(13.5)	(5.2)	(1.4)	(0.2)	(0.1)	(100.0)				
SLE腎炎	3	2	27	89	248	431	431	333	210	108	46	11	1		1,940	287	2,227	156.78	8.70
(%)	(0.2)	(0.1)	(1.4)	(4.6)	(12.8)	(22.2)	(22.2)	(17.2)	(10.8)	(5.6)	(2.4)	(0.6)	(0.1)		(100.0)				
アミロイド腎	1	1	10	35	53	71	66	55	47	36	7	1			383	79	462	156.26	10.01
(%)	(0.3)	(0.3)	(2.6)	(9.1)	(13.8)	(18.5)	(17.2)	(14.4)	(12.3)	(9.4)	(1.8)	(0.3)			(100.0)				
痛風腎	1		3	16	38	76	136	239	250	142	44	8	5		958	117	1,075	163.06	8.18
(%)	(0.1)		(0.3)	(1.7)	(4.0)	(7.9)	(14.2)	(24.9)	(26.1)	(14.8)	(4.6)	(0.8)	(0.5)		(100.0)				
先天性代謝異常による腎不全	4	4	14	7	28	34	44	42	42	20	10		1		250	24	274	156.91	12.71
(%)	(1.6)	(1.6)	(5.6)	(2.8)	(11.2)	(13.6)	(17.6)	(16.8)	(16.8)	(8.0)	(4.0)		(0.4)		(100.0)				
腎・尿路結核	1		4	9	26	30	32	36	20	12	1				171	24	195	156.53	8.85
(%)	(0.6)		(2.3)	(5.3)	(15.2)	(17.5)	(18.7)	(21.1)	(11.7)	(7.0)	(0.6)				(100.0)				
腎・尿路結石	4	3	8	19	50	83	103	106	78	43	16	3			516	65	581	158.46	9.50
(%)	(0.8)	(0.6)	(1.6)	(3.7)	(9.7)	(16.1)	(20.0)	(20.5)	(15.1)	(8.3)	(3.1)	(0.6)			(100.0)				
腎・尿路腫瘍		1	9	16	52	81	127	206	190	91	26	4	1		804	106	910	161.26	8.42
(%)		(0.1)	(1.1)	(2.0)	(6.5)	(10.1)	(15.8)	(25.6)	(23.6)	(11.3)	(3.2)	(0.5)	(0.1)		(100.0)				
閉塞性尿路障害	5	5	8	27	70	87	104	121	112	72	21	2	2	1	637	93	730	158.99	10.66
(%)	(0.8)	(0.8)	(1.3)	(4.2)	(11.0)	(13.7)	(16.3)	(19.0)	(17.6)	(11.3)	(3.3)	(0.3)	(0.3)	(0.2)	(100.0)				
骨髄腫	1	1	6	14	23	30	41	53	40	29	7	1			246	45	291	158.91	9.83
(%)	(0.4)	(0.4)	(2.4)	(5.7)	(9.3)	(12.2)	(16.7)	(21.5)	(16.3)	(11.8)	(2.8)	(0.4)			(100.0)				
腎形成不全	29	10	17	43	60	63	98	107	78	66	18	8			597	63	660	155.42	16.90
(%)	(4.9)	(1.7)	(2.8)	(7.2)	(10.1)	(10.6)	(16.4)	(17.9)	(13.1)	(11.1)	(3.0)	(1.3)			(100.0)				
不明	38	103	384	1,281	2,599	3,849	4,215	4,979	4,282	2,499	911	227	30	3	25,400	4,497	29,897	158.94	9.77
(%)	(0.1)	(0.4)	(1.5)	(5.0)	(10.2)	(15.2)	(16.6)	(19.6)	(16.9)	(9.8)	(3.6)	(0.9)	(0.1)	(0.0)	(100.0)				
再導入	3	8	12	33	138	232	310	390	378	260	113	35	3	1	1,916	267	2,183	161.69	9.64
(%)	(0.2)	(0.4)	(0.6)	(1.7)	(7.2)	(12.1)	(16.2)	(20.4)	(19.7)	(13.6)	(5.9)	(1.8)	(0.2)	(0.1)	(100.0)				
その他	32	26	87	348	663	971	1,057	1,274	1,167	745	272	84	16	1	6,743	1,301	8,044	159.28	10.59
(%)	(0.5)	(0.4)	(1.3)	(5.2)	(9.8)	(14.4)	(15.7)	(18.9)	(17.3)	(11.0)	(4.0)	(1.2)	(0.2)	(0.0)	(100.0)				
合計	355	669	2,806	10,387	23,585	38,513	44,878	54,630	49,443	32,350	12,028	3,177	516	80	273,417	39,779	313,196	160.20	9.66
(%)	(0.1)	(0.2)	(1.0)	(3.8)	(8.6)	(14.1)	(16.4)	(20.0)	(18.1)	(11.8)	(4.4)	(1.2)	(0.2)	(0.0)	(100.0)				
記載なし					2	2		2	2	2					10	11	21	160.70	9.55
(%)					(20.0)	(20.0)		(20.0)	(20.0)	(20.0)					(100.0)				
総計	355	669	2,806	10,387	23,587	38,515	44,878	54,632	49,445	32,352	12,028	3,177	516	80	273,427	39,790	313,217	160.20	9.66
(%)	(0.1)	(0.2)	(1.0)	(3.8)	(8.6)	(14.1)	(16.4)	(20.0)	(18.1)	(11.8)	(4.4)	(1.2)	(0.2)	(0.0)	(100.0)				

数値下のかっこ内は行方向の合計に対する％です。

(C)Japanese Society for Dialysis Therapy

2015年調査