

表 2215 身長(cm) 原疾患別（腹膜透析患者）

原疾患	130未満	130～	135～	140～	145～	150～	155～	160～	165～	170～	175～	180～	185～	190～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	7	3	15	53	164	300	373	394	418	286	120	34	2	1	2,170	660	2,830	161.10	10.00
(%)	(0.3)	(0.1)	(0.7)	(2.4)	(7.6)	(13.8)	(17.2)	(18.2)	(19.3)	(13.2)	(5.5)	(1.6)	(0.1)	(0.0)	(100.0)				
慢性腎盂腎炎		1	1	7	11	18	17	11	9	8	2	1			86	11	97	156.57	10.10
(%)		(1.2)	(1.2)	(8.1)	(12.8)	(20.9)	(19.8)	(12.8)	(10.5)	(9.3)	(2.3)	(1.2)			(100.0)				
急速進行性糸球体腎炎	1			1	4	4	13	8	7	8	2	1			49	17	66	160.51	10.49
(%)	(2.0)			(2.0)	(8.2)	(8.2)	(26.5)	(16.3)	(14.3)	(16.3)	(4.1)	(2.0)			(100.0)				
妊娠腎/妊娠中毒症					4	13	6	3							26	1	27	153.62	4.61
(%)					(15.4)	(50.0)	(23.1)	(11.5)							(100.0)				
その他の分類不能の腎炎	6		2	2	5	7	10	11	11	5	3	2			64	14	78	154.81	22.25
(%)	(9.4)		(3.1)	(3.1)	(7.8)	(10.9)	(15.6)	(17.2)	(17.2)	(7.8)	(4.7)	(3.1)			(100.0)				
多発性囊胞腎	6		2	5	11	22	38	29	29	30	21	4	1		198	65	263	160.31	16.60
(%)	(3.0)		(1.0)	(2.5)	(5.6)	(11.1)	(19.2)	(14.6)	(14.6)	(15.2)	(10.6)	(2.0)	(0.5)		(100.0)				
腎硬化症	2	1	12	40	82	109	135	165	137	102	33	3	1		822	273	1,095	159.54	9.61
(%)	(0.2)	(0.1)	(1.5)	(4.9)	(10.0)	(13.3)	(16.4)	(20.1)	(16.7)	(12.4)	(4.0)	(0.4)	(0.1)		(100.0)				
悪性高血圧			3		2	13	9	14	15	11	5				72	35	107	161.51	9.38
(%)			(4.2)		(2.8)	(18.1)	(12.5)	(19.4)	(20.8)	(15.3)	(6.9)				(100.0)				
糖尿病性腎症	3		9	57	109	213	277	429	453	334	155	35	4	3	2,081	718	2,799	162.92	9.47
(%)	(0.1)		(0.4)	(2.7)	(5.2)	(10.2)	(13.3)	(20.6)	(21.8)	(16.0)	(7.4)	(1.7)	(0.2)	(0.1)	(100.0)				
SLE腎炎			1		2	6	9	6	8	1					33	13	46	158.36	7.51
(%)			(3.0)		(6.1)	(18.2)	(27.3)	(18.2)	(24.2)	(3.0)					(100.0)				
アミロイド腎			1	1	1		3	1	2	2	1				12	3	15	159.42	12.36
(%)			(8.3)	(8.3)	(8.3)		(25.0)	(8.3)	(16.7)	(16.7)	(8.3)				(100.0)				
痛風腎				1		3	3	5	5	7	3				27	5	32	164.53	8.63
(%)				(3.7)		(11.1)	(11.1)	(18.5)	(18.5)	(25.9)	(11.1)				(100.0)				
先天性代謝異常による腎不全	1		2		1	1	3	3	2	2					15	2	17	155.00	14.20
(%)	(6.7)		(13.3)		(6.7)	(6.7)	(20.0)	(20.0)	(13.3)	(13.3)					(100.0)				
腎・尿路結核							1								1	1	2	155.00	
(%)							(100.0)								(100.0)				
腎・尿路結石					4	3	3	3	5	1					19	2	21	157.89	8.14
(%)					(21.1)	(15.8)	(15.8)	(15.8)	(26.3)	(5.3)					(100.0)				
腎・尿路腫瘍			1		1	1	1	3	4	2					13	1	14	161.23	9.66
(%)			(7.7)		(7.7)	(7.7)	(7.7)	(23.1)	(30.8)	(15.4)					(100.0)				
閉塞性尿路障害	2					3	1	2	5	3	1				17	4	21	156.41	23.49
(%)	(11.8)					(17.6)	(5.9)	(11.8)	(29.4)	(17.6)	(5.9)				(100.0)				
骨髄腫			1		2	1			1						5	2	7	149.80	9.96
(%)			(20.0)		(40.0)	(20.0)			(20.0)						(100.0)				
腎形成不全	21	1	1	4	4	7	7	4	2	5	1	1			58	14	72	134.07	34.65
(%)	(36.2)	(1.7)	(1.7)	(6.9)	(6.9)	(12.1)	(12.1)	(6.9)	(3.4)	(8.6)	(1.7)	(1.7)			(100.0)				
不明	2	1	9	26	42	93	105	149	105	65	29	4	1		631	211	842	159.99	9.31
(%)	(0.3)	(0.2)	(1.4)	(4.1)	(6.7)	(14.7)	(16.6)	(23.6)	(16.6)	(10.3)	(4.6)	(0.6)	(0.2)		(100.0)				
再導入				1	1	7	8	6	10	3	4	2			42	13	55	162.86	9.71
(%)				(2.4)	(2.4)	(16.7)	(19.0)	(14.3)	(23.8)	(7.1)	(9.5)	(4.8)			(100.0)				
その他	17	1	6	9	14	23	21	32	29	16	14	2	1		185	78	263	154.13	21.88
(%)	(9.2)	(0.5)	(3.2)	(4.9)	(7.6)	(12.4)	(11.4)	(17.3)	(15.7)	(8.6)	(7.6)	(1.1)	(0.5)		(100.0)				
合計	68	8	65	208	464	847	1,043	1,278	1,257	891	394	89	10	4	6,626	2,143	8,769	160.74	11.58
(%)	(1.0)	(0.1)	(1.0)	(3.1)	(7.0)	(12.8)	(15.7)	(19.3)	(19.0)	(13.4)	(5.9)	(1.3)	(0.2)	(0.1)	(100.0)				
記載なし																			
(%)																			
総計	68	8	65	208	464	847	1,043	1,278	1,257	891	394	89	10	4	6,626	2,143	8,769	160.74	11.58
(%)	(1.0)	(0.1)	(1.0)	(3.1)	(7.0)	(12.8)	(15.7)	(19.3)	(19.0)	(13.4)	(5.9)	(1.3)	(0.2)	(0.1)	(100.0)				

数値下のかっこ内は行方向の合計に対する％です。
(C)Japanese Society for Dialysis Therapy
2015年調査