

表 2241 透析後体重(kg) 医療機関別（施設血液透析患者 透析回数週3回）

医療機関	20未満	20～	30～	40～	50～	60～	70～	80～	90～	100～	110～	120～	合計	記載なし	総計	平均	標準偏差
国公立大学			37	105	134	78	21	8	3	2			388	20	408	54.49	12.04
(%)			(9.5)	(27.1)	(34.5)	(20.1)	(5.4)	(2.1)	(0.8)	(0.5)			(100.0)				
私立大学		5	97	392	490	309	118	35	13	3	1	2	1,465	31	1,496	55.93	12.32
(%)		(0.3)	(6.6)	(26.8)	(33.4)	(21.1)	(8.1)	(2.4)	(0.9)	(0.2)	(0.1)	(0.1)	(100.0)				
国立		5	32	126	169	90	23	8	5	1	1		460	55	515	54.71	12.04
(%)		(1.1)	(7.0)	(27.4)	(36.7)	(19.6)	(5.0)	(1.7)	(1.1)	(0.2)	(0.2)		(100.0)				
県市町村立	5	34	1,059	3,493	4,187	2,280	865	304	102	32	10	10	12,381	589	12,970	54.84	12.40
(%)	(0.0)	(0.3)	(8.6)	(28.2)	(33.8)	(18.4)	(7.0)	(2.5)	(0.8)	(0.3)	(0.1)	(0.1)	(100.0)				
社会保険	1	6	227	742	866	493	202	75	23	7	1		2,643	32	2,675	54.98	12.36
(%)	(0.0)	(0.2)	(8.6)	(28.1)	(32.8)	(18.7)	(7.6)	(2.8)	(0.9)	(0.3)	(0.0)		(100.0)				
厚生連	2	21	549	1,686	2,057	1,189	475	150	53	14	8	5	6,209	214	6,423	55.09	12.57
(%)	(0.0)	(0.3)	(8.8)	(27.2)	(33.1)	(19.1)	(7.7)	(2.4)	(0.9)	(0.2)	(0.1)	(0.1)	(100.0)				
その他公的	1	21	530	1,830	2,178	1,249	490	145	73	20	8	5	6,550	366	6,916	55.23	12.60
(%)	(0.0)	(0.3)	(8.1)	(27.9)	(33.3)	(19.1)	(7.5)	(2.2)	(1.1)	(0.3)	(0.1)	(0.1)	(100.0)				
私立総合	2	21	406	1,246	1,436	895	372	123	41	22	7	2	4,573	216	4,789	55.41	13.10
(%)	(0.0)	(0.5)	(8.9)	(27.2)	(31.4)	(19.6)	(8.1)	(2.7)	(0.9)	(0.5)	(0.2)	(0.0)	(100.0)				
私立	31	378	7,025	18,571	20,035	11,528	4,326	1,498	567	195	81	55	64,290	2,307	66,597	54.15	12.93
(%)	(0.0)	(0.6)	(10.9)	(28.9)	(31.2)	(17.9)	(6.7)	(2.3)	(0.9)	(0.3)	(0.1)	(0.1)	(100.0)				
私立診療所	45	301	8,626	29,785	38,364	25,139	10,212	3,857	1,404	510	215	110	118,568	2,495	121,063	56.56	13.15
(%)	(0.0)	(0.3)	(7.3)	(25.1)	(32.4)	(21.2)	(8.6)	(3.3)	(1.2)	(0.4)	(0.2)	(0.1)	(100.0)				
総計	87	792	18,588	57,976	69,916	43,250	17,104	6,203	2,284	806	332	189	217,527	6,325	223,852	55.61	13.03
(%)	(0.0)	(0.4)	(8.5)	(26.7)	(32.1)	(19.9)	(7.9)	(2.9)	(1.0)	(0.4)	(0.2)	(0.1)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2015年調査