

表 2250 体重(kg) 透析歴(年)別（腹膜透析患者）

透析歴	20未満	20～	30～	40～	50～	60～	70～	80～	90～	100～	110～	120～	合計	記載なし	総計	平均	標準偏差
2年未満	13	11	66	366	616	654	347	126	50	15	10	1	2,275	1,405	3,680	61.43	14.36
(%)	(0.6)	(0.5)	(2.9)	(16.1)	(27.1)	(28.7)	(15.3)	(5.5)	(2.2)	(0.7)	(0.4)	(0.0)	(100.0)				
2年～	20	7	49	240	552	607	324	99	39	9	6	2	1,954	1,148	3,102	61.88	14.04
(%)	(1.0)	(0.4)	(2.5)	(12.3)	(28.2)	(31.1)	(16.6)	(5.1)	(2.0)	(0.5)	(0.3)	(0.1)	(100.0)				
5年～	8	5	22	194	311	274	127	36	16	6	3	1	1,003	517	1,520	59.40	13.44
(%)	(0.8)	(0.5)	(2.2)	(19.3)	(31.0)	(27.3)	(12.7)	(3.6)	(1.6)	(0.6)	(0.3)	(0.1)	(100.0)				
10年～	2	2	4	46	82	64	23	8	3	1			235	79	314	58.39	12.41
(%)	(0.9)	(0.9)	(1.7)	(19.6)	(34.9)	(27.2)	(9.8)	(3.4)	(1.3)	(0.4)			(100.0)				
15年～			2	18	21	17	3	1					62	25	87	55.85	9.67
(%)			(3.2)	(29.0)	(33.9)	(27.4)	(4.8)	(1.6)					(100.0)				
20年～		1	1	6	5	3	1						17	9	26	50.83	13.24
(%)		(5.9)	(5.9)	(35.3)	(29.4)	(17.6)	(5.9)						(100.0)				
25年～				2	1	3							6	6	12	55.38	9.70
(%)				(33.3)	(16.7)	(50.0)							(100.0)				
30年～				1	2								3	3	6	49.70	7.54
(%)				(33.3)	(66.7)								(100.0)				
35年～			1				1						2		2	58.20	29.84
(%)			(50.0)				(50.0)						(100.0)				
40年～				1									1	1	2	46.30	
(%)				(100.0)									(100.0)				
合計	43	26	145	874	1,590	1,622	826	270	108	31	19	4	5,558	3,193	8,751	60.98	14.01
(%)	(0.8)	(0.5)	(2.6)	(15.7)	(28.6)	(29.2)	(14.9)	(4.9)	(1.9)	(0.6)	(0.3)	(0.1)	(100.0)				
不明				1	1	1							3	15	18	54.37	6.50
(%)				(33.3)	(33.3)	(33.3)							(100.0)				
記載なし																	
(%)																	
総計	43	26	145	875	1,591	1,623	826	270	108	31	19	4	5,561	3,208	8,769	60.98	14.01
(%)	(0.8)	(0.5)	(2.6)	(15.7)	(28.6)	(29.2)	(14.9)	(4.9)	(1.9)	(0.6)	(0.3)	(0.1)	(100.0)				
平均	3.00	4.00	3.06	3.62	3.34	3.02	2.70	2.49	2.40	2.58	2.26	2.75	3.12	2.78	2.99		
標準偏差	2.89	4.96	4.58	4.39	3.73	3.39	3.05	2.72	2.64	2.60	2.54	2.22	3.62	3.42	3.55		

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2015年調査