

表 2257 体重減少量(kg) 医療機関別（施設血液透析患者 透析回数週3回 透析歴2年以上）

医療機関	0.0未満	0.0～	1.0～	2.0～	3.0～	4.0～	5.0～	6.0～	7.0～	8.0～	合計	記載なし	総計	平均	標準偏差
国公立大学	2	28	91	80	31	2	2				236	9	245	1.99	0.95
(%)	(0.8)	(11.9)	(38.6)	(33.9)	(13.1)	(0.8)	(0.8)				(100.0)				
私立大学	11	70	259	389	177	48	6	2		1	963	17	980	2.32	1.07
(%)	(1.1)	(7.3)	(26.9)	(40.4)	(18.4)	(5.0)	(0.6)	(0.2)		(0.1)	(100.0)				
国立	3	15	63	114	84	10	6	1			296	41	337	2.53	1.07
(%)	(1.0)	(5.1)	(21.3)	(38.5)	(28.4)	(3.4)	(2.0)	(0.3)			(100.0)				
県市町村立	58	474	2,256	4,105	2,021	411	63	5	1	8	9,402	415	9,817	2.43	0.98
(%)	(0.6)	(5.0)	(24.0)	(43.7)	(21.5)	(4.4)	(0.7)	(0.1)	(0.0)	(0.1)	(100.0)				
社会保険	10	98	491	872	446	85	13	2	1	1	2,019	26	2,045	2.43	0.95
(%)	(0.5)	(4.9)	(24.3)	(43.2)	(22.1)	(4.2)	(0.6)	(0.1)	(0.0)	(0.0)	(100.0)				
厚生連	16	171	1,002	2,079	1,112	263	50	11	1	5	4,710	163	4,873	2.56	1.00
(%)	(0.3)	(3.6)	(21.3)	(44.1)	(23.6)	(5.6)	(1.1)	(0.2)	(0.0)	(0.1)	(100.0)				
その他公的	28	255	1,195	2,223	1,071	207	45	7	3	3	5,037	217	5,254	2.42	0.99
(%)	(0.6)	(5.1)	(23.7)	(44.1)	(21.3)	(4.1)	(0.9)	(0.1)	(0.1)	(0.1)	(100.0)				
私立総合	12	171	837	1,445	714	162	37	5	2	3	3,388	69	3,457	2.45	1.01
(%)	(0.4)	(5.0)	(24.7)	(42.7)	(21.1)	(4.8)	(1.1)	(0.1)	(0.1)	(0.1)	(100.0)				
私立	257	3,315	13,020	19,680	9,782	2,495	504	110	27	41	49,231	1,764	50,995	2.38	1.06
(%)	(0.5)	(6.7)	(26.4)	(40.0)	(19.9)	(5.1)	(1.0)	(0.2)	(0.1)	(0.1)	(100.0)				
私立診療所	342	3,511	20,900	39,728	21,214	5,616	1,161	225	59	70	92,826	1,915	94,741	2.55	1.01
(%)	(0.4)	(3.8)	(22.5)	(42.8)	(22.9)	(6.1)	(1.3)	(0.2)	(0.1)	(0.1)	(100.0)				
総計	739	8,108	40,114	70,715	36,652	9,299	1,887	368	94	132	168,108	4,636	172,744	2.48	1.03
(%)	(0.4)	(4.8)	(23.9)	(42.1)	(21.8)	(5.5)	(1.1)	(0.2)	(0.1)	(0.1)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査