

表 2261 体重減少量(kg) 原疾患別（血液透析濾過患者 透析回数週3回 透析歴2年以上）

原疾患	0.0未満	0.0～	1.0～	2.0～	3.0～	4.0～	5.0～	6.0～	7.0～	8.0～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	46	518	3,732	7,059	3,631	805	131	27	7	6	15,962	364	16,326	2.51	0.94
(%)	(0.3)	(3.2)	(23.4)	(44.2)	(22.7)	(5.0)	(0.8)	(0.2)	(0.0)	(0.0)	(100.0)				
慢性腎盂腎炎	1	16	128	192	86	25	1				449	7	456	2.40	0.89
(%)	(0.2)	(3.6)	(28.5)	(42.8)	(19.2)	(5.6)	(0.2)				(100.0)				
急速進行性糸球体腎炎	6	12	59	140	54	23	1				295		295	2.44	1.00
(%)	(2.0)	(4.1)	(20.0)	(47.5)	(18.3)	(7.8)	(0.3)				(100.0)				
妊娠腎/妊娠中毒症	2	17	108	186	59	7	1				380	9	389	2.25	0.78
(%)	(0.5)	(4.5)	(28.4)	(48.9)	(15.5)	(1.8)	(0.3)				(100.0)				
その他の分類不能の腎炎	4	8	46	108	52	16	4				238	4	242	2.52	1.02
(%)	(1.7)	(3.4)	(19.3)	(45.4)	(21.8)	(6.7)	(1.7)				(100.0)				
多発性嚢胞腎	11	58	435	732	368	60	11	1	2	1	1,679	29	1,708	2.44	1.02
(%)	(0.7)	(3.5)	(25.9)	(43.6)	(21.9)	(3.6)	(0.7)	(0.1)	(0.1)	(0.1)	(100.0)				
腎硬化症	10	129	721	1,221	637	147	25	10	2		2,902	86	2,988	2.47	0.98
(%)	(0.3)	(4.4)	(24.8)	(42.1)	(22.0)	(5.1)	(0.9)	(0.3)	(0.1)		(100.0)				
悪性高血圧	1	5	52	133	97	33	5	1		1	328	7	335	2.86	1.11
(%)	(0.3)	(1.5)	(15.9)	(40.5)	(29.6)	(10.1)	(1.5)	(0.3)		(0.3)	(100.0)				
糖尿病性腎症	46	437	2,445	5,432	3,709	1,215	269	50	16	12	13,631	344	13,975	2.73	1.06
(%)	(0.3)	(3.2)	(17.9)	(39.9)	(27.2)	(8.9)	(2.0)	(0.4)	(0.1)	(0.1)	(100.0)				
SLE腎炎	2	19	115	161	56	16					369	9	378	2.26	0.86
(%)	(0.5)	(5.1)	(31.2)	(43.6)	(15.2)	(4.3)					(100.0)				
アミロイド腎	1	2	13	22	8						46	3	49	2.12	0.84
(%)	(2.2)	(4.3)	(28.3)	(47.8)	(17.4)						(100.0)				
痛風腎	1	5	44	66	31	9					156	2	158	2.43	0.98
(%)	(0.6)	(3.2)	(28.2)	(42.3)	(19.9)	(5.8)					(100.0)				
先天性代謝異常による腎不全		1	8	24	8	2	1				44	2	46	2.62	0.96
(%)		(2.3)	(18.2)	(54.5)	(18.2)	(4.5)	(2.3)				(100.0)				
腎・尿路結核	1	1	11	9	6						28		28	2.02	1.00
(%)	(3.6)	(3.6)	(39.3)	(32.1)	(21.4)						(100.0)				
腎・尿路結石		2	17	30	17	3					69	1	70	2.44	0.89
(%)		(2.9)	(24.6)	(43.5)	(24.6)	(4.3)					(100.0)				
腎・尿路腫瘍		3	17	41	25	1	1				88	2	90	2.50	0.82
(%)		(3.4)	(19.3)	(46.6)	(28.4)	(1.1)	(1.1)				(100.0)				
閉塞性尿路障害		6	27	33	22	7		1			96		96	2.48	1.06
(%)		(6.3)	(28.1)	(34.4)	(22.9)	(7.3)		(1.0)			(100.0)				
骨髄腫		5	7	5	7						24	1	25	2.09	1.13
(%)		(20.8)	(29.2)	(20.8)	(29.2)						(100.0)				
腎形成不全		3	28	59	29	4	1				124	2	126	2.50	0.84
(%)		(2.4)	(22.6)	(47.6)	(23.4)	(3.2)	(0.8)				(100.0)				
不明	10	171	874	1,629	860	223	44	5	1	4	3,821	120	3,941	2.53	0.99
(%)	(0.3)	(4.5)	(22.9)	(42.6)	(22.5)	(5.8)	(1.2)	(0.1)	(0.0)	(0.1)	(100.0)				
再導入	1	9	83	191	107	31	4	1		1	428	8	436	2.65	1.08
(%)	(0.2)	(2.1)	(19.4)	(44.6)	(25.0)	(7.2)	(0.9)	(0.2)		(0.2)	(100.0)				
その他	4	46	232	367	222	53	12	3			939	20	959	2.51	1.02
(%)	(0.4)	(4.9)	(24.7)	(39.1)	(23.6)	(5.6)	(1.3)	(0.3)			(100.0)				
合計	147	1,473	9,202	17,840	10,091	2,680	511	99	28	25	42,096	1,020	43,116	2.57	1.00
(%)	(0.3)	(3.5)	(21.9)	(42.4)	(24.0)	(6.4)	(1.2)	(0.2)	(0.1)	(0.1)	(100.0)				
記載なし				4	1	1					6		6	2.95	0.82
(%)				(66.7)	(16.7)	(16.7)					(100.0)				
総計	147	1,473	9,202	17,844	10,092	2,681	511	99	28	25	42,102	1,020	43,122	2.57	1.00
(%)	(0.3)	(3.5)	(21.9)	(42.4)	(24.0)	(6.4)	(1.2)	(0.2)	(0.1)	(0.1)	(100.0)				

数値下のかつこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査