

表 2266 体重減少率(%) 原疾患別（施設血液透析患者 透析回数週3回 透析歴2年以上）

原疾患	1.0未満	1.0～	2.0～	3.0～	4.0～	5.0～	6.0～	7.0～	8.0～	9.0～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	914	1,842	4,971	10,330	14,258	11,564	6,056	2,302	769	384	53,390	1,397	54,787	4.62	1.69
(%)	(1.7)	(3.5)	(9.3)	(19.3)	(26.7)	(21.7)	(11.3)	(4.3)	(1.4)	(0.7)	(100.0)				
慢性腎盂腎炎	50	73	174	336	411	344	171	69	34	19	1,681	34	1,715	4.51	1.78
(%)	(3.0)	(4.3)	(10.4)	(20.0)	(24.4)	(20.5)	(10.2)	(4.1)	(2.0)	(1.1)	(100.0)				
急速進行性糸球体腎炎	24	46	117	222	273	274	169	65	18	13	1,221	36	1,257	4.70	1.71
(%)	(2.0)	(3.8)	(9.6)	(18.2)	(22.4)	(22.4)	(13.8)	(5.3)	(1.5)	(1.1)	(100.0)				
妊娠腎/妊娠中毒症	10	27	77	165	252	195	133	27	12	11	909	22	931	4.73	1.57
(%)	(1.1)	(3.0)	(8.5)	(18.2)	(27.7)	(21.5)	(14.6)	(3.0)	(1.3)	(1.2)	(100.0)				
その他の分類不能の腎炎	22	19	83	131	188	144	80	39	16	6	728	9	737	4.60	1.76
(%)	(3.0)	(2.6)	(11.4)	(18.0)	(25.8)	(19.8)	(11.0)	(5.4)	(2.2)	(0.8)	(100.0)				
多発性嚢胞腎	122	278	755	1,516	1,865	1,220	500	165	37	23	6,481	157	6,638	4.29	1.51
(%)	(1.9)	(4.3)	(11.6)	(23.4)	(28.8)	(18.8)	(7.7)	(2.5)	(0.6)	(0.4)	(100.0)				
腎硬化症	323	602	1,550	3,058	3,967	2,854	1,381	534	180	82	14,531	351	14,882	4.44	1.67
(%)	(2.2)	(4.1)	(10.7)	(21.0)	(27.3)	(19.6)	(9.5)	(3.7)	(1.2)	(0.6)	(100.0)				
悪性高血圧	28	40	125	247	357	340	168	67	24	15	1,411	40	1,451	4.74	1.77
(%)	(2.0)	(2.8)	(8.9)	(17.5)	(25.3)	(24.1)	(11.9)	(4.7)	(1.7)	(1.1)	(100.0)				
糖尿病性腎症	1,263	2,522	6,538	13,329	17,436	12,977	6,409	2,356	761	393	63,984	1,908	65,892	4.48	1.65
(%)	(2.0)	(3.9)	(10.2)	(20.8)	(27.3)	(20.3)	(10.0)	(3.7)	(1.2)	(0.6)	(100.0)				
SLE腎炎	33	38	116	196	283	243	146	62	15	10	1,142	26	1,168	4.64	1.72
(%)	(2.9)	(3.3)	(10.2)	(17.2)	(24.8)	(21.3)	(12.8)	(5.4)	(1.3)	(0.9)	(100.0)				
アミロイド腎	11	5	19	37	52	41	24	16	2	2	209	7	216	4.57	2.01
(%)	(5.3)	(2.4)	(9.1)	(17.7)	(24.9)	(19.6)	(11.5)	(7.7)	(1.0)	(1.0)	(100.0)				
痛風腎	16	28	61	138	150	157	59	18	8	1	636	16	652	4.42	1.60
(%)	(2.5)	(4.4)	(9.6)	(21.7)	(23.6)	(24.7)	(9.3)	(2.8)	(1.3)	(0.2)	(100.0)				
先天性代謝異常による腎不全	6	7	14	22	35	22	23	10	5	4	148	5	153	4.78	2.05
(%)	(4.1)	(4.7)	(9.5)	(14.9)	(23.6)	(14.9)	(15.5)	(6.8)	(3.4)	(2.7)	(100.0)				
腎・尿路結核	5	3	16	21	29	32	10	2	2		120	3	123	4.37	1.63
(%)	(4.2)	(2.5)	(13.3)	(17.5)	(24.2)	(26.7)	(8.3)	(1.7)	(1.7)		(100.0)				
腎・尿路結石	10	16	35	61	89	63	26	11	1	1	313	6	319	4.29	1.59
(%)	(3.2)	(5.1)	(11.2)	(19.5)	(28.4)	(20.1)	(8.3)	(3.5)	(0.3)	(0.3)	(100.0)				
腎・尿路腫瘍	9	15	46	91	126	103	41	13	4	3	451	8	459	4.49	1.83
(%)	(2.0)	(3.3)	(10.2)	(20.2)	(27.9)	(22.8)	(9.1)	(2.9)	(0.9)	(0.7)	(100.0)				
閉塞性尿路障害	14	22	41	63	83	72	42	24	7	4	372	5	377	4.54	1.90
(%)	(3.8)	(5.9)	(11.0)	(16.9)	(22.3)	(19.4)	(11.3)	(6.5)	(1.9)	(1.1)	(100.0)				
骨髄腫	9	7	8	11	23	12	8	5			83	4	87	3.97	2.05
(%)	(10.8)	(8.4)	(9.6)	(13.3)	(27.7)	(14.5)	(9.6)	(6.0)			(100.0)				
腎形成不全	7	7	19	66	78	89	39	22	10	2	339	5	344	4.88	1.62
(%)	(2.1)	(2.1)	(5.6)	(19.5)	(23.0)	(26.3)	(11.5)	(6.5)	(2.9)	(0.6)	(100.0)				
不明	411	666	1,507	2,972	3,816	2,963	1,574	576	225	118	14,828	465	15,293	4.48	1.75
(%)	(2.8)	(4.5)	(10.2)	(20.0)	(25.7)	(20.0)	(10.6)	(3.9)	(1.5)	(0.8)	(100.0)				
再導入	7	29	88	216	272	252	140	55	16	12	1,087	21	1,108	4.80	1.60
(%)	(0.6)	(2.7)	(8.1)	(19.9)	(25.0)	(23.2)	(12.9)	(5.1)	(1.5)	(1.1)	(100.0)				
その他	112	198	408	800	996	810	415	172	77	31	4,019	133	4,152	4.47	1.78
(%)	(2.8)	(4.9)	(10.2)	(19.9)	(24.8)	(20.2)	(10.3)	(4.3)	(1.9)	(0.8)	(100.0)				
合計	3,406	6,490	16,768	34,028	45,039	34,771	17,614	6,610	2,223	1,134	168,083	4,658	172,741	4.52	1.68
(%)	(2.0)	(3.9)	(10.0)	(20.2)	(26.8)	(20.7)	(10.5)	(3.9)	(1.3)	(0.7)	(100.0)				
記載なし		1	1							1	3		3	4.65	4.60
(%)		(33.3)	(33.3)							(33.3)	(100.0)				
総計	3,406	6,491	16,769	34,028	45,039	34,771	17,614	6,610	2,223	1,135	168,086	4,658	172,744	4.52	1.68
(%)	(2.0)	(3.9)	(10.0)	(20.2)	(26.8)	(20.7)	(10.5)	(3.9)	(1.3)	(0.7)	(100.0)				

数値下のかつこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査