

表 2300 体重減少率(%) 原疾患別（血液透析濾過患者 透析回数週3回 透析歴2年以上）

原疾患	1.0未満	1.0～	2.0～	3.0～	4.0～	5.0～	6.0～	7.0～	8.0～	9.0～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	208	429	1,449	3,143	4,440	3,551	1,823	638	197	83	15,961	365	16,326	4.63	1.55
(%)	(1.3)	(2.7)	(9.1)	(19.7)	(27.8)	(22.2)	(11.4)	(4.0)	(1.2)	(0.5)	(100.0)				
慢性腎盂腎炎	4	16	39	90	103	112	56	21	8		449	7	456	4.68	1.56
(%)	(0.9)	(3.6)	(8.7)	(20.0)	(22.9)	(24.9)	(12.5)	(4.7)	(1.8)		(100.0)				
急速進行性糸球体腎炎	8	11	24	47	81	66	35	15	5	3	295		295	4.66	1.78
(%)	(2.7)	(3.7)	(8.1)	(15.9)	(27.5)	(22.4)	(11.9)	(5.1)	(1.7)	(1.0)	(100.0)				
妊娠腎/妊娠中毒症	6	12	30	52	123	89	42	20	2	4	380	9	389	4.73	1.57
(%)	(1.6)	(3.2)	(7.9)	(13.7)	(32.4)	(23.4)	(11.1)	(5.3)	(0.5)	(1.1)	(100.0)				
その他の分類不能の腎炎	5	6	19	48	60	58	23	14	3	2	238	4	242	4.68	1.71
(%)	(2.1)	(2.5)	(8.0)	(20.2)	(25.2)	(24.4)	(9.7)	(5.9)	(1.3)	(0.8)	(100.0)				
多発性嚢胞腎	33	60	205	396	479	334	113	39	12	7	1,678	30	1,708	4.29	1.56
(%)	(2.0)	(3.6)	(12.2)	(23.6)	(28.5)	(19.9)	(6.7)	(2.3)	(0.7)	(0.4)	(100.0)				
腎硬化症	54	98	283	638	818	582	289	96	27	17	2,902	86	2,988	4.48	1.57
(%)	(1.9)	(3.4)	(9.8)	(22.0)	(28.2)	(20.1)	(10.0)	(3.3)	(0.9)	(0.6)	(100.0)				
悪性高血圧	2	8	20	56	118	75	32	13	3		327	8	335	4.68	1.33
(%)	(0.6)	(2.4)	(6.1)	(17.1)	(36.1)	(22.9)	(9.8)	(4.0)	(0.9)		(100.0)				
糖尿病性腎症	211	437	1,275	2,905	3,942	2,855	1,299	471	167	69	13,631	344	13,975	4.53	1.59
(%)	(1.5)	(3.2)	(9.4)	(21.3)	(28.9)	(20.9)	(9.5)	(3.5)	(1.2)	(0.5)	(100.0)				
SLE腎炎	5	10	40	63	97	76	51	18	5	4	369	9	378	4.68	1.64
(%)	(1.4)	(2.7)	(10.8)	(17.1)	(26.3)	(20.6)	(13.8)	(4.9)	(1.4)	(1.1)	(100.0)				
アミロイド腎	1	3	4	5	10	16	5	1	1		46	3	49	4.54	1.71
(%)	(2.2)	(6.5)	(8.7)	(10.9)	(21.7)	(34.8)	(10.9)	(2.2)	(2.2)		(100.0)				
痛風腎	3	6	23	45	34	28	13	3	1		156	2	158	4.16	1.62
(%)	(1.9)	(3.8)	(14.7)	(28.8)	(21.8)	(17.9)	(8.3)	(1.9)	(0.6)		(100.0)				
先天性代謝異常による腎不全			4	4	12	10	6	6	1	1	44	2	46	5.31	1.75
(%)			(9.1)	(9.1)	(27.3)	(22.7)	(13.6)	(13.6)	(2.3)	(2.3)	(100.0)				
腎・尿路結核	2	2	2	7	5	3	6	1			28		28	4.12	2.07
(%)	(7.1)	(7.1)	(7.1)	(25.0)	(17.9)	(10.7)	(21.4)	(3.6)			(100.0)				
腎・尿路結石	1	2	4	15	20	20	7				69	1	70	4.55	1.30
(%)	(1.4)	(2.9)	(5.8)	(21.7)	(29.0)	(29.0)	(10.1)				(100.0)				
腎・尿路腫瘍	1	3	8	19	30	18	7	2			88	2	90	4.42	1.35
(%)	(1.1)	(3.4)	(9.1)	(21.6)	(34.1)	(20.5)	(8.0)	(2.3)			(100.0)				
閉塞性尿路障害	2	4	11	17	23	21	10	4	3	1	96		96	4.64	1.82
(%)	(2.1)	(4.2)	(11.5)	(17.7)	(24.0)	(21.9)	(10.4)	(4.2)	(3.1)	(1.0)	(100.0)				
骨髄腫	2	3	4	3	5	5	1	1			24	1	25	3.67	1.79
(%)	(8.3)	(12.5)	(16.7)	(12.5)	(20.8)	(20.8)	(4.2)	(4.2)			(100.0)				
腎形成不全		3	10	20	29	38	15	5	3	1	124	2	126	4.89	1.54
(%)		(2.4)	(8.1)	(16.1)	(23.4)	(30.6)	(12.1)	(4.0)	(2.4)	(0.8)	(100.0)				
不明	57	150	335	751	1,043	833	438	140	55	19	3,821	120	3,941	4.60	1.63
(%)	(1.5)	(3.9)	(8.8)	(19.7)	(27.3)	(21.8)	(11.5)	(3.7)	(1.4)	(0.5)	(100.0)				
再導入	5	5	29	89	99	114	50	25	9	2	427	9	436	4.85	1.53
(%)	(1.2)	(1.2)	(6.8)	(20.8)	(23.2)	(26.7)	(11.7)	(5.9)	(2.1)	(0.5)	(100.0)				
その他	15	47	82	207	242	186	105	35	12	8	939	20	959	4.53	1.65
(%)	(1.6)	(5.0)	(8.7)	(22.0)	(25.8)	(19.8)	(11.2)	(3.7)	(1.3)	(0.9)	(100.0)				
合計	625	1,315	3,900	8,620	11,813	9,090	4,426	1,568	514	221	42,092	1,024	43,116	4.57	1.58
(%)	(1.5)	(3.1)	(9.3)	(20.5)	(28.1)	(21.6)	(10.5)	(3.7)	(1.2)	(0.5)	(100.0)				
記載なし				1	2	2	1				6		6	5.02	1.15
(%)				(16.7)	(33.3)	(33.3)	(16.7)				(100.0)				
総計	625	1,315	3,900	8,621	11,815	9,092	4,427	1,568	514	221	42,098	1,024	43,122	4.57	1.58
(%)	(1.5)	(3.1)	(9.3)	(20.5)	(28.1)	(21.6)	(10.5)	(3.7)	(1.2)	(0.5)	(100.0)				

数値下のかつこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy
2015年調査