

表 2332 Body Mass Index 治療方法(新分類)別(透析患者全体)

治療方法 (新分類)	14未満	14～	16～	18～	20～	22～	24～	26～	28～	30～	合計	記載なし	総計	平均	標準偏差
施設血液透析	1,444	7,878	24,923	43,673	47,938	37,004	22,316	11,935	6,233	5,449	208,793	39,932	248,725	21.47	3.69
(%)	(0.7)	(3.8)	(11.9)	(20.9)	(23.0)	(17.7)	(10.7)	(5.7)	(3.0)	(2.6)	(100.0)				
Off-line HDF	29	200	637	1,006	1,043	729	460	236	124	91	4,555	777	5,332	21.16	3.63
(%)	(0.6)	(4.4)	(14.0)	(22.1)	(22.9)	(16.0)	(10.1)	(5.2)	(2.7)	(2.0)	(100.0)				
On-line HDF	147	1,014	3,963	7,713	8,721	7,090	4,302	2,509	1,384	1,206	38,049	6,478	44,527	21.84	3.71
(%)	(0.4)	(2.7)	(10.4)	(20.3)	(22.9)	(18.6)	(11.3)	(6.6)	(3.6)	(3.2)	(100.0)				
Push/Pull HDF	1	4	11	23	23	15	10	6	1	2	96	14	110	21.16	3.54
(%)	(1.0)	(4.2)	(11.5)	(24.0)	(24.0)	(15.6)	(10.4)	(6.3)	(1.0)	(2.1)	(100.0)				
ハイオフィльтраーション	4	7	27	45	64	36	23	14	4	6	230	37	267	21.21	3.83
(%)	(1.7)	(3.0)	(11.7)	(19.6)	(27.8)	(15.7)	(10.0)	(6.1)	(1.7)	(2.6)	(100.0)				
IHDF	15	104	391	682	764	599	346	202	90	76	3,269	271	3,540	21.50	3.58
(%)	(0.5)	(3.2)	(12.0)	(20.9)	(23.4)	(18.3)	(10.6)	(6.2)	(2.8)	(2.3)	(100.0)				
血液濾過	1		3	1	1	4	3				13	3	16	20.49	4.40
(%)	(7.7)		(23.1)	(7.7)	(7.7)	(30.8)	(23.1)				(100.0)				
血液吸着透析 (リクセル等使用)	7	40	249	363	313	172	87	37	9	8	1,285	91	1,376	20.22	2.95
(%)	(0.5)	(3.1)	(19.4)	(28.2)	(24.4)	(13.4)	(6.8)	(2.9)	(0.7)	(0.6)	(100.0)				
在宅血液透析	1	2	27	63	76	74	62	32	12	14	363	192	555	22.57	3.51
(%)	(0.3)	(0.6)	(7.4)	(17.4)	(20.9)	(20.4)	(17.1)	(8.8)	(3.3)	(3.9)	(100.0)				
腹膜透析 (CAPD)	11	22	120	338	595	719	512	276	165	115	2,873	2,029	4,902	23.21	3.51
(%)	(0.4)	(0.8)	(4.2)	(11.8)	(20.7)	(25.0)	(17.8)	(9.6)	(5.7)	(4.0)	(100.0)				
腹膜透析 (APD)	12	43	88	184	370	375	289	175	109	83	1,728	1,047	2,775	23.11	3.90
(%)	(0.7)	(2.5)	(5.1)	(10.6)	(21.4)	(21.7)	(16.7)	(10.1)	(6.3)	(4.8)	(100.0)				
腹膜透析 (CCPD)	2	7	18	84	138	193	147	63	40	34	726	366	1,092	23.36	3.49
(%)	(0.3)	(1.0)	(2.5)	(11.6)	(19.0)	(26.6)	(20.2)	(8.7)	(5.5)	(4.7)	(100.0)				
合計	1,674	9,321	30,457	54,175	60,046	47,010	28,557	15,485	8,171	7,084	261,980	51,237	313,217	21.55	3.70
(%)	(0.6)	(3.6)	(11.6)	(20.7)	(22.9)	(17.9)	(10.9)	(5.9)	(3.1)	(2.7)	(100.0)				
記載なし															
(%)															
総計	1,674	9,321	30,457	54,175	60,046	47,010	28,557	15,485	8,171	7,084	261,980	51,237	313,217	21.55	3.70
(%)	(0.6)	(3.6)	(11.6)	(20.7)	(22.9)	(17.9)	(10.9)	(5.9)	(3.1)	(2.7)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

*腹膜透析(CCPD)は、APDと手動のバック交換の両方を行うもの

(C)Japanese Society for Dialysis Therapy

2015年調査