

表 2387 Body Mass Index 原疾患別（血液透析濾過患者 透析回数週3回）

原疾患	14未満	14～	16～	18～	20～	22～	24～	26～	28～	30～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	76	582	2,187	3,860	3,950	2,792	1,389	713	351	273	16,173	1,289	17,462	21.03	3.39
(%)	(0.5)	(3.6)	(13.5)	(23.9)	(24.4)	(17.3)	(8.6)	(4.4)	(2.2)	(1.7)	(100.0)				
慢性腎盂腎炎	4	27	75	107	97	73	36	14	9	7	449	35	484	20.58	3.46
(%)	(0.9)	(6.0)	(16.7)	(23.8)	(21.6)	(16.3)	(8.0)	(3.1)	(2.0)	(1.6)	(100.0)				
急速進行性糸球体腎炎	3	17	56	85	67	59	20	6	3	10	326	27	353	20.53	3.48
(%)	(0.9)	(5.2)	(17.2)	(26.1)	(20.6)	(18.1)	(6.1)	(1.8)	(0.9)	(3.1)	(100.0)				
妊娠腎/妊娠中毒症	4	23	67	96	76	45	24	17	6	10	368	33	401	20.47	3.71
(%)	(1.1)	(6.3)	(18.2)	(26.1)	(20.7)	(12.2)	(6.5)	(4.6)	(1.6)	(2.7)	(100.0)				
その他の分類不能の腎炎	5	11	27	74	54	29	25	14	3	2	244	17	261	20.72	3.45
(%)	(2.0)	(4.5)	(11.1)	(30.3)	(22.1)	(11.9)	(10.2)	(5.7)	(1.2)	(0.8)	(100.0)				
多発性嚢胞腎	1	39	154	397	485	360	166	88	32	24	1,746	134	1,880	21.44	3.10
(%)	(0.1)	(2.2)	(8.8)	(22.7)	(27.8)	(20.6)	(9.5)	(5.0)	(1.8)	(1.4)	(100.0)				
腎硬化症	17	99	366	707	833	652	379	223	106	79	3,461	359	3,820	21.66	3.55
(%)	(0.5)	(2.9)	(10.6)	(20.4)	(24.1)	(18.8)	(11.0)	(6.4)	(3.1)	(2.3)	(100.0)				
悪性高血圧		11	31	58	82	67	45	27	24	18	363	28	391	22.55	4.05
(%)		(3.0)	(8.5)	(16.0)	(22.6)	(18.5)	(12.4)	(7.4)	(6.6)	(5.0)	(100.0)				
糖尿病性腎症	38	240	1,072	2,340	3,180	3,074	2,292	1,422	832	770	15,260	1,700	16,960	22.85	3.87
(%)	(0.2)	(1.6)	(7.0)	(15.3)	(20.8)	(20.1)	(15.0)	(9.3)	(5.5)	(5.0)	(100.0)				
SLE腎炎	6	28	82	107	82	41	22	7	11	4	390	36	426	19.96	3.43
(%)	(1.5)	(7.2)	(21.0)	(27.4)	(21.0)	(10.5)	(5.6)	(1.8)	(2.8)	(1.0)	(100.0)				
アミロイド腎		2	18	20	16	6	2	1	1		66	8	74	19.58	2.83
(%)		(3.0)	(27.3)	(30.3)	(24.2)	(9.1)	(3.0)	(1.5)	(1.5)		(100.0)				
痛風腎	1	3	18	35	31	30	20	9	4	3	154	12	166	21.60	3.48
(%)	(0.6)	(1.9)	(11.7)	(22.7)	(20.1)	(19.5)	(13.0)	(5.8)	(2.6)	(1.9)	(100.0)				
先天性代謝異常による腎不全	1		11	16	8	5				5	46	3	49	20.46	4.62
(%)	(2.2)		(23.9)	(34.8)	(17.4)	(10.9)				(10.9)	(100.0)				
腎・尿路結核		2	4	9	7	1	2	2		1	28	3	31	20.74	3.91
(%)		(7.1)	(14.3)	(32.1)	(25.0)	(3.6)	(7.1)	(7.1)		(3.6)	(100.0)				
腎・尿路結石	2	3	8	9	26	10	9	5			72	7	79	20.82	3.70
(%)	(2.8)	(4.2)	(11.1)	(12.5)	(36.1)	(13.9)	(12.5)	(6.9)			(100.0)				
腎・尿路腫瘍	1	3	11	27	27	22	13	2	4	1	111	8	119	21.34	3.24
(%)	(0.9)	(2.7)	(9.9)	(24.3)	(24.3)	(19.8)	(11.7)	(1.8)	(3.6)	(0.9)	(100.0)				
閉塞性尿路障害	1	2	16	21	24	17	7	6	3	2	99	5	104	21.26	3.54
(%)	(1.0)	(2.0)	(16.2)	(21.2)	(24.2)	(17.2)	(7.1)	(6.1)	(3.0)	(2.0)	(100.0)				
骨髄腫			3	9	5	8	3	1			29	4	33	21.17	2.64
(%)			(10.3)	(31.0)	(17.2)	(27.6)	(10.3)	(3.4)			(100.0)				
腎形成不全		7	16	27	32	24	16	4	3	2	131	5	136	21.14	3.53
(%)		(5.3)	(12.2)	(20.6)	(24.4)	(18.3)	(12.2)	(3.1)	(2.3)	(1.5)	(100.0)				
不明	17	148	502	894	951	743	424	253	135	113	4,180	523	4,703	21.51	3.70
(%)	(0.4)	(3.5)	(12.0)	(21.4)	(22.8)	(17.8)	(10.1)	(6.1)	(3.2)	(2.7)	(100.0)				
再導入	4	17	74	113	110	66	37	19	9	4	453	30	483	20.72	3.31
(%)	(0.9)	(3.8)	(16.3)	(24.9)	(24.3)	(14.6)	(8.2)	(4.2)	(2.0)	(0.9)	(100.0)				
その他	9	32	133	227	247	169	102	71	31	31	1,052	108	1,160	21.49	3.75
(%)	(0.9)	(3.0)	(12.6)	(21.6)	(23.5)	(16.1)	(9.7)	(6.7)	(2.9)	(2.9)	(100.0)				
合計	190	1,296	4,931	9,238	10,390	8,293	5,033	2,904	1,567	1,359	45,201	4,374	49,575	21.75	3.71
(%)	(0.4)	(2.9)	(10.9)	(20.4)	(23.0)	(18.3)	(11.1)	(6.4)	(3.5)	(3.0)	(100.0)				
記載なし				1		1	1		1		4	2	6	23.40	4.15
(%)				(25.0)		(25.0)	(25.0)		(25.0)		(100.0)				
総計	190	1,296	4,931	9,239	10,390	8,294	5,034	2,904	1,568	1,359	45,205	4,376	49,581	21.75	3.71
(%)	(0.4)	(2.9)	(10.9)	(20.4)	(23.0)	(18.3)	(11.1)	(6.4)	(3.5)	(3.0)	(100.0)				

数値下のかつこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査