

表 2440 透析前BUN(mg/dL) 原疾患別（施設血液透析患者 透析回数週3回 透析歴2年以上）

原疾患	20未満	20～	30～	40～	50～	60～	70～	80～	90～	100～	110～	120～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	102	598	2,453	7,014	12,335	13,966	9,803	4,812	1,702	504	152	54	53,495	1,292	54,787	62.81	15.44
(%)	(0.2)	(1.1)	(4.6)	(13.1)	(23.1)	(26.1)	(18.3)	(9.0)	(3.2)	(0.9)	(0.3)	(0.1)	(100.0)				
慢性腎盂腎炎	4	31	91	208	418	419	299	137	65	8	3		1,683	32	1,715	61.91	15.56
(%)	(0.2)	(1.8)	(5.4)	(12.4)	(24.8)	(24.9)	(17.8)	(8.1)	(3.9)	(0.5)	(0.2)		(100.0)				
急速進行性糸球体腎炎	1	20	67	179	303	261	225	121	34	14	1		1,226	31	1,257	61.96	15.72
(%)	(0.1)	(1.6)	(5.5)	(14.6)	(24.7)	(21.3)	(18.4)	(9.9)	(2.8)	(1.1)	(0.1)		(100.0)				
妊娠腎/妊娠中毒症		8	41	96	196	235	171	103	42	16	3	1	912	19	931	65.03	16.05
(%)		(0.9)	(4.5)	(10.5)	(21.5)	(25.8)	(18.8)	(11.3)	(4.6)	(1.8)	(0.3)	(0.1)	(100.0)				
その他の分類不能の腎炎	1	7	32	112	169	184	111	75	22	8	2		723	14	737	62.56	15.43
(%)	(0.1)	(1.0)	(4.4)	(15.5)	(23.4)	(25.4)	(15.4)	(10.4)	(3.0)	(1.1)	(0.3)		(100.0)				
多発性嚢胞腎	11	66	255	777	1,621	1,789	1,203	542	166	45	10	5	6,490	148	6,638	62.67	14.53
(%)	(0.2)	(1.0)	(3.9)	(12.0)	(25.0)	(27.6)	(18.5)	(8.4)	(2.6)	(0.7)	(0.2)	(0.1)	(100.0)				
腎硬化症	32	223	922	2,327	3,627	3,514	2,338	1,068	349	106	29	14	14,549	333	14,882	60.45	15.50
(%)	(0.2)	(1.5)	(6.3)	(16.0)	(24.9)	(24.2)	(16.1)	(7.3)	(2.4)	(0.7)	(0.2)	(0.1)	(100.0)				
悪性高血圧	3	13	78	200	339	364	231	119	54	10	2	1	1,414	37	1,451	62.06	15.26
(%)	(0.2)	(0.9)	(5.5)	(14.1)	(24.0)	(25.7)	(16.3)	(8.4)	(3.8)	(0.7)	(0.1)	(0.1)	(100.0)				
糖尿病性腎症	185	1,198	4,805	11,577	16,481	15,005	9,015	3,952	1,361	403	130	58	64,170	1,722	65,892	58.87	15.48
(%)	(0.3)	(1.9)	(7.5)	(18.0)	(25.7)	(23.4)	(14.0)	(6.2)	(2.1)	(0.6)	(0.2)	(0.1)	(100.0)				
SLE腎炎	3	17	57	130	251	283	220	114	39	18	3	1	1,136	32	1,168	63.52	16.02
(%)	(0.3)	(1.5)	(5.0)	(11.4)	(22.1)	(24.9)	(19.4)	(10.0)	(3.4)	(1.6)	(0.3)	(0.1)	(100.0)				
アミロイド腎	1	7	12	31	49	54	32	16	6	1	2		211	5	216	60.61	16.70
(%)	(0.5)	(3.3)	(5.7)	(14.7)	(23.2)	(25.6)	(15.2)	(7.6)	(2.8)	(0.5)	(0.9)		(100.0)				
痛風腎	1	9	28	87	150	158	97	64	31	11	2	1	639	13	652	63.46	16.60
(%)	(0.2)	(1.4)	(4.4)	(13.6)	(23.5)	(24.7)	(15.2)	(10.0)	(4.9)	(1.7)	(0.3)	(0.2)	(100.0)				
先天性代謝異常による腎不全	1	4	6	20	29	33	27	17	5	6	1		149	4	153	64.00	18.80
(%)	(0.7)	(2.7)	(4.0)	(13.4)	(19.5)	(22.1)	(18.1)	(11.4)	(3.4)	(4.0)	(0.7)		(100.0)				
腎・尿路結核		5	5	15	27	30	23	11	4	1			121	2	123	61.95	15.92
(%)		(4.1)	(4.1)	(12.4)	(22.3)	(24.8)	(19.0)	(9.1)	(3.3)	(0.8)			(100.0)				
腎・尿路結石	1	3	16	46	81	86	53	15	8	4			313	6	319	60.66	14.98
(%)	(0.3)	(1.0)	(5.1)	(14.7)	(25.9)	(27.5)	(16.9)	(4.8)	(2.6)	(1.3)			(100.0)				
腎・尿路腫瘍	1	3	23	64	111	128	72	36	9	4			451	8	459	61.23	14.49
(%)	(0.2)	(0.7)	(5.1)	(14.2)	(24.6)	(28.4)	(16.0)	(8.0)	(2.0)	(0.9)			(100.0)				
閉塞性尿路障害	2	5	22	49	87	88	59	34	18	5		1	370	7	377	62.74	16.85
(%)	(0.5)	(1.4)	(5.9)	(13.2)	(23.5)	(23.8)	(15.9)	(9.2)	(4.9)	(1.4)		(0.3)	(100.0)				
骨髄腫		4	12	8	20	23	13	3	3				86	1	87	57.25	16.86
(%)		(4.7)	(14.0)	(9.3)	(23.3)	(26.7)	(15.1)	(3.5)	(3.5)				(100.0)				
腎形成不全	1	1	10	23	71	98	70	42	14	5	3	1	339	5	344	67.12	15.39
(%)	(0.3)	(0.3)	(2.9)	(6.8)	(20.9)	(28.9)	(20.6)	(12.4)	(4.1)	(1.5)	(0.9)	(0.3)	(100.0)				
不明	47	294	920	2,306	3,559	3,559	2,328	1,160	426	132	40	22	14,793	500	15,293	60.73	16.12
(%)	(0.3)	(2.0)	(6.2)	(15.6)	(24.1)	(24.1)	(15.7)	(7.8)	(2.9)	(0.9)	(0.3)	(0.1)	(100.0)				
再導入	3	12	44	98	220	290	251	117	37	12	3	2	1,089	19	1,108	65.10	15.39
(%)	(0.3)	(1.1)	(4.0)	(9.0)	(20.2)	(26.6)	(23.0)	(10.7)	(3.4)	(1.1)	(0.3)	(0.2)	(100.0)				
その他	15	78	260	611	947	927	693	320	111	40	9	8	4,019	133	4,152	61.03	16.30
(%)	(0.4)	(1.9)	(6.5)	(15.2)	(23.6)	(23.1)	(17.2)	(8.0)	(2.8)	(1.0)	(0.2)	(0.2)	(100.0)				
合計	415	2,606	10,159	25,978	41,091	41,494	27,334	12,878	4,506	1,353	395	169	168,378	4,363	172,741	60.88	15.63
(%)	(0.2)	(1.5)	(6.0)	(15.4)	(24.4)	(24.6)	(16.2)	(7.6)	(2.7)	(0.8)	(0.2)	(0.1)	(100.0)				
記載なし				1		1		1					3		3	67.00	20.66
(%)				(33.3)		(33.3)		(33.3)					(100.0)				
総計	415	2,606	10,159	25,979	41,091	41,495	27,334	12,879	4,506	1,353	395	169	168,381	4,363	172,744	60.88	15.63
(%)	(0.2)	(1.5)	(6.0)	(15.4)	(24.4)	(24.6)	(16.2)	(7.6)	(2.7)	(0.8)	(0.2)	(0.1)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2015年調査