

表 2462 透析前BUN(mg/dL) 原疾患別（血液透析濾過患者 透析回数週3回 透析歴2年以上）

原疾患	20未満	20～	30～	40～	50～	60～	70～	80～	90～	100～	110～	120～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	14	95	578	1,864	3,718	4,245	3,178	1,529	530	134	45	18	15,948	378	16,326	63.92	14.87
(%)	(0.1)	(0.6)	(3.6)	(11.7)	(23.3)	(26.6)	(19.9)	(9.6)	(3.3)	(0.8)	(0.3)	(0.1)	(100.0)				
慢性腎盂腎炎	1	4	17	57	115	121	78	37	15	6			451	5	456	62.55	14.70
(%)	(0.2)	(0.9)	(3.8)	(12.6)	(25.5)	(26.8)	(17.3)	(8.2)	(3.3)	(1.3)			(100.0)				
急速進行性糸球体腎炎		4	12	25	71	73	70	30	8	2			295		295	64.17	14.63
(%)		(1.4)	(4.1)	(8.5)	(24.1)	(24.7)	(23.7)	(10.2)	(2.7)	(0.7)			(100.0)				
妊娠腎/妊娠中毒症	1	3	13	51	70	115	67	39	10	3	2	2	376	13	389	64.08	15.62
(%)	(0.3)	(0.8)	(3.5)	(13.6)	(18.6)	(30.6)	(17.8)	(10.4)	(2.7)	(0.8)	(0.5)	(0.5)	(100.0)				
その他の分類不能の腎炎		4	13	37	57	47	46	20	10	3	1		238	4	242	62.03	16.35
(%)		(1.7)	(5.5)	(15.5)	(23.9)	(19.7)	(19.3)	(8.4)	(4.2)	(1.3)	(0.4)		(100.0)				
多発性嚢胞腎		5	53	207	388	453	343	145	56	19	5	1	1,675	33	1,708	64.04	14.49
(%)		(0.3)	(3.2)	(12.4)	(23.2)	(27.0)	(20.5)	(8.7)	(3.3)	(1.1)	(0.3)	(0.1)	(100.0)				
腎硬化症	7	23	156	424	710	760	515	196	84	28	6	2	2,911	77	2,988	61.60	15.17
(%)	(0.2)	(0.8)	(5.4)	(14.6)	(24.4)	(26.1)	(17.7)	(6.7)	(2.9)	(1.0)	(0.2)	(0.1)	(100.0)				
悪性高血圧	1	3	10	36	84	94	53	28	15	3	1		328	7	335	63.42	15.02
(%)	(0.3)	(0.9)	(3.0)	(11.0)	(25.6)	(28.7)	(16.2)	(8.5)	(4.6)	(0.9)	(0.3)		(100.0)				
糖尿病性腎症	24	195	855	2,358	3,628	3,288	1,993	842	321	88	28	11	13,631	344	13,975	59.64	15.10
(%)	(0.2)	(1.4)	(6.3)	(17.3)	(26.6)	(24.1)	(14.6)	(6.2)	(2.4)	(0.6)	(0.2)	(0.1)	(100.0)				
SLE腎炎		3	14	41	86	90	66	47	12	6	1	1	367	11	378	64.87	15.86
(%)		(0.8)	(3.8)	(11.2)	(23.4)	(24.5)	(18.0)	(12.8)	(3.3)	(1.6)	(0.3)	(0.3)	(100.0)				
アミロイド腎	2	2	9	7	15	6	4	2					47	2	49	60.62	15.76
(%)		(4.3)	(4.3)	(19.1)	(14.9)	(31.9)	(12.8)	(8.5)	(4.3)				(100.0)				
痛風腎	2	8	20	42	41	17	16	8	1		1		156	2	158	62.40	16.13
(%)		(1.3)	(5.1)	(12.8)	(26.9)	(26.3)	(10.9)	(10.3)	(5.1)	(0.6)	(0.6)		(100.0)				
先天性代謝異常による腎不全		2	4	7	10	12	6	3	1				45	1	46	67.96	16.04
(%)			(4.4)	(8.9)	(15.6)	(22.2)	(26.7)	(13.3)	(6.7)	(2.2)			(100.0)				
腎・尿路結核		1	4	5	8	6	2			1			27	1	28	63.52	14.92
(%)			(3.7)	(14.8)	(18.5)	(29.6)	(22.2)	(7.4)		(3.7)			(100.0)				
腎・尿路結石		2	6	18	14	20	5	3	2				70		70	66.40	15.01
(%)			(2.9)	(8.6)	(25.7)	(20.0)	(28.6)	(7.1)	(4.3)	(2.9)			(100.0)				
腎・尿路腫瘍		3	10	24	24	16	8	3					88	2	90	62.76	13.86
(%)			(3.4)	(11.4)	(27.3)	(27.3)	(18.2)	(9.1)	(3.4)				(100.0)				
閉塞性尿路障害		6	17	19	24	19	8	2	1				96		96	61.97	14.72
(%)			(6.3)	(17.7)	(19.8)	(25.0)	(19.8)	(8.3)	(2.1)	(1.0)			(100.0)				
骨髄腫		2	2	6	6	3	3	1					23	2	25	64.13	15.82
(%)			(8.7)	(8.7)	(26.1)	(26.1)	(13.0)	(13.0)	(4.3)				(100.0)				
腎形成不全		3	17	28	29	20	15	9	3				124	2	126	65.96	16.12
(%)			(2.4)	(13.7)	(22.6)	(23.4)	(16.1)	(12.1)	(7.3)	(2.4)			(100.0)				
不明	9	39	203	529	925	968	641	322	127	36	7	4	3,810	131	3,941	62.11	15.58
(%)	(0.2)	(1.0)	(5.3)	(13.9)	(24.3)	(25.4)	(16.8)	(8.5)	(3.3)	(0.9)	(0.2)	(0.1)	(100.0)				
再導入		1	10	39	92	115	97	42	17	7	5	1	426	10	436	66.59	15.71
(%)			(0.2)	(2.3)	(9.2)	(21.6)	(27.0)	(22.8)	(9.9)	(4.0)	(1.6)	(1.2)	(100.0)				
その他		8	38	125	218	269	161	80	33	7	1		940	19	959	62.67	14.80
(%)			(0.9)	(4.0)	(13.3)	(23.2)	(28.6)	(17.1)	(8.5)	(3.5)	(0.7)	(0.1)	(100.0)				
合計	57	391	2,001	5,882	10,318	10,809	7,427	3,424	1,269	351	103	40	42,072	1,044	43,116	62.19	15.18
(%)	(0.1)	(0.9)	(4.8)	(14.0)	(24.5)	(25.7)	(17.7)	(8.1)	(3.0)	(0.8)	(0.2)	(0.1)	(100.0)				
記載なし					2	1	2	1					6		6	69.50	11.93
(%)					(33.3)	(16.7)	(33.3)	(16.7)					(100.0)				
総計	57	391	2,001	5,882	10,320	10,810	7,429	3,425	1,269	351	103	40	42,078	1,044	43,122	62.19	15.18
(%)	(0.1)	(0.9)	(4.8)	(14.0)	(24.5)	(25.7)	(17.7)	(8.1)	(3.0)	(0.8)	(0.2)	(0.1)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2015年調査