

表 2463 透析前BUN(mg/dL) 医療機関別（血液透析濾過患者 透析回数週3回 透析歴2年以上）

医療機関	20未満	20～	30～	40～	50～	60～	70～	80～	90～	100～	110～	120～	合計	記載なし	総計	平均	標準偏差
国公立大学			5	8	6	5	8	3					35	2	37	58.27	16.73
(%)			(14.3)	(22.9)	(17.1)	(14.3)	(22.9)	(8.6)					(100.0)				
私立大学	1	5	12	19	39	34	27	8	7	1			153	2	155	59.32	16.36
(%)	(0.7)	(3.3)	(7.8)	(12.4)	(25.5)	(22.2)	(17.6)	(5.2)	(4.6)	(0.7)			(100.0)				
国立		6	2	9	9	3	6		4				39		39	53.74	20.02
(%)		(15.4)	(5.1)	(23.1)	(23.1)	(7.7)	(15.4)		(10.3)				(100.0)				
県市町村立	7	19	98	284	429	440	280	143	51	13	4	1	1,769	48	1,817	61.17	15.52
(%)	(0.4)	(1.1)	(5.5)	(16.1)	(24.3)	(24.9)	(15.8)	(8.1)	(2.9)	(0.7)	(0.2)	(0.1)	(100.0)				
社会保険		3	27	59	97	107	51	19	9	2			374	14	388	59.56	14.27
(%)		(0.8)	(7.2)	(15.8)	(25.9)	(28.6)	(13.6)	(5.1)	(2.4)	(0.5)			(100.0)				
厚生連		11	49	149	239	213	148	57	18	5	1		890	38	928	60.12	14.37
(%)		(1.2)	(5.5)	(16.7)	(26.9)	(23.9)	(16.6)	(6.4)	(2.0)	(0.6)	(0.1)		(100.0)				
その他公的	1	11	33	118	220	169	108	39	15	9			723	35	758	59.89	14.70
(%)	(0.1)	(1.5)	(4.6)	(16.3)	(30.4)	(23.4)	(14.9)	(5.4)	(2.1)	(1.2)			(100.0)				
私立総合	1	10	39	158	245	283	171	82	31	9	3	2	1,034	14	1,048	62.25	15.21
(%)	(0.1)	(1.0)	(3.8)	(15.3)	(23.7)	(27.4)	(16.5)	(7.9)	(3.0)	(0.9)	(0.3)	(0.2)	(100.0)				
私立	21	131	614	1,562	2,451	2,560	1,739	769	274	78	25	13	10,237	448	10,685	61.15	15.51
(%)	(0.2)	(1.3)	(6.0)	(15.3)	(23.9)	(25.0)	(17.0)	(7.5)	(2.7)	(0.8)	(0.2)	(0.1)	(100.0)				
私立診療所	26	195	1,122	3,516	6,585	6,996	4,891	2,305	860	234	70	24	26,824	443	27,267	62.86	15.01
(%)	(0.1)	(0.7)	(4.2)	(13.1)	(24.5)	(26.1)	(18.2)	(8.6)	(3.2)	(0.9)	(0.3)	(0.1)	(100.0)				
総計	57	391	2,001	5,882	10,320	10,810	7,429	3,425	1,269	351	103	40	42,078	1,044	43,122	62.19	15.18
(%)	(0.1)	(0.9)	(4.8)	(14.0)	(24.5)	(25.7)	(17.7)	(8.1)	(3.0)	(0.8)	(0.2)	(0.1)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2015年調査