

表 2483 BUN(mg/dL) 透析歴(年)別 (腹膜透析患者)

透析歴	20未満	20～	30～	40～	50～	60～	70～	80～	90～	100～	110～	120～	合計	記載なし	総計	平均	標準偏差
2年未満	6	76	297	533	611	481	253	98	34	14	11	4	2,418	1,262	3,680	55.37	16.08
(%)	(0.2)	(3.1)	(12.3)	(22.0)	(25.3)	(19.9)	(10.5)	(4.1)	(1.4)	(0.6)	(0.5)	(0.2)	(100.0)				
2年～	6	59	202	463	564	429	203	79	37	10	6	1	2,059	1,043	3,102	55.61	15.16
(%)	(0.3)	(2.9)	(9.8)	(22.5)	(27.4)	(20.8)	(9.9)	(3.8)	(1.8)	(0.5)	(0.3)	(0.0)	(100.0)				
5年～	1	21	100	266	296	201	111	33	11	5	4		1,049	471	1,520	55.31	14.46
(%)	(0.1)	(2.0)	(9.5)	(25.4)	(28.2)	(19.2)	(10.6)	(3.1)	(1.0)	(0.5)	(0.4)		(100.0)				
10年～	1	7	30	55	76	51	19	8	1	2			250	64	314	53.50	14.04
(%)	(0.4)	(2.8)	(12.0)	(22.0)	(30.4)	(20.4)	(7.6)	(3.2)	(0.4)	(0.8)			(100.0)				
15年～		1	10	20	18	13	2		1				65	22	87	51.30	12.22
(%)		(1.5)	(15.4)	(30.8)	(27.7)	(20.0)	(3.1)		(1.5)				(100.0)				
20年～		3	2	3	6	1		1					16	10	26	47.38	15.84
(%)		(18.8)	(12.5)	(18.8)	(37.5)	(6.3)		(6.3)					(100.0)				
25年～		1		3	2		1	1					8	4	12	51.88	20.53
(%)		(12.5)		(37.5)	(25.0)		(12.5)	(12.5)					(100.0)				
30年～				1		3							4	2	6	57.25	10.31
(%)				(25.0)		(75.0)							(100.0)				
35年～				2									2		2	43.00	4.24
(%)				(100.0)									(100.0)				
40年～			1										1	1	2	36.00	
(%)			(100.0)										(100.0)				
合計	14	168	642	1,346	1,573	1,179	589	220	84	31	21	5	5,872	2,879	8,751	55.29	15.37
(%)	(0.2)	(2.9)	(10.9)	(22.9)	(26.8)	(20.1)	(10.0)	(3.7)	(1.4)	(0.5)	(0.4)	(0.1)	(100.0)				
不明		1				1							2	16	18	48.50	28.99
(%)		(50.0)				(50.0)							(100.0)				
記載なし																	
(%)																	
総計	14	169	642	1,346	1,573	1,180	589	220	84	31	21	5	5,874	2,895	8,769	55.28	15.37
(%)	(0.2)	(2.9)	(10.9)	(22.9)	(26.8)	(20.1)	(10.0)	(3.7)	(1.4)	(0.5)	(0.4)	(0.1)	(100.0)				
平均	2.71	3.10	3.01	3.30	3.24	3.08	2.83	2.77	2.56	2.74	2.14	1.20	3.11	2.75	2.99		
標準偏差	2.61	4.25	3.86	3.88	3.62	3.55	3.05	3.50	2.67	3.27	2.50	1.64	3.64	3.36	3.55		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査