

表 2484 BUN(mg/dL) 原疾患別（腹膜透析患者）

原疾患	20未満	20～	30～	40～	50～	60～	70～	80～	90～	100～	110～	120～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	2	32	179	387	555	421	228	76	34	9	6		1,929	901	2,830	56.85	14.69
(%)	(0.1)	(1.7)	(9.3)	(20.1)	(28.8)	(21.8)	(11.8)	(3.9)	(1.8)	(0.5)	(0.3)		(100.0)				
慢性腎盂腎炎		6	11	22	13	12	9	2			1	1	77	20	97	52.75	18.87
(%)		(7.8)	(14.3)	(28.6)	(16.9)	(15.6)	(11.7)	(2.6)			(1.3)	(1.3)	(100.0)				
急速進行性糸球体腎炎	2	5	12	10	9	3			1		1		43	23	66	53.63	17.17
(%)	(4.7)	(11.6)	(27.9)	(23.3)	(20.9)	(7.0)			(2.3)		(2.3)		(100.0)				
妊娠腎/妊娠中毒症	1	2	10	7	4		1						25	2	27	51.36	11.76
(%)	(4.0)	(8.0)	(40.0)	(28.0)	(16.0)		(4.0)						(100.0)				
その他の分類不能の腎炎	1		16	16	9	4	3	2			1		52	26	78	58.47	16.00
(%)		(1.9)		(30.8)	(30.8)	(17.3)	(7.7)	(5.8)	(3.8)		(1.9)		(100.0)				
多発性嚢胞腎	1	2	10	37	44	31	28	10	3	1			167	96	263	58.33	15.21
(%)	(0.6)	(1.2)	(6.0)	(22.2)	(26.3)	(18.6)	(16.8)	(6.0)	(1.8)	(0.6)			(100.0)				
腎硬化症	1	26	81	168	210	143	63	35	8	5	3		743	352	1,095	54.89	15.40
(%)	(0.1)	(3.5)	(10.9)	(22.6)	(28.3)	(19.2)	(8.5)	(4.7)	(1.1)	(0.7)	(0.4)		(100.0)				
悪性高血圧			7	19	16	11	8	4			1		66	41	107	56.19	14.95
(%)			(10.6)	(28.8)	(24.2)	(16.7)	(12.1)	(6.1)			(1.5)		(100.0)				
糖尿病性腎症	8	74	223	463	457	368	154	47	27	8	4	3	1,836	963	2,799	53.68	15.44
(%)	(0.4)	(4.0)	(12.1)	(25.2)	(24.9)	(20.0)	(8.4)	(2.6)	(1.5)	(0.4)	(0.2)	(0.2)	(100.0)				
SLE腎炎			4	9	5	8	3	1	1				31	15	46	55.35	15.96
(%)			(12.9)	(29.0)	(16.1)	(25.8)	(9.7)	(3.2)	(3.2)				(100.0)				
アミロイド腎		1	1	7				2					11	4	15	57.09	13.60
(%)			(9.1)	(9.1)	(63.6)			(18.2)					(100.0)				
痛風腎		4	2	7	6	3	1	2					25	7	32	60.19	17.41
(%)			(16.0)	(8.0)	(28.0)	(24.0)	(12.0)	(4.0)	(8.0)				(100.0)				
先天性代謝異常による腎不全				3	2	2	4	1					12	5	17	64.58	14.43
(%)				(25.0)	(16.7)	(16.7)	(33.3)	(8.3)					(100.0)				
腎・尿路結核				1									1	1	2	48.00	
(%)				(100.0)									(100.0)				
腎・尿路結石		1	3	4	7	1	1						17	4	21	48.29	12.04
(%)		(5.9)	(17.6)	(23.5)	(41.2)	(5.9)	(5.9)						(100.0)				
腎・尿路腫瘍			1	4	4		1	1					11	3	14	52.19	13.40
(%)			(9.1)	(36.4)	(36.4)		(9.1)	(9.1)					(100.0)				
閉塞性尿路障害			2	3	4	4				1			14	7	21	55.00	16.88
(%)			(14.3)	(21.4)	(28.6)	(28.6)				(7.1)			(100.0)				
骨髄腫			3					1					4	3	7	47.75	25.71
(%)			(75.0)					(25.0)					(100.0)				
腎形成不全		2	4	8	19	8	4	3	1	1			50	22	72	57.17	15.78
(%)		(4.0)	(8.0)	(16.0)	(38.0)	(16.0)	(8.0)	(6.0)	(2.0)	(2.0)			(100.0)				
不明	1	16	73	129	139	101	61	24	4	5	2	1	556	286	842	55.06	15.97
(%)	(0.2)	(2.9)	(13.1)	(23.2)	(25.0)	(18.2)	(11.0)	(4.3)	(0.7)	(0.9)	(0.4)	(0.2)	(100.0)				
再導入			4	3	9	6	2	2			1		27	28	55	57.85	17.91
(%)			(14.8)	(11.1)	(33.3)	(22.2)	(7.4)	(7.4)			(3.7)		(100.0)				
その他	1	6	25	45	42	36	13	6	1		2		177	86	263	53.78	15.44
(%)	(0.6)	(3.4)	(14.1)	(25.4)	(23.7)	(20.3)	(7.3)	(3.4)	(0.6)		(1.1)		(100.0)				
合計	14	169	642	1,346	1,573	1,180	589	220	84	31	21	5	5,874	2,895	8,769	55.28	15.37
(%)	(0.2)	(2.9)	(10.9)	(22.9)	(26.8)	(20.1)	(10.0)	(3.7)	(1.4)	(0.5)	(0.4)	(0.1)	(100.0)				
記載なし																	
総計	14	169	642	1,346	1,573	1,180	589	220	84	31	21	5	5,874	2,895	8,769	55.28	15.37
(%)	(0.2)	(2.9)	(10.9)	(22.9)	(26.8)	(20.1)	(10.0)	(3.7)	(1.4)	(0.5)	(0.4)	(0.1)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2015年調査