

表 2485 BUN(mg/dL) 医療機関別（腹膜透析患者）

医療機関	20未満	20～	30～	40～	50～	60～	70～	80～	90～	100～	110～	120～	合計	記載なし	総計	平均	標準偏差
国公立大学		7	24	77	84	96	45	21	4	1	1		360	303	663	58.07	14.27
(%)		(1.9)	(6.7)	(21.4)	(23.3)	(26.7)	(12.5)	(5.8)	(1.1)	(0.3)	(0.3)		(100.0)				
私立大学		17	61	158	183	152	73	33	7	3	2	1	690	416	1,106	56.20	15.05
(%)		(2.5)	(8.8)	(22.9)	(26.5)	(22.0)	(10.6)	(4.8)	(1.0)	(0.4)	(0.3)	(0.1)	(100.0)				
国立		3	20	41	48	25	13	6	3				159	63	222	53.68	14.18
(%)		(1.9)	(12.6)	(25.8)	(30.2)	(15.7)	(8.2)	(3.8)	(1.9)				(100.0)				
県市町村立	1	27	146	250	307	214	99	39	14	4	5	1	1,107	466	1,573	54.76	15.06
(%)	(0.1)	(2.4)	(13.2)	(22.6)	(27.7)	(19.3)	(8.9)	(3.5)	(1.3)	(0.4)	(0.5)	(0.1)	(100.0)				
社会保険	2	3	22	41	43	50	19	4	2	2	1		189	333	522	55.51	15.32
(%)	(1.1)	(1.6)	(11.6)	(21.7)	(22.8)	(26.5)	(10.1)	(2.1)	(1.1)	(1.1)	(0.5)		(100.0)				
厚生連	2	14	70	107	103	77	36	9	7	3	1	1	430	130	560	53.20	16.16
(%)	(0.5)	(3.3)	(16.3)	(24.9)	(24.0)	(17.9)	(8.4)	(2.1)	(1.6)	(0.7)	(0.2)	(0.2)	(100.0)				
その他公的	1	24	101	239	262	175	98	34	11	4	2	1	952	407	1,359	54.87	14.79
(%)	(0.1)	(2.5)	(10.6)	(25.1)	(27.5)	(18.4)	(10.3)	(3.6)	(1.2)	(0.4)	(0.2)	(0.1)	(100.0)				
私立総合		13	28	61	88	68	45	11	5	1	1		321	42	363	56.24	15.33
(%)		(4.0)	(8.7)	(19.0)	(27.4)	(21.2)	(14.0)	(3.4)	(1.6)	(0.3)	(0.3)		(100.0)				
私立	6	46	115	250	290	213	96	46	21	12	7	1	1,103	596	1,699	55.26	16.76
(%)	(0.5)	(4.2)	(10.4)	(22.7)	(26.3)	(19.3)	(8.7)	(4.2)	(1.9)	(1.1)	(0.6)	(0.1)	(100.0)				
私立診療所	2	15	55	122	165	110	65	17	10	1	1		563	139	702	55.56	14.65
(%)	(0.4)	(2.7)	(9.8)	(21.7)	(29.3)	(19.5)	(11.5)	(3.0)	(1.8)	(0.2)	(0.2)		(100.0)				
総計	14	169	642	1,346	1,573	1,180	589	220	84	31	21	5	5,874	2,895	8,769	55.28	15.37
(%)	(0.2)	(2.9)	(10.9)	(22.9)	(26.8)	(20.1)	(10.0)	(3.7)	(1.4)	(0.5)	(0.4)	(0.1)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2015年調査