

表 2625 nPCR(g/kg/day) 原疾患別（施設血液透析患者 透析回数週3回 透析歴2年以上）

原疾患	0.4未満	0.4～	0.6～	0.8～	1.0～	1.2～	1.4～	1.6～	1.8～	2.0～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	71	1,847	12,825	22,753	10,883	2,062	230	37	4	18	50,730	4,057	54,787	0.89	0.18
(%)	(0.1)	(3.6)	(25.3)	(44.9)	(21.5)	(4.1)	(0.5)	(0.1)	(0.0)	(0.0)	(100.0)				
慢性腎盂腎炎	2	74	407	721	319	67	9				1,599	116	1,715	0.89	0.18
(%)	(0.1)	(4.6)	(25.5)	(45.1)	(19.9)	(4.2)	(0.6)				(100.0)				
急速進行性糸球体腎炎		49	322	486	270	43	5	1			1,176	81	1,257	0.89	0.18
(%)		(4.2)	(27.4)	(41.3)	(23.0)	(3.7)	(0.4)	(0.1)			(100.0)				
妊娠腎/妊娠中毒症		26	168	352	247	64	12			1	870	61	931	0.94	0.20
(%)		(3.0)	(19.3)	(40.5)	(28.4)	(7.4)	(1.4)			(0.1)	(100.0)				
その他の分類不能の腎炎	1	22	182	309	149	34	2				699	38	737	0.89	0.18
(%)	(0.1)	(3.1)	(26.0)	(44.2)	(21.3)	(4.9)	(0.3)				(100.0)				
多発性嚢胞腎	12	201	1,659	2,957	1,163	175	13	1		2	6,183	455	6,638	0.88	0.16
(%)	(0.2)	(3.3)	(26.8)	(47.8)	(18.8)	(2.8)	(0.2)	(0.0)		(0.0)	(100.0)				
腎硬化症	26	735	4,507	5,932	2,314	365	43	4		6	13,932	950	14,882	0.86	0.17
(%)	(0.2)	(5.3)	(32.3)	(42.6)	(16.6)	(2.6)	(0.3)	(0.0)		(0.0)	(100.0)				
悪性高血圧	3	58	385	604	245	50	7			1	1,353	98	1,451	0.88	0.18
(%)	(0.2)	(4.3)	(28.5)	(44.6)	(18.1)	(3.7)	(0.5)			(0.1)	(100.0)				
糖尿病性腎症	135	4,152	22,307	25,142	7,964	1,140	139	18	9	16	61,022	4,870	65,892	0.83	0.17
(%)	(0.2)	(6.8)	(36.6)	(41.2)	(13.1)	(1.9)	(0.2)	(0.0)	(0.0)	(0.0)	(100.0)				
SLE腎炎	1	41	236	446	289	59	10				1,082	86	1,168	0.92	0.18
(%)	(0.1)	(3.8)	(21.8)	(41.2)	(26.7)	(5.5)	(0.9)				(100.0)				
アミロイド腎	2	8	49	89	35	10					193	23	216	0.87	0.18
(%)	(1.0)	(4.1)	(25.4)	(46.1)	(18.1)	(5.2)					(100.0)				
痛風腎		20	179	257	112	35	2			1	606	46	652	0.89	0.20
(%)		(3.3)	(29.5)	(42.4)	(18.5)	(5.8)	(0.3)			(0.2)	(100.0)				
先天性代謝異常による腎不全	1	5	34	52	33	11	2				138	15	153	0.92	0.21
(%)	(0.7)	(3.6)	(24.6)	(37.7)	(23.9)	(8.0)	(1.4)				(100.0)				
腎・尿路結核		7	27	50	27	6					117	6	123	0.88	0.19
(%)		(6.0)	(23.1)	(42.7)	(23.1)	(5.1)					(100.0)				
腎・尿路結石	2	12	94	130	51	11	1				301	18	319	0.86	0.17
(%)	(0.7)	(4.0)	(31.2)	(43.2)	(16.9)	(3.7)	(0.3)				(100.0)				
腎・尿路腫瘍	1	16	135	203	65	13	1				434	25	459	0.86	0.16
(%)	(0.2)	(3.7)	(31.1)	(46.8)	(15.0)	(3.0)	(0.2)				(100.0)				
閉塞性尿路障害	1	22	92	141	74	17	4				351	26	377	0.89	0.20
(%)	(0.3)	(6.3)	(26.2)	(40.2)	(21.1)	(4.8)	(1.1)				(100.0)				
骨髄腫		12	20	30	15	1					78	9	87	0.82	0.19
(%)		(15.4)	(25.6)	(38.5)	(19.2)	(1.3)					(100.0)				
腎形成不全		7	55	137	100	18	4	1			322	22	344	0.94	0.18
(%)		(2.2)	(17.1)	(42.5)	(31.1)	(5.6)	(1.2)	(0.3)			(100.0)				
不明	36	814	4,323	5,752	2,569	474	53	6	2	6	14,035	1,258	15,293	0.86	0.18
(%)	(0.3)	(5.8)	(30.8)	(41.0)	(18.3)	(3.4)	(0.4)	(0.0)	(0.0)	(0.0)	(100.0)				
再導入	2	36	197	452	281	52	1	1			1,022	86	1,108	0.92	0.18
(%)	(0.2)	(3.5)	(19.3)	(44.2)	(27.5)	(5.1)	(0.1)	(0.1)			(100.0)				
その他	13	236	1,135	1,538	747	141	21	4			3,835	317	4,152	0.87	0.19
(%)	(0.3)	(6.2)	(29.6)	(40.1)	(19.5)	(3.7)	(0.5)	(0.1)			(100.0)				
合計	309	8,400	49,338	68,533	27,952	4,848	559	73	15	51	160,078	12,663	172,741	0.86	0.18
(%)	(0.2)	(5.2)	(30.8)	(42.8)	(17.5)	(3.0)	(0.3)	(0.0)	(0.0)	(0.0)	(100.0)				
記載なし			1	1		1					3		3	0.96	0.35
(%)			(33.3)	(33.3)		(33.3)					(100.0)				
総計	309	8,400	49,339	68,534	27,952	4,849	559	73	15	51	160,081	12,663	172,744	0.86	0.18
(%)	(0.2)	(5.2)	(30.8)	(42.8)	(17.5)	(3.0)	(0.3)	(0.0)	(0.0)	(0.0)	(100.0)				

数値下のカッコ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査