

表 2665 %クレアチニン産生速度(%) 年齢(歳)別(施設血液透析患者 透析回数週3回 透析歴2年以上)

年齢	60未満	60～	70～	80～	90～	100～	110～	120～	130～	140～	合計	記載なし	総計	平均	標準偏差
15歳未満	1	2									3	1	4	57.43	8.55
(%)	(33.3)	(66.7)									(100.0)				
15歳～	16	13	39	40	58	67	59	40	7	4	343	25	368	97.95	22.04
(%)	(4.7)	(3.8)	(11.4)	(11.7)	(16.9)	(19.5)	(17.2)	(11.7)	(2.0)	(1.2)	(100.0)				
30歳～	292	236	413	712	1,015	1,253	1,135	754	346	158	6,314	446	6,760	101.76	22.85
(%)	(4.6)	(3.7)	(6.5)	(11.3)	(16.1)	(19.8)	(18.0)	(11.9)	(5.5)	(2.5)	(100.0)				
45歳～	1,470	1,220	2,226	3,386	4,436	4,868	4,423	3,116	1,527	826	27,498	2,080	29,578	100.35	23.58
(%)	(5.3)	(4.4)	(8.1)	(12.3)	(16.1)	(17.7)	(16.1)	(11.3)	(5.6)	(3.0)	(100.0)				
60歳～	5,295	3,946	6,225	8,916	11,102	12,139	11,073	7,971	4,585	3,101	74,353	5,814	80,167	99.53	25.69
(%)	(7.1)	(5.3)	(8.4)	(12.0)	(14.9)	(16.3)	(14.9)	(10.7)	(6.2)	(4.2)	(100.0)				
75歳～	5,260	3,211	4,586	6,058	7,210	7,258	6,155	4,316	2,500	1,988	48,542	4,111	52,653	95.42	27.75
(%)	(10.8)	(6.6)	(9.4)	(12.5)	(14.9)	(15.0)	(12.7)	(8.9)	(5.2)	(4.1)	(100.0)				
90歳～	486	255	312	383	386	384	293	197	112	108	2,916	297	3,213	89.34	29.90
(%)	(16.7)	(8.7)	(10.7)	(13.1)	(13.2)	(13.2)	(10.0)	(6.8)	(3.8)	(3.7)	(100.0)				
合計	12,820	8,883	13,801	19,495	24,207	25,969	23,138	16,394	9,077	6,185	159,969	12,774	172,743	98.32	26.07
(%)	(8.0)	(5.6)	(8.6)	(12.2)	(15.1)	(16.2)	(14.5)	(10.2)	(5.7)	(3.9)	(100.0)				
不明												1	1		
(%)															
記載なし															
(%)															
総計	12,820	8,883	13,801	19,495	24,207	25,969	23,138	16,394	9,077	6,185	159,969	12,775	172,744	98.32	26.07
(%)	(8.0)	(5.6)	(8.6)	(12.2)	(15.1)	(16.2)	(14.5)	(10.2)	(5.7)	(3.9)	(100.0)				
平均	71.75	70.29	69.16	68.43	67.79	67.21	66.93	66.97	67.81	69.45	68.21	68.97	68.26		
標準偏差	11.75	11.87	12.00	12.08	12.24	12.32	12.15	11.97	11.41	10.97	12.10	12.17	12.11		

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2015年調査