

表 2667 %クレアチニン産生速度(%) 原疾患別（施設血液透析患者 透析回数週3回 透析歴2年以上）

原疾患	60未満	60～	70～	80～	90～	100～	110～	120～	130～	140～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	2,489	1,753	3,019	4,864	6,752	8,548	8,868	7,137	4,310	2,953	50,693	4,094	54,787	105.32	24.81
(%)	(4.9)	(3.5)	(6.0)	(9.6)	(13.3)	(16.9)	(17.5)	(14.1)	(8.5)	(5.8)	(100.0)				
慢性腎盂腎炎	102	77	123	186	232	250	261	198	105	65	1,599	116	1,715	101.00	25.40
(%)	(6.4)	(4.8)	(7.7)	(11.6)	(14.5)	(15.6)	(16.3)	(12.4)	(6.6)	(4.1)	(100.0)				
急速進行性糸球体腎炎	156	60	86	124	161	148	154	163	77	46	1,175	82	1,257	96.30	30.66
(%)	(13.3)	(5.1)	(7.3)	(10.6)	(13.7)	(12.6)	(13.1)	(13.9)	(6.6)	(3.9)	(100.0)				
妊娠腎/妊娠中毒症	32	30	42	85	121	158	149	118	88	46	869	62	931	106.43	22.95
(%)	(3.7)	(3.5)	(4.8)	(9.8)	(13.9)	(18.2)	(17.1)	(13.6)	(10.1)	(5.3)	(100.0)				
その他の分類不能の腎炎	48	42	35	68	92	115	122	86	49	42	699	38	737	102.69	27.05
(%)	(6.9)	(6.0)	(5.0)	(9.7)	(13.2)	(16.5)	(17.5)	(12.3)	(7.0)	(6.0)	(100.0)				
多発性嚢胞腎	226	239	441	708	1,054	1,225	1,088	646	367	182	6,176	462	6,638	102.08	21.88
(%)	(3.7)	(3.9)	(7.1)	(11.5)	(17.1)	(19.8)	(17.6)	(10.5)	(5.9)	(2.9)	(100.0)				
腎硬化症	1,053	700	1,095	1,565	2,070	2,248	2,031	1,522	922	719	13,925	957	14,882	100.24	26.70
(%)	(7.6)	(5.0)	(7.9)	(11.2)	(14.9)	(16.1)	(14.6)	(10.9)	(6.6)	(5.2)	(100.0)				
悪性高血圧	103	58	91	134	206	200	229	167	102	62	1,352	99	1,451	101.41	26.87
(%)	(7.6)	(4.3)	(6.7)	(9.9)	(15.2)	(14.8)	(16.9)	(12.4)	(7.5)	(4.6)	(100.0)				
糖尿病性腎症	6,496	4,688	7,027	9,284	10,354	9,457	6,927	3,945	1,712	1,096	60,986	4,906	65,892	91.45	24.90
(%)	(10.7)	(7.7)	(11.5)	(15.2)	(17.0)	(15.5)	(11.4)	(6.5)	(2.8)	(1.8)	(100.0)				
SLE腎炎	106	67	109	114	177	178	161	100	49	19	1,080	88	1,168	95.11	26.52
(%)	(9.8)	(6.2)	(10.1)	(10.6)	(16.4)	(16.5)	(14.9)	(9.3)	(4.5)	(1.8)	(100.0)				
アミロイド腎	31	23	21	20	36	27	21	6	4	4	193	23	216	86.58	29.26
(%)	(16.1)	(11.9)	(10.9)	(10.4)	(18.7)	(14.0)	(10.9)	(3.1)	(2.1)	(2.1)	(100.0)				
痛風腎	22	19	37	60	84	108	101	84	53	37	605	47	652	106.27	23.88
(%)	(3.6)	(3.1)	(6.1)	(9.9)	(13.9)	(17.9)	(16.7)	(13.9)	(8.8)	(6.1)	(100.0)				
先天性代謝異常による腎不全	11	8	16	16	25	20	22	11	4	5	138	15	153	96.26	25.62
(%)	(8.0)	(5.8)	(11.6)	(11.6)	(18.1)	(14.5)	(15.9)	(8.0)	(2.9)	(3.6)	(100.0)				
腎・尿路結核	13	8	8	9	23	21	18	9	3	5	117	6	123	95.14	26.33
(%)	(11.1)	(6.8)	(6.8)	(7.7)	(19.7)	(17.9)	(15.4)	(7.7)	(2.6)	(4.3)	(100.0)				
腎・尿路結石	14	17	16	32	39	51	54	40	20	18	301	18	319	104.19	25.32
(%)	(4.7)	(5.6)	(5.3)	(10.6)	(13.0)	(16.9)	(17.9)	(13.3)	(6.6)	(6.0)	(100.0)				
腎・尿路腫瘍	28	14	27	48	36	74	68	70	28	41	434	25	459	106.07	27.50
(%)	(6.5)	(3.2)	(6.2)	(11.1)	(8.3)	(17.1)	(15.7)	(16.1)	(6.5)	(9.4)	(100.0)				
閉塞性尿路障害	49	29	28	37	46	54	48	33	19	8	351	26	377	92.45	29.46
(%)	(14.0)	(8.3)	(8.0)	(10.5)	(13.1)	(15.4)	(13.7)	(9.4)	(5.4)	(2.3)	(100.0)				
骨髄腫	11	9	9	12	11	8	7	6	5		78	9	87	87.27	27.64
(%)	(14.1)	(11.5)	(11.5)	(15.4)	(14.1)	(10.3)	(9.0)	(7.7)	(6.4)		(100.0)				
腎形成不全	11	13	10	34	53	62	64	41	24	10	322	22	344	104.97	21.32
(%)	(3.4)	(4.0)	(3.1)	(10.6)	(16.5)	(19.3)	(19.9)	(12.7)	(7.5)	(3.1)	(100.0)				
不明	1,303	752	1,185	1,530	1,957	2,240	2,024	1,491	891	650	14,023	1,270	15,293	98.57	27.66
(%)	(9.3)	(5.4)	(8.5)	(10.9)	(14.0)	(16.0)	(14.4)	(10.6)	(6.4)	(4.6)	(100.0)				
再導入	67	39	58	116	155	169	187	140	60	31	1,022	86	1,108	101.37	24.97
(%)	(6.6)	(3.8)	(5.7)	(11.4)	(15.2)	(16.5)	(18.3)	(13.7)	(5.9)	(3.0)	(100.0)				
その他	449	237	318	447	523	608	534	381	185	146	3,828	324	4,152	95.38	28.62
(%)	(11.7)	(6.2)	(8.3)	(11.7)	(13.7)	(15.9)	(13.9)	(10.0)	(4.8)	(3.8)	(100.0)				
合計	12,820	8,882	13,801	19,493	24,207	25,969	23,138	16,394	9,077	6,185	159,966	12,775	172,741	98.32	26.07
(%)	(8.0)	(5.6)	(8.6)	(12.2)	(15.1)	(16.2)	(14.5)	(10.2)	(5.7)	(3.9)	(100.0)				
記載なし		1		2							3		3	77.97	12.82
(%)		(33.3)		(66.7)							(100.0)				
総計	12,820	8,883	13,801	19,495	24,207	25,969	23,138	16,394	9,077	6,185	159,969	12,775	172,744	98.32	26.07
(%)	(8.0)	(5.6)	(8.6)	(12.2)	(15.1)	(16.2)	(14.5)	(10.2)	(5.7)	(3.9)	(100.0)				

数値下のカッコ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査