

表 2687 %クレアチニン産生速度(%) 原疾患別（血液透析濾過患者 透析回数週3回 透析歴2年以上）

原疾患	60未満	60～	70～	80～	90～	100～	110～	120～	130～	140～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	459	491	784	1,421	2,154	2,830	2,826	2,282	1,323	884	15,454	872	16,326	107.10	22.78
(%)	(3.0)	(3.2)	(5.1)	(9.2)	(13.9)	(18.3)	(18.3)	(14.8)	(8.6)	(5.7)	(100.0)				
慢性腎盂腎炎	12	19	38	47	75	75	78	52	25	18	439	17	456	102.80	22.05
(%)	(2.7)	(4.3)	(8.7)	(10.7)	(17.1)	(17.1)	(17.8)	(11.8)	(5.7)	(4.1)	(100.0)				
急速進行性糸球体腎炎	18	10	11	26	43	49	60	27	21	17	282	13	295	103.30	26.02
(%)	(6.4)	(3.5)	(3.9)	(9.2)	(15.2)	(17.4)	(21.3)	(9.6)	(7.4)	(6.0)	(100.0)				
妊娠腎/妊娠中毒症	7	13	18	39	47	70	60	65	29	16	364	25	389	107.10	20.94
(%)	(1.9)	(3.6)	(4.9)	(10.7)	(12.9)	(19.2)	(16.5)	(17.9)	(8.0)	(4.4)	(100.0)				
その他の分類不能の腎炎	11	12	11	19	23	51	53	32	9	7	228	14	242	103.36	22.95
(%)	(4.8)	(5.3)	(4.8)	(8.3)	(10.1)	(22.4)	(23.2)	(14.0)	(3.9)	(3.1)	(100.0)				
多発性嚢胞腎	40	52	115	188	281	319	290	192	90	49	1,616	92	1,708	103.43	21.81
(%)	(2.5)	(3.2)	(7.1)	(11.6)	(17.4)	(19.7)	(17.9)	(11.9)	(5.6)	(3.0)	(100.0)				
腎硬化症	141	119	180	326	416	468	450	351	208	167	2,826	162	2,988	103.62	25.25
(%)	(5.0)	(4.2)	(6.4)	(11.5)	(14.7)	(16.6)	(15.9)	(12.4)	(7.4)	(5.9)	(100.0)				
悪性高血圧	10	11	26	33	45	56	59	46	20	14	320	15	335	104.62	24.01
(%)	(3.1)	(3.4)	(8.1)	(10.3)	(14.1)	(17.5)	(18.4)	(14.4)	(6.3)	(4.4)	(100.0)				
糖尿病性腎症	996	907	1,440	2,068	2,317	2,179	1,616	894	432	269	13,118	857	13,975	94.07	23.60
(%)	(7.6)	(6.9)	(11.0)	(15.8)	(17.7)	(16.6)	(12.3)	(6.8)	(3.3)	(2.1)	(100.0)				
SLE腎炎	34	24	26	48	52	70	52	29	15	7	357	21	378	95.55	24.40
(%)	(9.5)	(6.7)	(7.3)	(13.4)	(14.6)	(19.6)	(14.6)	(8.1)	(4.2)	(2.0)	(100.0)				
アミロイド腎	7	3	4	6	7	9	4	2	2	1	45	4	49	90.38	27.77
(%)	(15.6)	(6.7)	(8.9)	(13.3)	(15.6)	(20.0)	(8.9)	(4.4)	(4.4)	(2.2)	(100.0)				
痛風腎	3	5	12	18	21	23	17	31	11	12	153	5	158	106.72	23.62
(%)	(2.0)	(3.3)	(7.8)	(11.8)	(13.7)	(15.0)	(11.1)	(20.3)	(7.2)	(7.8)	(100.0)				
先天性代謝異常による腎不全	2		4	4	13	5	7	6		2	43	3	46	101.23	21.02
(%)	(4.7)		(9.3)	(9.3)	(30.2)	(11.6)	(16.3)	(14.0)		(4.7)	(100.0)				
腎・尿路結核	1	1	2	4	6	5	3	3	1	1	27	1	28	99.57	20.31
(%)	(3.7)	(3.7)	(7.4)	(14.8)	(22.2)	(18.5)	(11.1)	(11.1)	(3.7)	(3.7)	(100.0)				
腎・尿路結石	2	5	4	4	10	11	15	7	6	4	68	2	70	105.10	22.02
(%)	(2.9)	(7.4)	(5.9)	(5.9)	(14.7)	(16.2)	(22.1)	(10.3)	(8.8)	(5.9)	(100.0)				
腎・尿路腫瘍	2	4	7	6	11	15	8	14	11	7	85	5	90	108.60	27.06
(%)	(2.4)	(4.7)	(8.2)	(7.1)	(12.9)	(17.6)	(9.4)	(16.5)	(12.9)	(8.2)	(100.0)				
閉塞性尿路障害	5	2	9	14	14	15	13	10	6	3	91	5	96	99.61	24.07
(%)	(5.5)	(2.2)	(9.9)	(15.4)	(15.4)	(16.5)	(14.3)	(11.0)	(6.6)	(3.3)	(100.0)				
骨髄腫	2	2	4	2		8	2	1		1	22	3	25	90.14	31.28
(%)	(9.1)	(9.1)	(18.2)	(9.1)		(36.4)	(9.1)	(4.5)		(4.5)	(100.0)				
腎形成不全	3	2	4	15	13	37	21	12	10	3	120	6	126	105.31	19.31
(%)	(2.5)	(1.7)	(3.3)	(12.5)	(10.8)	(30.8)	(17.5)	(10.0)	(8.3)	(2.5)	(100.0)				
不明	193	155	247	385	560	641	614	451	231	198	3,675	266	3,941	102.97	24.77
(%)	(5.3)	(4.2)	(6.7)	(10.5)	(15.2)	(17.4)	(16.7)	(12.3)	(6.3)	(5.4)	(100.0)				
再導入	14	14	22	49	70	68	64	65	34	12	412	24	436	104.79	22.26
(%)	(3.4)	(3.4)	(5.3)	(11.9)	(17.0)	(16.5)	(15.5)	(15.8)	(8.3)	(2.9)	(100.0)				
その他	57	39	66	103	140	156	148	101	56	42	908	51	959	101.19	25.03
(%)	(6.3)	(4.3)	(7.3)	(11.3)	(15.4)	(17.2)	(16.3)	(11.1)	(6.2)	(4.6)	(100.0)				
合計	2,019	1,890	3,034	4,825	6,318	7,160	6,460	4,673	2,540	1,734	40,653	2,463	43,116	101.70	24.12
(%)	(5.0)	(4.6)	(7.5)	(11.9)	(15.5)	(17.6)	(15.9)	(11.5)	(6.2)	(4.3)	(100.0)				
記載なし				1		1	2	2			6		6	114.67	14.50
(%)				(16.7)		(16.7)	(33.3)	(33.3)			(100.0)				
総計	2,019	1,890	3,034	4,826	6,318	7,161	6,462	4,675	2,540	1,734	40,659	2,463	43,122	101.70	24.12
(%)	(5.0)	(4.6)	(7.5)	(11.9)	(15.5)	(17.6)	(15.9)	(11.5)	(6.2)	(4.3)	(100.0)				

数値下のカッコ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査