

表 2726 透析前アルブミン濃度(g/dL) 治療方法(新分類)別(透析患者全体)

治療方法 (新分類)	1.5未満	1.5～	2.0～	2.5～	3.0～	3.5～	4.0～	4.5～	5.0～	合計	記載なし	総計	平均	標準偏差
施設血液透析	177	918	3,852	13,605	56,978	108,967	37,192	1,915	109	223,713	25,012	248,725	3.57	0.45
(%)	(0.1)	(0.4)	(1.7)	(6.1)	(25.5)	(48.7)	(16.6)	(0.9)	(0.0)	(100.0)				
Off-line HDF	2	23	87	322	1,399	2,278	642	27	3	4,783	549	5,332	3.52	0.44
(%)	(0.0)	(0.5)	(1.8)	(6.7)	(29.2)	(47.6)	(13.4)	(0.6)	(0.1)	(100.0)				
On-line HDF	5	51	281	1,516	10,207	22,210	6,334	233	13	40,850	3,677	44,527	3.61	0.37
(%)	(0.0)	(0.1)	(0.7)	(3.7)	(25.0)	(54.4)	(15.5)	(0.6)	(0.0)	(100.0)				
Push/Pull HDF			1	3	27	53	19	4		107	3	110	3.66	0.40
(%)			(0.9)	(2.8)	(25.2)	(49.5)	(17.8)	(3.7)		(100.0)				
ハイオフィルトレーション		1	3	23	83	107	33	2		252	15	267	3.49	0.42
(%)		(0.4)	(1.2)	(9.1)	(32.9)	(42.5)	(13.1)	(0.8)		(100.0)				
IHDF		8	26	136	863	1,790	582	26	1	3,432	108	3,540	3.61	0.39
(%)		(0.2)	(0.8)	(4.0)	(25.1)	(52.2)	(17.0)	(0.8)	(0.0)	(100.0)				
血液濾過		1	1	3	5	3				13	3	16	3.00	0.61
(%)		(7.7)	(7.7)	(23.1)	(38.5)	(23.1)				(100.0)				
血液吸着透析 (リクセル等使用)		2	10	63	355	713	157	6		1,306	70	1,376	3.57	0.38
(%)		(0.2)	(0.8)	(4.8)	(27.2)	(54.6)	(12.0)	(0.5)		(100.0)				
在宅血液透析				3	35	165	196	40	1	440	115	555	3.95	0.38
(%)				(0.7)	(8.0)	(37.5)	(44.5)	(9.1)	(0.2)	(100.0)				
腹膜透析 (CAPD)	12	46	180	593	1,262	938	173	12	2	3,218	1,684	4,902	3.23	0.52
(%)	(0.4)	(1.4)	(5.6)	(18.4)	(39.2)	(29.1)	(5.4)	(0.4)	(0.1)	(100.0)				
腹膜透析 (APD)	7	27	91	339	714	590	148	14	2	1,932	843	2,775	3.28	0.53
(%)	(0.4)	(1.4)	(4.7)	(17.5)	(37.0)	(30.5)	(7.7)	(0.7)	(0.1)	(100.0)				
腹膜透析 (CCPD)		3	20	125	339	237	49	4		777	315	1,092	3.31	0.44
(%)		(0.4)	(2.6)	(16.1)	(43.6)	(30.5)	(6.3)	(0.5)		(100.0)				
合計	203	1,080	4,552	16,731	72,267	138,051	45,525	2,283	131	280,823	32,394	313,217	3.57	0.44
(%)	(0.1)	(0.4)	(1.6)	(6.0)	(25.7)	(49.2)	(16.2)	(0.8)	(0.0)	(100.0)				
記載なし														
(%)														
総計	203	1,080	4,552	16,731	72,267	138,051	45,525	2,283	131	280,823	32,394	313,217	3.57	0.44
(%)	(0.1)	(0.4)	(1.6)	(6.0)	(25.7)	(49.2)	(16.2)	(0.8)	(0.0)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

*腹膜透析(CCPD)は、APDと手動のバック交換の両方を行うもの

(C)Japanese Society for Dialysis Therapy

2015年調査