

表 2727 透析前アルブミン濃度(g/dL) 併用の状況別 (透析患者全体)

併用の状況	1.5未満	1.5～	2.0～	2.5～	3.0～	3.5～	4.0～	4.5～	5.0～	合計	記載なし	総計	平均	標準偏差
HD(F)単独	182	997	4,248	15,632	69,874	136,194	45,134	2,253	127	274,641	29,522	304,163	3.58	0.44
(%)	(0.1)	(0.4)	(1.5)	(5.7)	(25.4)	(49.6)	(16.4)	(0.8)	(0.0)	(100.0)				
PD単独	17	68	246	815	1,727	1,374	330	28	3	4,608	2,496	7,104	3.26	0.53
(%)	(0.4)	(1.5)	(5.3)	(17.7)	(37.5)	(29.8)	(7.2)	(0.6)	(0.1)	(100.0)				
HD(F)単独.PDカテあり	1	4	13	31	46	51	12			158	24	182	3.21	0.59
(%)	(0.6)	(2.5)	(8.2)	(19.6)	(29.1)	(32.3)	(7.6)			(100.0)				
PD.HD併用週1 HD(F)	2	8	37	219	527	340	32	1	1	1,167	303	1,470	3.24	0.44
(%)	(0.2)	(0.7)	(3.2)	(18.8)	(45.2)	(29.1)	(2.7)	(0.1)	(0.1)	(100.0)				
PD.HD併用週2 HD(F)		1	3	14	52	60	10			140	32	172	3.41	0.42
(%)		(0.7)	(2.1)	(10.0)	(37.1)	(42.9)	(7.1)			(100.0)				
PD.HD併用週3 HD(F)	1			3	8	12	2			26	3	29	3.37	0.56
(%)	(3.8)			(11.5)	(30.8)	(46.2)	(7.7)			(100.0)				
PD.HD併用週4 HD(F)														
(%)														
PD.HD併用その他			5	16	27	9	2	1		60	11	71	3.16	0.45
(%)			(8.3)	(26.7)	(45.0)	(15.0)	(3.3)	(1.7)		(100.0)				
合計	203	1,078	4,552	16,730	72,261	138,040	45,522	2,283	131	280,800	32,391	313,191	3.57	0.44
(%)	(0.1)	(0.4)	(1.6)	(6.0)	(25.7)	(49.2)	(16.2)	(0.8)	(0.0)	(100.0)				
不明		2		1	6	11	3			23	2	25	3.43	0.62
(%)		(8.7)		(4.3)	(26.1)	(47.8)	(13.0)			(100.0)				
記載なし											1	1		
(%)														
総計	203	1,080	4,552	16,731	72,267	138,051	45,525	2,283	131	280,823	32,394	313,217	3.57	0.44
(%)	(0.1)	(0.4)	(1.6)	(6.0)	(25.7)	(49.2)	(16.2)	(0.8)	(0.0)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査