

表 2731 透析前アルブミン濃度(g/dL) 原疾患別（透析患者全体）

原疾患	1.5未満	1.5～	2.0～	2.5～	3.0～	3.5～	4.0～	4.5～	5.0～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	41	216	969	4,124	20,731	42,314	14,462	756	47	83,660	9,687	93,347	3.60	0.42
(%)	(0.0)	(0.3)	(1.2)	(4.9)	(24.8)	(50.6)	(17.3)	(0.9)	(0.1)	(100.0)				
慢性腎盂腎炎	1	11	37	172	719	1,262	438	23	3	2,666	269	2,935	3.56	0.44
(%)	(0.0)	(0.4)	(1.4)	(6.5)	(27.0)	(47.3)	(16.4)	(0.9)	(0.1)	(100.0)				
急速進行性糸球体腎炎	1	14	66	203	705	950	236	20	1	2,196	282	2,478	3.45	0.47
(%)	(0.0)	(0.6)	(3.0)	(9.2)	(32.1)	(43.3)	(10.7)	(0.9)	(0.0)	(100.0)				
妊娠腎/妊娠中毒症	1	1	10	45	336	778	233	12	2	1,418	127	1,545	3.63	0.38
(%)	(0.1)	(0.1)	(0.7)	(3.2)	(23.7)	(54.9)	(16.4)	(0.8)	(0.1)	(100.0)				
その他の分類不能の腎炎	1	6	21	61	291	603	264	26		1,273	132	1,405	3.63	0.45
(%)	(0.1)	(0.5)	(1.6)	(4.8)	(22.9)	(47.4)	(20.7)	(2.0)		(100.0)				
多発性嚢胞腎	4	31	109	474	2,654	5,317	1,539	61	2	10,191	1,065	11,256	3.58	0.41
(%)	(0.0)	(0.3)	(1.1)	(4.7)	(26.0)	(52.2)	(15.1)	(0.6)	(0.0)	(100.0)				
腎硬化症	18	91	500	1,859	7,931	12,677	3,496	194	8	26,774	3,031	29,805	3.52	0.44
(%)	(0.1)	(0.3)	(1.9)	(6.9)	(29.6)	(47.3)	(13.1)	(0.7)	(0.0)	(100.0)				
悪性高血圧		5	27	101	537	1,137	471	44	5	2,327	291	2,618	3.65	0.43
(%)		(0.2)	(1.2)	(4.3)	(23.1)	(48.9)	(20.2)	(1.9)	(0.2)	(100.0)				
糖尿病性腎症	90	460	1,912	6,661	27,214	53,073	17,837	752	41	108,040	12,238	120,278	3.57	0.45
(%)	(0.1)	(0.4)	(1.8)	(6.2)	(25.2)	(49.1)	(16.5)	(0.7)	(0.0)	(100.0)				
SLE腎炎	2	9	26	144	650	903	242	16	2	1,994	233	2,227	3.50	0.43
(%)	(0.1)	(0.5)	(1.3)	(7.2)	(32.6)	(45.3)	(12.1)	(0.8)	(0.1)	(100.0)				
アミロイド腎		9	20	66	116	151	40	1		403	59	462	3.34	0.54
(%)		(2.2)	(5.0)	(16.4)	(28.8)	(37.5)	(9.9)	(0.2)		(100.0)				
痛風腎	1	2	11	43	227	530	152	8		974	101	1,075	3.60	0.40
(%)	(0.1)	(0.2)	(1.1)	(4.4)	(23.3)	(54.4)	(15.6)	(0.8)		(100.0)				
先天性代謝異常による腎不全		3	5	14	44	91	73	8		238	36	274	3.69	0.52
(%)		(1.3)	(2.1)	(5.9)	(18.5)	(38.2)	(30.7)	(3.4)		(100.0)				
腎・尿路結核		1	7	15	45	78	26			172	23	195	3.47	0.49
(%)		(0.6)	(4.1)	(8.7)	(26.2)	(45.3)	(15.1)			(100.0)				
腎・尿路結石		6	13	38	148	249	71	3		528	53	581	3.51	0.48
(%)		(1.1)	(2.5)	(7.2)	(28.0)	(47.2)	(13.4)	(0.6)		(100.0)				
腎・尿路腫瘍		9	21	47	256	393	89	5		820	90	910	3.49	0.45
(%)		(1.1)	(2.6)	(5.7)	(31.2)	(47.9)	(10.9)	(0.6)		(100.0)				
閉塞性尿路障害	3	1	11	36	159	328	106	10		654	76	730	3.58	0.47
(%)	(0.5)	(0.2)	(1.7)	(5.5)	(24.3)	(50.2)	(16.2)	(1.5)		(100.0)				
骨髄腫		4	10	30	83	90	34	2		253	38	291	3.41	0.54
(%)		(1.6)	(4.0)	(11.9)	(32.8)	(35.6)	(13.4)	(0.8)		(100.0)				
腎形成不全			5	18	97	268	181	20	1	590	70	660	3.76	0.42
(%)			(0.8)	(3.1)	(16.4)	(45.4)	(30.7)	(3.4)	(0.2)	(100.0)				
不明	25	141	578	1,938	7,031	12,589	3,982	224	10	26,518	3,379	29,897	3.53	0.46
(%)	(0.1)	(0.5)	(2.2)	(7.3)	(26.5)	(47.5)	(15.0)	(0.8)	(0.0)	(100.0)				
再導入		6	22	77	420	992	396	15	4	1,932	251	2,183	3.64	0.42
(%)		(0.3)	(1.1)	(4.0)	(21.7)	(51.3)	(20.5)	(0.8)	(0.2)	(100.0)				
その他	15	54	172	561	1,869	3,272	1,156	83	5	7,187	857	8,044	3.53	0.50
(%)	(0.2)	(0.8)	(2.4)	(7.8)	(26.0)	(45.5)	(16.1)	(1.2)	(0.1)	(100.0)				
合計	203	1,080	4,552	16,727	72,263	138,045	45,524	2,283	131	280,808	32,388	313,196	3.57	0.44
(%)	(0.1)	(0.4)	(1.6)	(6.0)	(25.7)	(49.2)	(16.2)	(0.8)	(0.0)	(100.0)				
記載なし				4	4	6	1			15	6	21	3.35	0.48
(%)				(26.7)	(26.7)	(40.0)	(6.7)			(100.0)				
総計	203	1,080	4,552	16,731	72,267	138,051	45,525	2,283	131	280,823	32,394	313,217	3.57	0.44
(%)	(0.1)	(0.4)	(1.6)	(6.0)	(25.7)	(49.2)	(16.2)	(0.8)	(0.0)	(100.0)				

数値下のかつこ内は行方向の合計に対する％です。
(C)Japanese Society for Dialysis Therapy
2015年調査