

表 2732 透析前アルブミン濃度(g/dL) 医療機関別（透析患者全体）

医療機関	1.5未満	1.5～	2.0～	2.5～	3.0～	3.5～	4.0～	4.5～	5.0～	合計	記載なし	総計	平均	標準偏差
国公立大学	5	23	88	170	272	218	56	3	1	836	372	1,208	3.14	0.61
(%)	(0.6)	(2.8)	(10.5)	(20.3)	(32.5)	(26.1)	(6.7)	(0.4)	(0.1)	(100.0)				
私立大学	9	48	137	301	705	909	355	15		2,479	1,085	3,564	3.38	0.58
(%)	(0.4)	(1.9)	(5.5)	(12.1)	(28.4)	(36.7)	(14.3)	(0.6)		(100.0)				
国立	6	13	46	110	225	252	90	2	3	747	177	924	3.33	0.61
(%)	(0.8)	(1.7)	(6.2)	(14.7)	(30.1)	(33.7)	(12.0)	(0.3)	(0.4)	(100.0)				
県市町村立	19	143	505	1,424	4,917	7,162	2,195	115	6	16,486	3,032	19,518	3.47	0.49
(%)	(0.1)	(0.9)	(3.1)	(8.6)	(29.8)	(43.4)	(13.3)	(0.7)	(0.0)	(100.0)				
社会保険	7	18	108	311	1,016	1,517	418	20		3,415	484	3,899	3.47	0.48
(%)	(0.2)	(0.5)	(3.2)	(9.1)	(29.8)	(44.4)	(12.2)	(0.6)		(100.0)				
厚生連	9	53	177	651	2,370	3,455	1,148	80	13	7,956	1,045	9,001	3.51	0.48
(%)	(0.1)	(0.7)	(2.2)	(8.2)	(29.8)	(43.4)	(14.4)	(1.0)	(0.2)	(100.0)				
その他公的	15	71	268	793	2,500	3,884	1,127	47	4	8,709	2,215	10,924	3.47	0.49
(%)	(0.2)	(0.8)	(3.1)	(9.1)	(28.7)	(44.6)	(12.9)	(0.5)	(0.0)	(100.0)				
私立総合	16	34	139	528	1,862	2,888	879	42	5	6,393	1,055	7,448	3.50	0.48
(%)	(0.3)	(0.5)	(2.2)	(8.3)	(29.1)	(45.2)	(13.7)	(0.7)	(0.1)	(100.0)				
私立	94	544	2,144	6,834	22,684	36,053	11,218	535	45	80,151	9,914	90,065	3.49	0.48
(%)	(0.1)	(0.7)	(2.7)	(8.5)	(28.3)	(45.0)	(14.0)	(0.7)	(0.1)	(100.0)				
私立診療所	23	133	940	5,609	35,716	81,713	28,039	1,424	54	153,651	13,015	166,666	3.64	0.38
(%)	(0.0)	(0.1)	(0.6)	(3.7)	(23.2)	(53.2)	(18.2)	(0.9)	(0.0)	(100.0)				
総計	203	1,080	4,552	16,731	72,267	138,051	45,525	2,283	131	280,823	32,394	313,217	3.57	0.44
(%)	(0.1)	(0.4)	(1.6)	(6.0)	(25.7)	(49.2)	(16.2)	(0.8)	(0.0)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査