

表 2748 透析前アルブミン濃度(g/dL) 原疾患別（施設血液透析患者 透析回数週3回）

原疾患	1.5未満	1.5～	2.0～	2.5～	3.0～	3.5～	4.0～	4.5～	5.0～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	31	166	747	2,997	14,907	30,722	10,871	584	35	61,060	2,414	63,474	3.60	0.42
(%)	(0.1)	(0.3)	(1.2)	(4.9)	(24.4)	(50.3)	(17.8)	(1.0)	(0.1)	(100.0)				
慢性腎盂腎炎	1	9	28	123	545	932	319	16	1	1,974	66	2,040	3.56	0.44
(%)	(0.1)	(0.5)	(1.4)	(6.2)	(27.6)	(47.2)	(16.2)	(0.8)	(0.1)	(100.0)				
急速進行性糸球体腎炎		10	48	160	562	719	187	14	1	1,701	70	1,771	3.45	0.47
(%)		(0.6)	(2.8)	(9.4)	(33.0)	(42.3)	(11.0)	(0.8)	(0.1)	(100.0)				
妊娠腎/妊娠中毒症	1	1	7	33	209	505	174	10	2	942	39	981	3.64	0.40
(%)	(0.1)	(0.1)	(0.7)	(3.5)	(22.2)	(53.6)	(18.5)	(1.1)	(0.2)	(100.0)				
その他の分類不能の腎炎		5	16	45	204	426	186	16		898	31	929	3.62	0.45
(%)		(0.6)	(1.8)	(5.0)	(22.7)	(47.4)	(20.7)	(1.8)		(100.0)				
多発性嚢胞腎	4	21	88	347	1,971	4,057	1,216	47	2	7,753	263	8,016	3.59	0.41
(%)	(0.1)	(0.3)	(1.1)	(4.5)	(25.4)	(52.3)	(15.7)	(0.6)	(0.0)	(100.0)				
腎硬化症	11	70	389	1,475	6,200	10,109	2,821	151	7	21,233	774	22,007	3.52	0.44
(%)	(0.1)	(0.3)	(1.8)	(6.9)	(29.2)	(47.6)	(13.3)	(0.7)	(0.0)	(100.0)				
悪性高血圧		4	23	78	415	870	353	37	3	1,783	78	1,861	3.64	0.43
(%)		(0.2)	(1.3)	(4.4)	(23.3)	(48.8)	(19.8)	(2.1)	(0.2)	(100.0)				
糖尿病性腎症	71	362	1,543	5,311	21,422	42,260	14,498	631	31	86,129	3,297	89,426	3.57	0.45
(%)	(0.1)	(0.4)	(1.8)	(6.2)	(24.9)	(49.1)	(16.8)	(0.7)	(0.0)	(100.0)				
SLE腎炎	2	9	19	106	468	648	174	14	1	1,441	63	1,504	3.49	0.44
(%)	(0.1)	(0.6)	(1.3)	(7.4)	(32.5)	(45.0)	(12.1)	(1.0)	(0.1)	(100.0)				
アミロイド腎		5	12	46	94	110	32	1		300	14	314	3.36	0.51
(%)		(1.7)	(4.0)	(15.3)	(31.3)	(36.7)	(10.7)	(0.3)		(100.0)				
痛風腎	1	1	9	32	167	418	118	7		753	25	778	3.61	0.40
(%)	(0.1)	(0.1)	(1.2)	(4.2)	(22.2)	(55.5)	(15.7)	(0.9)		(100.0)				
先天性代謝異常による腎不全		2	5	10	32	62	52	6		169	4	173	3.68	0.54
(%)		(1.2)	(3.0)	(5.9)	(18.9)	(36.7)	(30.8)	(3.6)		(100.0)				
腎・尿路結核		1	7	11	37	59	19			134	7	141	3.45	0.52
(%)		(0.7)	(5.2)	(8.2)	(27.6)	(44.0)	(14.2)			(100.0)				
腎・尿路結石		6	10	30	114	189	60	3		412	9	421	3.52	0.49
(%)		(1.5)	(2.4)	(7.3)	(27.7)	(45.9)	(14.6)	(0.7)		(100.0)				
腎・尿路腫瘍		6	19	40	206	314	71	5		661	17	678	3.48	0.45
(%)		(0.9)	(2.9)	(6.1)	(31.2)	(47.5)	(10.7)	(0.8)		(100.0)				
閉塞性尿路障害	1	1	9	26	115	245	73	5		475	19	494	3.57	0.46
(%)	(0.2)	(0.2)	(1.9)	(5.5)	(24.2)	(51.6)	(15.4)	(1.1)		(100.0)				
骨髄腫		3	8	25	61	60	25			182	8	190	3.36	0.54
(%)		(1.6)	(4.4)	(13.7)	(33.5)	(33.0)	(13.7)			(100.0)				
腎形成不全			2	9	63	179	122	12	1	388	12	400	3.76	0.40
(%)			(0.5)	(2.3)	(16.2)	(46.1)	(31.4)	(3.1)	(0.3)	(100.0)				
不明	17	104	439	1,541	5,369	9,466	3,065	165	10	20,176	1,000	21,176	3.53	0.47
(%)	(0.1)	(0.5)	(2.2)	(7.6)	(26.6)	(46.9)	(15.2)	(0.8)	(0.0)	(100.0)				
再導入		4	18	47	273	673	286	11	4	1,316	40	1,356	3.66	0.43
(%)		(0.3)	(1.4)	(3.6)	(20.7)	(51.1)	(21.7)	(0.8)	(0.3)	(100.0)				
その他	12	37	133	447	1,445	2,463	885	68	5	5,495	220	5,715	3.53	0.50
(%)	(0.2)	(0.7)	(2.4)	(8.1)	(26.3)	(44.8)	(16.1)	(1.2)	(0.1)	(100.0)				
合計	152	827	3,579	12,939	54,879	105,486	35,607	1,803	103	215,375	8,470	223,845	3.57	0.44
(%)	(0.1)	(0.4)	(1.7)	(6.0)	(25.5)	(49.0)	(16.5)	(0.8)	(0.0)	(100.0)				
記載なし				4	1	1				6	1	7	3.02	0.53
(%)				(66.7)	(16.7)	(16.7)				(100.0)				
総計	152	827	3,579	12,943	54,880	105,487	35,607	1,803	103	215,381	8,471	223,852	3.57	0.44
(%)	(0.1)	(0.4)	(1.7)	(6.0)	(25.5)	(49.0)	(16.5)	(0.8)	(0.0)	(100.0)				

数値下のかつこ内は行方向の合計に対する％です。
(C)Japanese Society for Dialysis Therapy
2015年調査