

表 2749 透析前アルブミン濃度(g/dL) 医療機関別（施設血液透析患者 透析回数週3回）

医療機関	1.5未満	1.5～	2.0～	2.5～	3.0～	3.5～	4.0～	4.5～	5.0～	合計	記載なし	総計	平均	標準偏差
国公立大学	4	14	67	90	105	90	25		1	396	12	408	3.03	0.66
(%)	(1.0)	(3.5)	(16.9)	(22.7)	(26.5)	(22.7)	(6.3)		(0.3)	(100.0)				
私立大学	5	36	91	165	316	575	243	7		1,438	58	1,496	3.41	0.61
(%)	(0.3)	(2.5)	(6.3)	(11.5)	(22.0)	(40.0)	(16.9)	(0.5)		(100.0)				
国立	2	7	29	58	144	186	69	1	3	499	16	515	3.40	0.59
(%)	(0.4)	(1.4)	(5.8)	(11.6)	(28.9)	(37.3)	(13.8)	(0.2)	(0.6)	(100.0)				
県市町村立	15	101	370	1,011	3,570	5,465	1,727	82	5	12,346	624	12,970	3.49	0.49
(%)	(0.1)	(0.8)	(3.0)	(8.2)	(28.9)	(44.3)	(14.0)	(0.7)	(0.0)	(100.0)				
社会保険	7	11	76	244	760	1,187	330	19		2,634	41	2,675	3.48	0.48
(%)	(0.3)	(0.4)	(2.9)	(9.3)	(28.9)	(45.1)	(12.5)	(0.7)		(100.0)				
厚生連	4	45	132	462	1,786	2,716	927	62	13	6,147	276	6,423	3.52	0.48
(%)	(0.1)	(0.7)	(2.1)	(7.5)	(29.1)	(44.2)	(15.1)	(1.0)	(0.2)	(100.0)				
その他公的	10	46	194	522	1,810	3,070	900	38	2	6,592	324	6,916	3.50	0.48
(%)	(0.2)	(0.7)	(2.9)	(7.9)	(27.5)	(46.6)	(13.7)	(0.6)	(0.0)	(100.0)				
私立総合	11	28	97	363	1,277	2,053	687	35	3	4,554	235	4,789	3.51	0.48
(%)	(0.2)	(0.6)	(2.1)	(8.0)	(28.0)	(45.1)	(15.1)	(0.8)	(0.1)	(100.0)				
私立	75	438	1,768	5,541	18,026	28,294	9,056	434	38	63,670	2,927	66,597	3.49	0.49
(%)	(0.1)	(0.7)	(2.8)	(8.7)	(28.3)	(44.4)	(14.2)	(0.7)	(0.1)	(100.0)				
私立診療所	19	101	755	4,487	27,086	61,851	21,643	1,125	38	117,105	3,958	121,063	3.64	0.38
(%)	(0.0)	(0.1)	(0.6)	(3.8)	(23.1)	(52.8)	(18.5)	(1.0)	(0.0)	(100.0)				
総計	152	827	3,579	12,943	54,880	105,487	35,607	1,803	103	215,381	8,471	223,852	3.57	0.44
(%)	(0.1)	(0.4)	(1.7)	(6.0)	(25.5)	(49.0)	(16.5)	(0.8)	(0.0)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査