

表 2766 透析前アルブミン濃度(g/dL) 医療機関別（血液透析濾過患者 透析回数週3回）

医療機関	1.5未満	1.5～	2.0～	2.5～	3.0～	3.5～	4.0～	4.5～	5.0～	合計	記載なし	総計	平均	標準偏差
国公立大学		2	4	12	14	11	4			47	2	49	3.13	0.60
(%)		(4.3)	(8.5)	(25.5)	(29.8)	(23.4)	(8.5)			(100.0)				
私立大学		2	5	8	53	78	26	1		173	2	175	3.50	0.47
(%)		(1.2)	(2.9)	(4.6)	(30.6)	(45.1)	(15.0)	(0.6)		(100.0)				
国立		2	6	6	15	16	6			51		51	3.25	0.62
(%)		(3.9)	(11.8)	(11.8)	(29.4)	(31.4)	(11.8)			(100.0)				
県市町村立		8	36	116	681	945	222	8	1	2,017	98	2,115	3.50	0.41
(%)		(0.4)	(1.8)	(5.8)	(33.8)	(46.9)	(11.0)	(0.4)	(0.0)	(100.0)				
社会保険			6	20	130	192	49	1		398	15	413	3.53	0.39
(%)			(1.5)	(5.0)	(32.7)	(48.2)	(12.3)	(0.3)		(100.0)				
厚生連		2	8	48	287	472	129	6		952	70	1,022	3.56	0.39
(%)		(0.2)	(0.8)	(5.0)	(30.1)	(49.6)	(13.6)	(0.6)		(100.0)				
その他公的		2	13	39	215	401	131	6	1	808	46	854	3.58	0.42
(%)		(0.2)	(1.6)	(4.8)	(26.6)	(49.6)	(16.2)	(0.7)	(0.1)	(100.0)				
私立総合	2	2	20	87	371	589	137	6	2	1,216	26	1,242	3.51	0.43
(%)	(0.2)	(0.2)	(1.6)	(7.2)	(30.5)	(48.4)	(11.3)	(0.5)	(0.2)	(100.0)				
私立	4	40	175	754	3,259	5,952	1,565	56	1	11,806	448	12,254	3.54	0.42
(%)	(0.0)	(0.3)	(1.5)	(6.4)	(27.6)	(50.4)	(13.3)	(0.5)	(0.0)	(100.0)				
私立診療所	1	11	108	854	7,306	17,227	5,105	195	12	30,819	587	31,406	3.64	0.35
(%)	(0.0)	(0.0)	(0.4)	(2.8)	(23.7)	(55.9)	(16.6)	(0.6)	(0.0)	(100.0)				
総計	7	71	381	1,944	12,331	25,883	7,374	279	17	48,287	1,294	49,581	3.60	0.38
(%)	(0.0)	(0.1)	(0.8)	(4.0)	(25.5)	(53.6)	(15.3)	(0.6)	(0.0)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査