

表 2798 透析前CRP濃度(mg/dL) 併用の状況別 (透析患者全体)

併用の状況	0.2未満	0.2～	0.5～	1.0～	2.0～	4.0～	6.0～	8.0～	10.0～	15.0～	合計	記載なし	総計	平均	標準偏差
HD(F)単独	131,774	45,826	23,359	15,151	8,903	3,270	1,577	814	926	604	232,204	71,959	304,163	0.62	1.82
(%)	(56.7)	(19.7)	(10.1)	(6.5)	(3.8)	(1.4)	(0.7)	(0.4)	(0.4)	(0.3)	(100.0)				
PD単独	2,300	789	414	261	179	63	28	13	17	7	4,071	3,033	7,104	0.62	1.60
(%)	(56.5)	(19.4)	(10.2)	(6.4)	(4.4)	(1.5)	(0.7)	(0.3)	(0.4)	(0.2)	(100.0)				
HD(F)単独.PDカテあり	69	28	12	11	8	4	2	1	3	2	140	42	182	1.30	3.33
(%)	(49.3)	(20.0)	(8.6)	(7.9)	(5.7)	(2.9)	(1.4)	(0.7)	(2.1)	(1.4)	(100.0)				
PD.HD併用週1 HD(F)	589	224	107	58	37	11	3	1	2	3	1,035	435	1,470	0.51	1.42
(%)	(56.9)	(21.6)	(10.3)	(5.6)	(3.6)	(1.1)	(0.3)	(0.1)	(0.2)	(0.3)	(100.0)				
PD.HD併用週2 HD(F)	58	27	20	10	3	4	1		1		124	48	172	0.68	1.37
(%)	(46.8)	(21.8)	(16.1)	(8.1)	(2.4)	(3.2)	(0.8)		(0.8)		(100.0)				
PD.HD併用週3 HD(F)	12	4	1	5			1	1			24	5	29	1.04	2.14
(%)	(50.0)	(16.7)	(4.2)	(20.8)			(4.2)	(4.2)			(100.0)				
PD.HD併用週4 HD(F)															
(%)															
PD.HD併用その他	34	11	6	2	6	1		1			61	10	71	0.71	1.51
(%)	(55.7)	(18.0)	(9.8)	(3.3)	(9.8)	(1.6)		(1.6)			(100.0)				
合計	134,836	46,909	23,919	15,498	9,136	3,353	1,612	831	949	616	237,659	75,532	313,191	0.62	1.81
(%)	(56.7)	(19.7)	(10.1)	(6.5)	(3.8)	(1.4)	(0.7)	(0.3)	(0.4)	(0.3)	(100.0)				
不明	11	4		4	2					1	22	3	25	1.84	5.73
(%)	(50.0)	(18.2)		(18.2)	(9.1)					(4.5)	(100.0)				
記載なし												1	1		
(%)															
総計	134,847	46,913	23,919	15,502	9,138	3,353	1,612	831	949	617	237,681	75,536	313,217	0.62	1.81
(%)	(56.7)	(19.7)	(10.1)	(6.5)	(3.8)	(1.4)	(0.7)	(0.3)	(0.4)	(0.3)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査