

表 2802 透析前CRP濃度(mg/dL) 原疾患別（透析患者全体）

原疾患	0.2未満	0.2～	0.5～	1.0～	2.0～	4.0～	6.0～	8.0～	10.0～	15.0～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	41,934	13,451	6,397	4,037	2,367	829	407	194	206	156	69,978	23,369	93,347	0.55	1.64
(%)	(59.9)	(19.2)	(9.1)	(5.8)	(3.4)	(1.2)	(0.6)	(0.3)	(0.3)	(0.2)	(100.0)				
慢性腎盂腎炎	1,272	438	214	152	80	33	15	10	14	9	2,237	698	2,935	0.73	2.83
(%)	(56.9)	(19.6)	(9.6)	(6.8)	(3.6)	(1.5)	(0.7)	(0.4)	(0.6)	(0.4)	(100.0)				
急速進行性糸球体腎炎	1,043	354	208	142	90	38	17	10	8	3	1,913	565	2,478	0.71	1.88
(%)	(54.5)	(18.5)	(10.9)	(7.4)	(4.7)	(2.0)	(0.9)	(0.5)	(0.4)	(0.2)	(100.0)				
妊娠腎/妊娠中毒症	752	216	90	59	30	12	2	3	3	1	1,168	377	1,545	0.43	1.16
(%)	(64.4)	(18.5)	(7.7)	(5.1)	(2.6)	(1.0)	(0.2)	(0.3)	(0.3)	(0.1)	(100.0)				
その他の分類不能の腎炎	653	191	75	71	39	15	2	2	7		1,055	350	1,405	0.52	1.30
(%)	(61.9)	(18.1)	(7.1)	(6.7)	(3.7)	(1.4)	(0.2)	(0.2)	(0.7)		(100.0)				
多発性嚢胞腎	4,749	1,769	915	576	361	113	63	32	30	33	8,641	2,615	11,256	0.66	1.95
(%)	(55.0)	(20.5)	(10.6)	(6.7)	(4.2)	(1.3)	(0.7)	(0.4)	(0.3)	(0.4)	(100.0)				
腎硬化症	12,450	4,730	2,504	1,600	896	344	149	80	103	56	22,912	6,893	29,805	0.64	1.74
(%)	(54.3)	(20.6)	(10.9)	(7.0)	(3.9)	(1.5)	(0.7)	(0.3)	(0.4)	(0.2)	(100.0)				
悪性高血圧	1,185	412	190	97	59	26	10		5	3	1,987	631	2,618	0.47	1.22
(%)	(59.6)	(20.7)	(9.6)	(4.9)	(3.0)	(1.3)	(0.5)		(0.3)	(0.2)	(100.0)				
糖尿病性腎症	50,952	18,417	9,610	6,214	3,720	1,352	643	332	401	243	91,884	28,394	120,278	0.65	1.91
(%)	(55.5)	(20.0)	(10.5)	(6.8)	(4.0)	(1.5)	(0.7)	(0.4)	(0.4)	(0.3)	(100.0)				
SLE腎炎	953	325	167	135	72	27	16	5	10	3	1,713	514	2,227	0.67	1.69
(%)	(55.6)	(19.0)	(9.7)	(7.9)	(4.2)	(1.6)	(0.9)	(0.3)	(0.6)	(0.2)	(100.0)				
アミロイド腎	187	75	29	30	22	14	1	2	4	4	368	94	462	1.06	2.83
(%)	(50.8)	(20.4)	(7.9)	(8.2)	(6.0)	(3.8)	(0.3)	(0.5)	(1.1)	(1.1)	(100.0)				
痛風腎	443	176	85	61	25	6	9	3	1	1	810	265	1,075	0.55	1.29
(%)	(54.7)	(21.7)	(10.5)	(7.5)	(3.1)	(0.7)	(1.1)	(0.4)	(0.1)	(0.1)	(100.0)				
先天性代謝異常による腎不全	132	34	16	9	6	4	2	2			205	69	274	0.55	1.37
(%)	(64.4)	(16.6)	(7.8)	(4.4)	(2.9)	(2.0)	(1.0)	(1.0)			(100.0)				
腎・尿路結核	76	31	12	6	5	5	2	5	2	1	145	50	195	1.17	2.72
(%)	(52.4)	(21.4)	(8.3)	(4.1)	(3.4)	(3.4)	(1.4)	(3.4)	(1.4)	(0.7)	(100.0)				
腎・尿路結石	250	87	44	32	17	7	6	4	2	1	450	131	581	0.72	1.77
(%)	(55.6)	(19.3)	(9.8)	(7.1)	(3.8)	(1.6)	(1.3)	(0.9)	(0.4)	(0.2)	(100.0)				
腎・尿路腫瘍	348	139	73	55	40	11	6	9	7	4	692	218	910	0.96	2.51
(%)	(50.3)	(20.1)	(10.5)	(7.9)	(5.8)	(1.6)	(0.9)	(1.3)	(1.0)	(0.6)	(100.0)				
閉塞性尿路障害	303	101	65	54	17	8	6	1	10	2	567	163	730	0.82	2.02
(%)	(53.4)	(17.8)	(11.5)	(9.5)	(3.0)	(1.4)	(1.1)	(0.2)	(1.8)	(0.4)	(100.0)				
骨髄腫	123	36	23	24	14	2	1	3	1		227	64	291	0.72	1.49
(%)	(54.2)	(15.9)	(10.1)	(10.6)	(6.2)	(0.9)	(0.4)	(1.3)	(0.4)		(100.0)				
腎形成不全	332	102	31	25	6	6	2	1			505	155	660	0.35	0.86
(%)	(65.7)	(20.2)	(6.1)	(5.0)	(1.2)	(1.2)	(0.4)	(0.2)			(100.0)				
不明	12,451	4,280	2,387	1,564	942	350	186	94	103	71	22,428	7,469	29,897	0.68	1.87
(%)	(55.5)	(19.1)	(10.6)	(7.0)	(4.2)	(1.6)	(0.8)	(0.4)	(0.5)	(0.3)	(100.0)				
再導入	926	345	154	120	61	23	10	3	7	6	1,655	528	2,183	0.62	1.71
(%)	(56.0)	(20.8)	(9.3)	(7.3)	(3.7)	(1.4)	(0.6)	(0.2)	(0.4)	(0.4)	(100.0)				
その他	3,328	1,203	627	437	269	127	57	36	24	20	6,128	1,916	8,044	0.72	1.85
(%)	(54.3)	(19.6)	(10.2)	(7.1)	(4.4)	(2.1)	(0.9)	(0.6)	(0.4)	(0.3)	(100.0)				
合計	134,842	46,912	23,916	15,500	9,138	3,352	1,612	831	948	617	237,668	75,528	313,196	0.62	1.81
(%)	(56.7)	(19.7)	(10.1)	(6.5)	(3.8)	(1.4)	(0.7)	(0.3)	(0.4)	(0.3)	(100.0)				
記載なし	5	1	3	2		1			1		13	8	21	1.67	2.97
(%)	(38.5)	(7.7)	(23.1)	(15.4)		(7.7)			(7.7)		(100.0)				
総計	134,847	46,913	23,919	15,502	9,138	3,353	1,612	831	949	617	237,681	75,536	313,217	0.62	1.81
(%)	(56.7)	(19.7)	(10.1)	(6.5)	(3.8)	(1.4)	(0.7)	(0.3)	(0.4)	(0.3)	(100.0)				

数値下のかつこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査