

表 2818 透析前CRP濃度(mg/dL) 原疾患別（施設血液透析患者 透析回数週3回）

原疾患	0.2未満	0.2～	0.5～	1.0～	2.0～	4.0～	6.0～	8.0～	10.0～	15.0～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	30,074	9,831	4,695	2,939	1,765	608	305	143	158	117	50,635	12,839	63,474	0.56	1.68
(%)	(59.4)	(19.4)	(9.3)	(5.8)	(3.5)	(1.2)	(0.6)	(0.3)	(0.3)	(0.2)	(100.0)				
慢性腎盂腎炎	920	331	159	114	56	23	13	7	11	8	1,642	398	2,040	0.77	3.17
(%)	(56.0)	(20.2)	(9.7)	(6.9)	(3.4)	(1.4)	(0.8)	(0.4)	(0.7)	(0.5)	(100.0)				
急速進行性糸球体腎炎	802	274	166	109	68	33	12	7	6	2	1,479	292	1,771	0.71	1.93
(%)	(54.2)	(18.5)	(11.2)	(7.4)	(4.6)	(2.2)	(0.8)	(0.5)	(0.4)	(0.1)	(100.0)				
妊娠腎/妊娠中毒症	489	145	60	36	18	6		3	3	1	761	220	981	0.43	1.28
(%)	(64.3)	(19.1)	(7.9)	(4.7)	(2.4)	(0.8)		(0.4)	(0.4)	(0.1)	(100.0)				
その他の分類不能の腎炎	460	147	53	52	25	12	2	2	6		759	170	929	0.55	1.37
(%)	(60.6)	(19.4)	(7.0)	(6.9)	(3.3)	(1.6)	(0.3)	(0.3)	(0.8)		(100.0)				
多発性嚢胞腎	3,557	1,367	675	429	282	86	48	28	23	28	6,523	1,493	8,016	0.68	2.05
(%)	(54.5)	(21.0)	(10.3)	(6.6)	(4.3)	(1.3)	(0.7)	(0.4)	(0.4)	(0.4)	(100.0)				
腎硬化症	9,797	3,762	1,928	1,270	727	276	131	64	83	46	18,084	3,923	22,007	0.65	1.78
(%)	(54.2)	(20.8)	(10.7)	(7.0)	(4.0)	(1.5)	(0.7)	(0.4)	(0.5)	(0.3)	(100.0)				
悪性高血圧	894	305	148	76	49	22	7		4	3	1,508	353	1,861	0.50	1.29
(%)	(59.3)	(20.2)	(9.8)	(5.0)	(3.2)	(1.5)	(0.5)		(0.3)	(0.2)	(100.0)				
糖尿病性腎症	40,247	14,578	7,640	4,938	2,976	1,076	507	269	321	192	72,744	16,682	89,426	0.65	1.88
(%)	(55.3)	(20.0)	(10.5)	(6.8)	(4.1)	(1.5)	(0.7)	(0.4)	(0.4)	(0.3)	(100.0)				
SLE腎炎	674	229	123	101	52	21	14	2	7	2	1,225	279	1,504	0.69	1.61
(%)	(55.0)	(18.7)	(10.0)	(8.2)	(4.2)	(1.7)	(1.1)	(0.2)	(0.6)	(0.2)	(100.0)				
アミロイド腎	139	58	25	21	10	11	1	1	4	4	274	40	314	1.12	3.16
(%)	(50.7)	(21.2)	(9.1)	(7.7)	(3.6)	(4.0)	(0.4)	(0.4)	(1.5)	(1.5)	(100.0)				
痛風腎	346	128	59	47	22	5	7	1	1	1	617	161	778	0.55	1.30
(%)	(56.1)	(20.7)	(9.6)	(7.6)	(3.6)	(0.8)	(1.1)	(0.2)	(0.2)	(0.2)	(100.0)				
先天性代謝異常による腎不全	93	20	11	6	5	3	2	1			141	32	173	0.58	1.41
(%)	(66.0)	(14.2)	(7.8)	(4.3)	(3.5)	(2.1)	(1.4)	(0.7)			(100.0)				
腎・尿路結核	59	24	10	6	5	5	2	4	2		117	24	141	1.18	2.48
(%)	(50.4)	(20.5)	(8.5)	(5.1)	(4.3)	(4.3)	(1.7)	(3.4)	(1.7)		(100.0)				
腎・尿路結石	193	71	30	23	12	5	6	2	2		344	77	421	0.68	1.59
(%)	(56.1)	(20.6)	(8.7)	(6.7)	(3.5)	(1.5)	(1.7)	(0.6)	(0.6)		(100.0)				
腎・尿路腫瘍	280	101	61	50	35	9	5	7	5	4	557	121	678	1.00	2.62
(%)	(50.3)	(18.1)	(11.0)	(9.0)	(6.3)	(1.6)	(0.9)	(1.3)	(0.9)	(0.7)	(100.0)				
閉塞性尿路障害	213	79	51	43	13	5	6	1	6	1	418	76	494	0.80	1.82
(%)	(51.0)	(18.9)	(12.2)	(10.3)	(3.1)	(1.2)	(1.4)	(0.2)	(1.4)	(0.2)	(100.0)				
骨髄腫	81	29	20	15	11			2	1		159	31	190	0.69	1.40
(%)	(50.9)	(18.2)	(12.6)	(9.4)	(6.9)			(1.3)	(0.6)		(100.0)				
腎形成不全	220	65	17	17	4	3	1	1			328	72	400	0.34	0.87
(%)	(67.1)	(19.8)	(5.2)	(5.2)	(1.2)	(0.9)	(0.3)	(0.3)			(100.0)				
不明	9,298	3,190	1,814	1,230	745	279	144	61	82	54	16,897	4,279	21,176	0.70	1.93
(%)	(55.0)	(18.9)	(10.7)	(7.3)	(4.4)	(1.7)	(0.9)	(0.4)	(0.5)	(0.3)	(100.0)				
再導入	620	226	103	86	38	17	7	2	5	4	1,108	248	1,356	0.63	1.70
(%)	(56.0)	(20.4)	(9.3)	(7.8)	(3.4)	(1.5)	(0.6)	(0.2)	(0.5)	(0.4)	(100.0)				
その他	2,482	918	483	355	205	102	46	30	19	15	4,655	1,060	5,715	0.75	1.89
(%)	(53.3)	(19.7)	(10.4)	(7.6)	(4.4)	(2.2)	(1.0)	(0.6)	(0.4)	(0.3)	(100.0)				
合計	101,938	35,878	18,331	11,963	7,123	2,607	1,266	638	749	482	180,975	42,870	223,845	0.63	1.84
(%)	(56.3)	(19.8)	(10.1)	(6.6)	(3.9)	(1.4)	(0.7)	(0.4)	(0.4)	(0.3)	(100.0)				
記載なし	2		2						1		5	2	7	2.43	4.58
(%)	(40.0)		(40.0)						(20.0)		(100.0)				
総計	101,940	35,878	18,333	11,963	7,123	2,607	1,266	638	750	482	180,980	42,872	223,852	0.63	1.84
(%)	(56.3)	(19.8)	(10.1)	(6.6)	(3.9)	(1.4)	(0.7)	(0.4)	(0.4)	(0.3)	(100.0)				

数値下のカッコ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査