

表 2834 透析前CRP濃度(mg/dL) 原疾患別（血液透析濾過患者 透析回数週3回）

原疾患	0.2未満	0.2～	0.5～	1.0～	2.0～	4.0～	6.0～	8.0～	10.0～	15.0～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	9,140	2,735	1,284	796	434	156	78	41	28	26	14,718	2,744	17,462	0.49	1.47
(%)	(62.1)	(18.6)	(8.7)	(5.4)	(2.9)	(1.1)	(0.5)	(0.3)	(0.2)	(0.2)	(100.0)				
慢性腎盂腎炎	254	77	37	27	14	6	1	1	2	1	420	64	484	0.56	1.53
(%)	(60.5)	(18.3)	(8.8)	(6.4)	(3.3)	(1.4)	(0.2)	(0.2)	(0.5)	(0.2)	(100.0)				
急速進行性糸球体腎炎	170	61	27	28	16	3	3	1	1	1	311	42	353	0.67	1.61
(%)	(54.7)	(19.6)	(8.7)	(9.0)	(5.1)	(1.0)	(1.0)	(0.3)	(0.3)	(0.3)	(100.0)				
妊娠腎/妊娠中毒症	209	61	27	18	10	6					331	70	401	0.41	0.84
(%)	(63.1)	(18.4)	(8.2)	(5.4)	(3.0)	(1.8)					(100.0)				
その他の分類不能の腎炎	139	28	17	9	11	2					206	55	261	0.41	0.83
(%)	(67.5)	(13.6)	(8.3)	(4.4)	(5.3)	(1.0)					(100.0)				
多発性嚢胞腎	906	325	192	112	62	20	12	3	6	2	1,640	240	1,880	0.58	1.45
(%)	(55.2)	(19.8)	(11.7)	(6.8)	(3.8)	(1.2)	(0.7)	(0.2)	(0.4)	(0.1)	(100.0)				
腎硬化症	1,856	660	391	226	107	38	9	11	8	6	3,312	508	3,820	0.55	1.54
(%)	(56.0)	(19.9)	(11.8)	(6.8)	(3.2)	(1.1)	(0.3)	(0.3)	(0.2)	(0.2)	(100.0)				
悪性高血圧	201	76	29	16	9	4	2		1		338	53	391	0.44	1.06
(%)	(59.5)	(22.5)	(8.6)	(4.7)	(2.7)	(1.2)	(0.6)		(0.3)		(100.0)				
糖尿病性腎症	8,266	2,931	1,499	959	535	192	94	42	55	35	14,608	2,352	16,960	0.60	1.80
(%)	(56.6)	(20.1)	(10.3)	(6.6)	(3.7)	(1.3)	(0.6)	(0.3)	(0.4)	(0.2)	(100.0)				
SLE腎炎	211	76	29	27	12	2	1	1	2	1	362	64	426	0.56	1.83
(%)	(58.3)	(21.0)	(8.0)	(7.5)	(3.3)	(0.6)	(0.3)	(0.3)	(0.6)	(0.3)	(100.0)				
アミロイド腎	34	14	4	6	4	2		1			65	9	74	0.73	1.47
(%)	(52.3)	(21.5)	(6.2)	(9.2)	(6.2)	(3.1)		(1.5)			(100.0)				
痛風腎	69	41	18	9	1	1	2	2			143	23	166	0.58	1.39
(%)	(48.3)	(28.7)	(12.6)	(6.3)	(0.7)	(0.7)	(1.4)	(1.4)			(100.0)				
先天性代謝異常による腎不全	26	9	5	2				1			43	6	49	0.44	1.36
(%)	(60.5)	(20.9)	(11.6)	(4.7)				(2.3)			(100.0)				
腎・尿路結核	14	4	2							1	21	10	31	1.04	3.88
(%)	(66.7)	(19.0)	(9.5)							(4.8)	(100.0)				
腎・尿路結石	36	12	8	7	2	2		1		1	69	10	79	0.95	2.53
(%)	(52.2)	(17.4)	(11.6)	(10.1)	(2.9)	(2.9)		(1.4)		(1.4)	(100.0)				
腎・尿路腫瘍	58	27	9	1	5	2	1	2			105	14	119	0.67	1.63
(%)	(55.2)	(25.7)	(8.6)	(1.0)	(4.8)	(1.9)	(1.0)	(1.9)			(100.0)				
閉塞性尿路障害	55	17	7	6	2	1			2		90	14	104	0.62	1.83
(%)	(61.1)	(18.9)	(7.8)	(6.7)	(2.2)	(1.1)			(2.2)		(100.0)				
骨髄腫	15	5		6	2	2					30	3	33	0.84	1.29
(%)	(50.0)	(16.7)		(20.0)	(6.7)	(6.7)					(100.0)				
腎形成不全	73	24	8	5	1	1					112	24	136	0.29	0.59
(%)	(65.2)	(21.4)	(7.1)	(4.5)	(0.9)	(0.9)					(100.0)				
不明	2,359	773	404	224	117	48	26	17	12	6	3,986	717	4,703	0.54	1.47
(%)	(59.2)	(19.4)	(10.1)	(5.6)	(2.9)	(1.2)	(0.7)	(0.4)	(0.3)	(0.2)	(100.0)				
再導入	237	88	40	27	16	6	1	1		1	417	66	483	0.53	1.44
(%)	(56.8)	(21.1)	(9.6)	(6.5)	(3.8)	(1.4)	(0.2)	(0.2)		(0.2)	(100.0)				
その他	597	207	93	51	30	15	9	4	3	4	1,013	147	1,160	0.60	1.75
(%)	(58.9)	(20.4)	(9.2)	(5.0)	(3.0)	(1.5)	(0.9)	(0.4)	(0.3)	(0.4)	(100.0)				
合計	24,925	8,251	4,130	2,562	1,390	509	239	129	120	85	42,340	7,235	49,575	0.55	1.60
(%)	(58.9)	(19.5)	(9.8)	(6.1)	(3.3)	(1.2)	(0.6)	(0.3)	(0.3)	(0.2)	(100.0)				
記載なし	3			2		1					6		6	1.40	1.80
(%)	(50.0)			(33.3)		(16.7)					(100.0)				
総計	24,928	8,251	4,130	2,564	1,390	510	239	129	120	85	42,346	7,235	49,581	0.55	1.60
(%)	(58.9)	(19.5)	(9.8)	(6.1)	(3.3)	(1.2)	(0.6)	(0.3)	(0.3)	(0.2)	(100.0)				

数値下のかつこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2015年調査