

表 2849 CRP濃度(mg/dL) 透析歴(年)別 (腹膜透析患者)

透析歴	0.2未満	0.2～	0.5～	1.0～	2.0～	4.0～	6.0～	8.0～	10.0～	15.0～	合計	記載なし	総計	平均	標準偏差
2年未満	1,224	428	202	135	91	30	13	7	7	3	2,140	1,540	3,680	0.58	1.47
(%)	(57.2)	(20.0)	(9.4)	(6.3)	(4.3)	(1.4)	(0.6)	(0.3)	(0.3)	(0.1)	(100.0)				
2年～	1,022	371	193	110	98	27	9	6	9	3	1,848	1,254	3,102	0.62	1.52
(%)	(55.3)	(20.1)	(10.4)	(6.0)	(5.3)	(1.5)	(0.5)	(0.3)	(0.5)	(0.2)	(100.0)				
5年～	546	177	107	63	25	15	8	1	4	1	947	573	1,520	0.58	1.61
(%)	(57.7)	(18.7)	(11.3)	(6.7)	(2.6)	(1.6)	(0.8)	(0.1)	(0.4)	(0.1)	(100.0)				
10年～	116	53	22	17	8	5	2			2	225	89	314	0.70	1.95
(%)	(51.6)	(23.6)	(9.8)	(7.6)	(3.6)	(2.2)	(0.9)			(0.9)	(100.0)				
15年～	33	9	8	5	1	2		1			59	28	87	0.66	1.42
(%)	(55.9)	(15.3)	(13.6)	(8.5)	(1.7)	(3.4)		(1.7)			(100.0)				
20年～	6	1	5								12	14	26	0.36	0.34
(%)	(50.0)	(8.3)	(41.7)								(100.0)				
25年～	2	1	2	1	1						7	5	12	0.88	0.93
(%)	(28.6)	(14.3)	(28.6)	(14.3)	(14.3)						(100.0)				
30年～	4										4	2	6	0.11	0.06
(%)	(100.0)										(100.0)				
35年～	1									1	2		2	12.58	17.74
(%)	(50.0)									(50.0)	(100.0)				
40年～												2	2		
(%)															
合計	2,954	1,040	539	331	224	79	32	15	20	10	5,244	3,507	8,751	0.61	1.57
(%)	(56.3)	(19.8)	(10.3)	(6.3)	(4.3)	(1.5)	(0.6)	(0.3)	(0.4)	(0.2)	(100.0)				
不明	2	1	1								4	14	18	0.30	0.33
(%)	(50.0)	(25.0)	(25.0)								(100.0)				
記載なし															
(%)															
総計	2,956	1,041	540	331	224	79	32	15	20	10	5,248	3,521	8,769	0.61	1.57
(%)	(56.3)	(19.8)	(10.3)	(6.3)	(4.3)	(1.5)	(0.6)	(0.3)	(0.4)	(0.2)	(100.0)				
平均	3.09	3.03	3.41	3.28	2.65	3.44	3.19	2.80	2.70	7.30	3.12	2.81	2.99		
標準偏差	3.59	3.43	3.94	3.67	3.09	3.81	3.28	4.16	2.39	10.53	3.61	3.46	3.55		

数値下のカッコ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査