

表 2850 CRP濃度(mg/dL) 原疾患別（腹膜透析患者）

原疾患	0.2未満	0.2～	0.5～	1.0～	2.0～	4.0～	6.0～	8.0～	10.0～	15.0～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	1,036	313	149	85	59	23	11	4	3	2	1,685	1,145	2,830	0.51	1.40
(%)	(61.5)	(18.6)	(8.8)	(5.0)	(3.5)	(1.4)	(0.7)	(0.2)	(0.2)	(0.1)	(100.0)				
慢性腎盂腎炎	33	13	5	5	2	2		1			61	36	97	0.68	1.46
(%)	(54.1)	(21.3)	(8.2)	(8.2)	(3.3)	(3.3)		(1.6)			(100.0)				
急速進行性糸球体腎炎	26	3	4	2	2	2					39	27	66	0.64	1.25
(%)	(66.7)	(7.7)	(10.3)	(5.1)	(5.1)	(5.1)					(100.0)				
妊娠腎/妊娠中毒症	14	2		3	1		1				21	6	27	0.68	1.46
(%)	(66.7)	(9.5)		(14.3)	(4.8)		(4.8)				(100.0)				
その他の分類不能の腎炎	30	8	3	5					1		47	31	78	0.57	1.91
(%)	(63.8)	(17.0)	(6.4)	(10.6)					(2.1)		(100.0)				
多発性嚢胞腎	97	21	12	10	5	1					146	117	263	0.38	0.69
(%)	(66.4)	(14.4)	(8.2)	(6.8)	(3.4)	(0.7)					(100.0)				
腎硬化症	339	154	82	42	21	13	4		4		659	436	1,095	0.57	1.24
(%)	(51.4)	(23.4)	(12.4)	(6.4)	(3.2)	(2.0)	(0.6)		(0.6)		(100.0)				
悪性高血圧	38	14	9	3							64	43	107	0.26	0.32
(%)	(59.4)	(21.9)	(14.1)	(4.7)							(100.0)				
糖尿病性腎症	899	333	182	118	89	27	13	4	10	4	1,679	1,120	2,799	0.70	1.84
(%)	(53.5)	(19.8)	(10.8)	(7.0)	(5.3)	(1.6)	(0.8)	(0.2)	(0.6)	(0.2)	(100.0)				
SLE腎炎	16	9	4		2						31	15	46	0.38	0.68
(%)	(51.6)	(29.0)	(12.9)		(6.5)						(100.0)				
アミロイド腎	3	3		1	3						10	5	15	1.12	1.29
(%)	(30.0)	(30.0)		(10.0)	(30.0)						(100.0)				
痛風腎	12	4	4	1							21	11	32	0.32	0.44
(%)	(57.1)	(19.0)	(19.0)	(4.8)							(100.0)				
先天性代謝異常による腎不全	9	2			1	1					13	4	17	0.64	1.35
(%)	(69.2)	(15.4)			(7.7)	(7.7)					(100.0)				
腎・尿路結核	1										1	1	2		
(%)	(100.0)										(100.0)				
腎・尿路結石	9	2	4		1			1			17	4	21	0.86	2.16
(%)	(52.9)	(11.8)	(23.5)		(5.9)			(5.9)			(100.0)				
腎・尿路腫瘍	1	4	2	1							8	6	14	0.56	0.55
(%)	(12.5)	(50.0)	(25.0)	(12.5)							(100.0)				
閉塞性尿路障害	10	1	2			1					14	7	21	0.52	1.18
(%)	(71.4)	(7.1)	(14.3)			(7.1)					(100.0)				
骨髄腫	3		1								4	3	7	0.24	0.33
(%)	(75.0)		(25.0)								(100.0)				
腎形成不全	30	9	4	1		2	1				47	25	72	0.53	1.31
(%)	(63.8)	(19.1)	(8.5)	(2.1)		(4.3)	(2.1)				(100.0)				
不明	252	113	54	38	26	5	2	4	2	3	499	343	842	0.73	1.89
(%)	(50.5)	(22.6)	(10.8)	(7.6)	(5.2)	(1.0)	(0.4)	(0.8)	(0.4)	(0.6)	(100.0)				
再導入	12	10	4	4	2						32	23	55	0.61	0.85
(%)	(37.5)	(31.3)	(12.5)	(12.5)	(6.3)						(100.0)				
その他	86	23	15	12	10	2		1		1	150	113	263	0.66	1.60
(%)	(57.3)	(15.3)	(10.0)	(8.0)	(6.7)	(1.3)		(0.7)		(0.7)	(100.0)				
合計	2,956	1,041	540	331	224	79	32	15	20	10	5,248	3,521	8,769	0.61	1.57
(%)	(56.3)	(19.8)	(10.3)	(6.3)	(4.3)	(1.5)	(0.6)	(0.3)	(0.4)	(0.2)	(100.0)				
記載なし															
(%)															
総計	2,956	1,041	540	331	224	79	32	15	20	10	5,248	3,521	8,769	0.61	1.57
(%)	(56.3)	(19.8)	(10.3)	(6.3)	(4.3)	(1.5)	(0.6)	(0.3)	(0.4)	(0.2)	(100.0)				

数値下のカッコ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査