

表 2864 透析前補正カルシウム濃度(mg/dL) 治療方法(新分類)別 (透析患者全体)

治療方法 (新分類)	7.0未満	7.0～	7.5～	8.0～	8.4～	8.5～	9.0～	9.5～	10.0～	10.1～	10.5～	11.0～	11.5～	12.0～	合計	記載なし	総計	平均	標準偏差
施設血液透析	1,060	1,367	5,468	17,003	4,883	54,473	66,996	42,861	5,404	13,274	6,530	2,202	799	583	222,903	25,822	248,725	9.18	0.75
(%)	(0.5)	(0.6)	(2.5)	(7.6)	(2.2)	(24.4)	(30.1)	(19.2)	(2.4)	(6.0)	(2.9)	(1.0)	(0.4)	(0.3)	(100.0)				
Off-line HDF	25	30	109	275	87	1,004	1,401	1,028	132	356	207	78	25	16	4,773	559	5,332	9.29	0.80
(%)	(0.5)	(0.6)	(2.3)	(5.8)	(1.8)	(21.0)	(29.4)	(21.5)	(2.8)	(7.5)	(4.3)	(1.6)	(0.5)	(0.3)	(100.0)				
On-line HDF	147	229	993	3,069	803	9,861	12,251	8,200	975	2,486	1,147	359	123	99	40,742	3,785	44,527	9.18	0.73
(%)	(0.4)	(0.6)	(2.4)	(7.5)	(2.0)	(24.2)	(30.1)	(20.1)	(2.4)	(6.1)	(2.8)	(0.9)	(0.3)	(0.2)	(100.0)				
Push/Pull HDF	1	2	5	7	1	20	31	22	4	7	5		1	1	107	3	110	9.24	0.88
(%)	(0.9)	(1.9)	(4.7)	(6.5)	(0.9)	(18.7)	(29.0)	(20.6)	(3.7)	(6.5)	(4.7)		(0.9)	(0.9)	(100.0)				
バイオフィльтраーション		3	5	17	6	49	79	54	10	17	9			2	251	16	267	9.24	0.74
(%)		(1.2)	(2.0)	(6.8)	(2.4)	(19.5)	(31.5)	(21.5)	(4.0)	(6.8)	(3.6)			(0.8)	(100.0)				
IHDF	14	20	67	298	79	907	1,041	628	50	189	85	30	15	3	3,426	114	3,540	9.13	0.72
(%)	(0.4)	(0.6)	(2.0)	(8.7)	(2.3)	(26.5)	(30.4)	(18.3)	(1.5)	(5.5)	(2.5)	(0.9)	(0.4)	(0.1)	(100.0)				
血液濾過	1					7	3	1	1						13	3	16	8.78	1.02
(%)	(7.7)					(53.8)	(23.1)	(7.7)	(7.7)						(100.0)				
血液吸着透析 (リクセル等使用)	3	13	33	99	20	272	365	303	31	99	50	14	2	1	1,305	71	1,376	9.24	0.73
(%)	(0.2)	(1.0)	(2.5)	(7.6)	(1.5)	(20.8)	(28.0)	(23.2)	(2.4)	(7.6)	(3.8)	(1.1)	(0.2)	(0.1)	(100.0)				
在宅血液透析	7	5	14	31	12	103	122	83	14	25	17	4		1	438	117	555	9.13	0.87
(%)	(1.6)	(1.1)	(3.2)	(7.1)	(2.7)	(23.5)	(27.9)	(18.9)	(3.2)	(5.7)	(3.9)	(0.9)		(0.2)	(100.0)				
腹膜透析 (CAPD)	9	19	53	161	43	542	874	836	117	304	161	53	21	8	3,201	1,701	4,902	9.40	0.77
(%)	(0.3)	(0.6)	(1.7)	(5.0)	(1.3)	(16.9)	(27.3)	(26.1)	(3.7)	(9.5)	(5.0)	(1.7)	(0.7)	(0.2)	(100.0)				
腹膜透析 (APD)	14	14	56	110	30	353	549	432	61	148	89	36	16	10	1,918	857	2,775	9.32	0.85
(%)	(0.7)	(0.7)	(2.9)	(5.7)	(1.6)	(18.4)	(28.6)	(22.5)	(3.2)	(7.7)	(4.6)	(1.9)	(0.8)	(0.5)	(100.0)				
腹膜透析 (CCPD)	1	7	14	34	10	145	189	208	28	86	39	7	5	1	774	318	1,092	9.39	0.76
(%)	(0.1)	(0.9)	(1.8)	(4.4)	(1.3)	(18.7)	(24.4)	(26.9)	(3.6)	(11.1)	(5.0)	(0.9)	(0.6)	(0.1)	(100.0)				
合計	1,282	1,709	6,817	21,104	5,974	67,736	83,901	54,656	6,827	16,991	8,339	2,783	1,007	725	279,851	33,366	313,217	9.18	0.75
(%)	(0.5)	(0.6)	(2.4)	(7.5)	(2.1)	(24.2)	(30.0)	(19.5)	(2.4)	(6.1)	(3.0)	(1.0)	(0.4)	(0.3)	(100.0)				
記載なし																			
(%)																			
総計	1,282	1,709	6,817	21,104	5,974	67,736	83,901	54,656	6,827	16,991	8,339	2,783	1,007	725	279,851	33,366	313,217	9.18	0.75
(%)	(0.5)	(0.6)	(2.4)	(7.5)	(2.1)	(24.2)	(30.0)	(19.5)	(2.4)	(6.1)	(3.0)	(1.0)	(0.4)	(0.3)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

*腹膜透析 (CCPD) は、APDと手動のバック交換の両方を行うもの

(C)Japanese Society for Dialysis Therapy

2015年調査