

表 2869 透析前補正カルシウム濃度(mg/dL) 原疾患別（透析患者全体）

原疾患	7.0未満	7.0～	7.5～	8.0～	8.4～	8.5～	9.0～	9.5～	10.0～	10.1～	10.5～	11.0～	11.5～	12.0～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	328	469	1,910	5,599	1,632	18,211	24,172	18,008	2,269	6,069	3,112	1,000	355	242	83,376	9,971	93,347	9.25	0.77
(%)	(0.4)	(0.6)	(2.3)	(6.7)	(2.0)	(21.8)	(29.0)	(21.6)	(2.7)	(7.3)	(3.7)	(1.2)	(0.4)	(0.3)	(100.0)				
慢性腎盂腎炎	15	21	57	187	36	567	803	555	86	186	102	31	9	7	2,662	273	2,935	9.24	0.77
(%)	(0.6)	(0.8)	(2.1)	(7.0)	(1.4)	(21.3)	(30.2)	(20.8)	(3.2)	(7.0)	(3.8)	(1.2)	(0.3)	(0.3)	(100.0)				
急速進行性糸球体腎炎	9	8	46	135	55	500	717	421	62	146	49	31	5	5	2,189	289	2,478	9.22	0.71
(%)	(0.4)	(0.4)	(2.1)	(6.2)	(2.5)	(22.8)	(32.8)	(19.2)	(2.8)	(6.7)	(2.2)	(1.4)	(0.2)	(0.2)	(100.0)				
妊娠腎/妊娠中毒症	6	9	36	93	23	304	372	340	48	114	45	17	5	4	1,416	129	1,545	9.28	0.76
(%)	(0.4)	(0.6)	(2.5)	(6.6)	(1.6)	(21.5)	(26.3)	(24.0)	(3.4)	(8.1)	(3.2)	(1.2)	(0.4)	(0.3)	(100.0)				
その他の分類不能の腎炎	7	10	28	94	34	277	363	273	36	81	48	11	4	2	1,268	137	1,405	9.21	0.77
(%)	(0.6)	(0.8)	(2.2)	(7.4)	(2.7)	(21.8)	(28.6)	(21.5)	(2.8)	(6.4)	(3.8)	(0.9)	(0.3)	(0.2)	(100.0)				
多発性嚢胞腎	41	54	186	635	187	2,245	3,048	2,256	292	703	329	125	46	18	10,165	1,091	11,256	9.25	0.74
(%)	(0.4)	(0.5)	(1.8)	(6.2)	(1.8)	(22.1)	(30.0)	(22.2)	(2.9)	(6.9)	(3.2)	(1.2)	(0.5)	(0.2)	(100.0)				
腎硬化症	126	136	571	2,048	582	6,693	8,290	5,076	681	1,445	686	210	83	68	26,695	3,110	29,805	9.16	0.73
(%)	(0.5)	(0.5)	(2.1)	(7.7)	(2.2)	(25.1)	(31.1)	(19.0)	(2.6)	(5.4)	(2.6)	(0.8)	(0.3)	(0.3)	(100.0)				
悪性高血圧	9	11	50	158	52	584	699	458	67	145	61	20	6	5	2,325	293	2,618	9.20	0.71
(%)	(0.4)	(0.5)	(2.2)	(6.8)	(2.2)	(25.1)	(30.1)	(19.7)	(2.9)	(6.2)	(2.6)	(0.9)	(0.3)	(0.2)	(100.0)				
糖尿病性腎症	515	721	2,899	9,111	2,498	28,313	32,762	19,089	2,218	5,454	2,676	871	321	215	107,663	12,615	120,278	9.12	0.73
(%)	(0.5)	(0.7)	(2.7)	(8.5)	(2.3)	(26.3)	(30.4)	(17.7)	(2.1)	(5.1)	(2.5)	(0.8)	(0.3)	(0.2)	(100.0)				
SLE腎炎	7	10	50	126	39	429	586	440	59	136	66	27	7	5	1,987	240	2,227	9.25	0.75
(%)	(0.4)	(0.5)	(2.5)	(6.3)	(2.0)	(21.6)	(29.5)	(22.1)	(3.0)	(6.8)	(3.3)	(1.4)	(0.4)	(0.3)	(100.0)				
アミロイド腎	2	1	11	21	10	85	131	89	8	34	5	3	1	2	403	59	462	9.23	0.73
(%)	(0.5)	(0.2)	(2.7)	(5.2)	(2.5)	(21.1)	(32.5)	(22.1)	(2.0)	(8.4)	(1.2)	(0.7)	(0.2)	(0.5)	(100.0)				
痛風腎	8	7	32	71	18	204	276	195	37	69	34	12	5	3	971	104	1,075	9.22	0.81
(%)	(0.8)	(0.7)	(3.3)	(7.3)	(1.9)	(21.0)	(28.4)	(20.1)	(3.8)	(7.1)	(3.5)	(1.2)	(0.5)	(0.3)	(100.0)				
先天性代謝異常による腎不全	4	1	10	16	3	53	62	62	3	13	6	3		2	238	36	274	9.18	0.92
(%)	(1.7)	(0.4)	(4.2)	(6.7)	(1.3)	(22.3)	(26.1)	(26.1)	(1.3)	(5.5)	(2.5)	(1.3)		(0.8)	(100.0)				
腎・尿路結核		2	2	13	4	30	58	42	2	10	8		1		172	23	195	9.25	0.68
(%)		(1.2)	(1.2)	(7.6)	(2.3)	(17.4)	(33.7)	(24.4)	(1.2)	(5.8)	(4.7)		(0.6)		(100.0)				
腎・尿路結石	3	2	10	26	13	129	161	101	11	42	15	10	2	2	527	54	581	9.24	0.75
(%)	(0.6)	(0.4)	(1.9)	(4.9)	(2.5)	(24.5)	(30.6)	(19.2)	(2.1)	(8.0)	(2.8)	(1.9)	(0.4)	(0.4)	(100.0)				
腎・尿路腫瘍	8	6	16	68	22	200	236	157	23	43	29	5	4		817	93	910	9.14	0.79
(%)	(1.0)	(0.7)	(2.0)	(8.3)	(2.7)	(24.5)	(28.9)	(19.2)	(2.8)	(5.3)	(3.5)	(0.6)	(0.5)		(100.0)				
閉塞性尿路障害	5	4	12	50	12	152	193	140	15	32	19	8	6	2	650	80	730	9.22	0.77
(%)	(0.8)	(0.6)	(1.8)	(7.7)	(1.8)	(23.4)	(29.7)	(21.5)	(2.3)	(4.9)	(2.9)	(1.2)	(0.9)	(0.3)	(100.0)				
骨髄腫	7	2	5	22	9	69	66	36	6	8	9	4	2	6	251	40	291	9.15	1.04
(%)	(2.8)	(0.8)	(2.0)	(8.8)	(3.6)	(27.5)	(26.3)	(14.3)	(2.4)	(3.2)	(3.6)	(1.6)	(0.8)	(2.4)	(100.0)				
腎形成不全	3	5	20	54	14	115	167	114	12	44	27	8	2	4	589	71	660	9.22	0.84
(%)	(0.5)	(0.8)	(3.4)	(9.2)	(2.4)	(19.5)	(28.4)	(19.4)	(2.0)	(7.5)	(4.6)	(1.4)	(0.3)	(0.7)	(100.0)				
不明	137	167	636	1,970	531	6,431	8,010	4,975	653	1,624	767	272	104	104	26,381	3,516	29,897	9.18	0.77
(%)	(0.5)	(0.6)	(2.4)	(7.5)	(2.0)	(24.4)	(30.4)	(18.9)	(2.5)	(6.2)	(2.9)	(1.0)	(0.4)	(0.4)	(100.0)				
再導入	9	16	62	168	55	424	574	344	55	125	57	27	9	4	1,929	254	2,183	9.17	0.77
(%)	(0.5)	(0.8)	(3.2)	(8.7)	(2.9)	(22.0)	(29.8)	(17.8)	(2.9)	(6.5)	(3.0)	(1.4)	(0.5)	(0.2)	(100.0)				
その他	33	47	167	439	145	1,720	2,149	1,482	183	467	188	88	29	25	7,162	882	8,044	9.21	0.76
(%)	(0.5)	(0.7)	(2.3)	(6.1)	(2.0)	(24.0)	(30.0)	(20.7)	(2.6)	(6.5)	(2.6)	(1.2)	(0.4)	(0.3)	(100.0)				
合計	1,282	1,709	6,816	21,104	5,974	67,736	83,895	54,653	6,826	16,990	8,338	2,783	1,006	725	279,836	33,360	313,196	9.18	0.75
(%)	(0.5)	(0.6)	(2.4)	(7.5)	(2.1)	(24.2)	(30.0)	(19.5)	(2.4)	(6.1)	(3.0)	(1.0)	(0.4)	(0.3)	(100.0)				
記載なし			1			1	6	3	1	1	1		1		15	6	21	9.53	0.83
(%)			(6.7)			(6.7)	(40.0)	(20.0)	(6.7)	(6.7)	(6.7)		(6.7)		(100.0)				
総計	1,282	1,709	6,817	21,104	5,974	67,736	83,901	54,656	6,827	16,991	8,339	2,783	1,007	725	279,851	33,366	313,217	9.18	0.75
(%)	(0.5)	(0.6)	(2.4)	(7.5)	(2.1)	(24.2)	(30.0)	(19.5)	(2.4)	(6.1)	(3.0)	(1.0)	(0.4)	(0.3)	(100.0)				

数値下のかっこ内は行方向の合計に対する％です。

(C)Japanese Society for Dialysis Therapy

2015年調査