

表 2872 透析前補正カルシウム濃度(mg/dL) 透析前リン濃度(mg/dL)別 (透析患者全体)

透析前リン濃度	7.0未満	7.0～	7.5～	8.0～	8.4～	8.5～	9.0～	9.5～	10.0～	10.1～	10.5～	11.0～	11.5～	12.0～	合計	記載なし	総計	平均	標準偏差
3未満	88	45	169	489	157	2,014	3,019	2,344	370	1,003	582	263	127	108	10,778	182	10,960	9.41	0.91
(%)	(0.8)	(0.4)	(1.6)	(4.5)	(1.5)	(18.7)	(28.0)	(21.7)	(3.4)	(9.3)	(5.4)	(2.4)	(1.2)	(1.0)	(100.0)				
3.0～	51	48	214	810	246	3,037	4,088	2,655	385	906	464	170	64	48	13,186	171	13,357	9.26	0.76
(%)	(0.4)	(0.4)	(1.6)	(6.1)	(1.9)	(23.0)	(31.0)	(20.1)	(2.9)	(6.9)	(3.5)	(1.3)	(0.5)	(0.4)	(100.0)				
3.5～	77	106	417	1,437	437	5,475	7,274	4,656	546	1,355	653	216	90	68	22,807	294	23,101	9.22	0.73
(%)	(0.3)	(0.5)	(1.8)	(6.3)	(1.9)	(24.0)	(31.9)	(20.4)	(2.4)	(5.9)	(2.9)	(0.9)	(0.4)	(0.3)	(100.0)				
4.0～	119	135	608	2,250	700	8,580	10,888	6,796	793	1,900	939	328	110	90	34,236	455	34,691	9.20	0.72
(%)	(0.3)	(0.4)	(1.8)	(6.6)	(2.0)	(25.1)	(31.8)	(19.9)	(2.3)	(5.5)	(2.7)	(1.0)	(0.3)	(0.3)	(100.0)				
4.5～	126	188	843	2,893	839	10,370	12,871	7,961	963	2,391	1,079	312	123	95	41,054	483	41,537	9.18	0.71
(%)	(0.3)	(0.5)	(2.1)	(7.0)	(2.0)	(25.3)	(31.4)	(19.4)	(2.3)	(5.8)	(2.6)	(0.8)	(0.3)	(0.2)	(100.0)				
5.0～	143	212	953	3,181	926	10,631	13,062	8,402	979	2,474	1,213	339	128	79	42,722	476	43,198	9.18	0.72
(%)	(0.3)	(0.5)	(2.2)	(7.4)	(2.2)	(24.9)	(30.6)	(19.7)	(2.3)	(5.8)	(2.8)	(0.8)	(0.3)	(0.2)	(100.0)				
5.5～	131	213	891	2,882	798	9,174	11,132	7,277	857	2,252	1,034	363	113	73	37,190	422	37,612	9.18	0.73
(%)	(0.4)	(0.6)	(2.4)	(7.7)	(2.1)	(24.7)	(29.9)	(19.6)	(2.3)	(6.1)	(2.8)	(1.0)	(0.3)	(0.2)	(100.0)				
6.0～	28	38	166	521	131	1,496	1,951	1,236	149	370	183	61	16	13	6,359	70	6,429	9.17	0.74
(%)	(0.4)	(0.6)	(2.6)	(8.2)	(2.1)	(23.5)	(30.7)	(19.4)	(2.3)	(5.8)	(2.9)	(1.0)	(0.3)	(0.2)	(100.0)				
6.1～	99	161	614	1,866	488	5,186	6,167	4,202	565	1,315	669	223	60	44	21,659	325	21,984	9.16	0.75
(%)	(0.5)	(0.7)	(2.8)	(8.6)	(2.3)	(23.9)	(28.5)	(19.4)	(2.6)	(6.1)	(3.1)	(1.0)	(0.3)	(0.2)	(100.0)				
6.5～	98	151	589	1,634	493	4,638	5,310	3,623	484	1,161	568	176	60	29	19,014	265	19,279	9.15	0.76
(%)	(0.5)	(0.8)	(3.1)	(8.6)	(2.6)	(24.4)	(27.9)	(19.1)	(2.5)	(6.1)	(3.0)	(0.9)	(0.3)	(0.2)	(100.0)				
7.0～	78	112	437	1,128	293	2,827	3,188	2,200	296	731	358	125	39	29	11,841	185	12,026	9.13	0.79
(%)	(0.7)	(0.9)	(3.7)	(9.5)	(2.5)	(23.9)	(26.9)	(18.6)	(2.5)	(6.2)	(3.0)	(1.1)	(0.3)	(0.2)	(100.0)				
7.5～	57	96	315	719	164	1,692	1,960	1,298	173	423	252	84	21	17	7,271	134	7,405	9.11	0.82
(%)	(0.8)	(1.3)	(4.3)	(9.9)	(2.3)	(23.3)	(27.0)	(17.9)	(2.4)	(5.8)	(3.5)	(1.2)	(0.3)	(0.2)	(100.0)				
8.0～	47	59	188	463	131	993	1,170	754	102	286	129	43	24	11	4,400	83	4,483	9.09	0.85
(%)	(1.1)	(1.3)	(4.3)	(10.5)	(3.0)	(22.6)	(26.6)	(17.1)	(2.3)	(6.5)	(2.9)	(1.0)	(0.5)	(0.3)	(100.0)				
8.5～	42	44	137	291	62	612	642	470	56	160	75	32	12	5	2,640	56	2,696	9.04	0.89
(%)	(1.6)	(1.7)	(5.2)	(11.0)	(2.3)	(23.2)	(24.3)	(17.8)	(2.1)	(6.1)	(2.8)	(1.2)	(0.5)	(0.2)	(100.0)				
9.0～	88	98	267	512	106	951	1,069	672	90	225	115	40	13	9	4,255	106	4,361	8.97	0.91
(%)	(2.1)	(2.3)	(6.3)	(12.0)	(2.5)	(22.4)	(25.1)	(15.8)	(2.1)	(5.3)	(2.7)	(0.9)	(0.3)	(0.2)	(100.0)				
合計	1,272	1,706	6,808	21,076	5,971	67,676	83,791	54,546	6,808	16,952	8,313	2,775	1,000	718	279,412	3,707	283,119	9.18	0.75
(%)	(0.5)	(0.6)	(2.4)	(7.5)	(2.1)	(24.2)	(30.0)	(19.5)	(2.4)	(6.1)	(3.0)	(1.0)	(0.4)	(0.3)	(100.0)				
記載なし	10	3	9	28	3	60	110	110	19	39	26	8	7	7	439	29,659	30,098	9.40	1.06
(%)	(2.3)	(0.7)	(2.1)	(6.4)	(0.7)	(13.7)	(25.1)	(25.1)	(4.3)	(8.9)	(5.9)	(1.8)	(1.6)	(1.6)	(100.0)				
総計	1,282	1,709	6,817	21,104	5,974	67,736	83,901	54,656	6,827	16,991	8,339	2,783	1,007	725	279,851	33,366	313,217	9.18	0.75
(%)	(0.5)	(0.6)	(2.4)	(7.5)	(2.1)	(24.2)	(30.0)	(19.5)	(2.4)	(6.1)	(3.0)	(1.0)	(0.4)	(0.3)	(100.0)				
平均	5.76	5.91	5.73	5.50	5.39	5.26	5.19	5.19	5.18	5.18	5.16	5.09	4.89	4.69	5.25	5.37	5.25		
標準偏差	2.01	1.81	1.65	1.50	1.44	1.39	1.38	1.42	1.46	1.48	1.56	1.63	1.71	1.76	1.44	1.64	1.44		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査