

表 2881 透析前補正カルシウム濃度(mg/dL) 性別（施設血液透析患者 透析回数週3回）

性別	7.0未満	7.0～	7.5～	8.0～	8.4～	8.5～	9.0～	9.5～	10.0～	10.1～	10.5～	11.0～	11.5～	12.0～	合計	記載なし	総計	平均	標準偏差
男性	664	863	3,740	11,734	3,368	35,944	41,342	24,839	3,102	7,407	3,610	1,182	399	282	138,476	5,964	144,440	9.13	0.74
(%)	(0.5)	(0.6)	(2.7)	(8.5)	(2.4)	(26.0)	(29.9)	(17.9)	(2.2)	(5.3)	(2.6)	(0.9)	(0.3)	(0.2)	(100.0)				
女性	334	434	1,499	4,588	1,305	16,343	23,179	16,567	2,136	5,467	2,721	948	365	274	76,160	3,252	79,412	9.27	0.76
(%)	(0.4)	(0.6)	(2.0)	(6.0)	(1.7)	(21.5)	(30.4)	(21.8)	(2.8)	(7.2)	(3.6)	(1.2)	(0.5)	(0.4)	(100.0)				
合計	998	1,297	5,239	16,322	4,673	52,287	64,521	41,406	5,238	12,874	6,331	2,130	764	556	214,636	9,216	223,852	9.18	0.75
(%)	(0.5)	(0.6)	(2.4)	(7.6)	(2.2)	(24.4)	(30.1)	(19.3)	(2.4)	(6.0)	(2.9)	(1.0)	(0.4)	(0.3)	(100.0)				
記載なし																			
(%)																			
総計	998	1,297	5,239	16,322	4,673	52,287	64,521	41,406	5,238	12,874	6,331	2,130	764	556	214,636	9,216	223,852	9.18	0.75
(%)	(0.5)	(0.6)	(2.4)	(7.6)	(2.2)	(24.4)	(30.1)	(19.3)	(2.4)	(6.0)	(2.9)	(1.0)	(0.4)	(0.3)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2015年調査