

表 2883 透析前補正カルシウム濃度(mg/dL) 透析歴(年)別 (施設血液透析患者 透析回数週3回)

透析歴	7.0未満	7.0～	7.5～	8.0～	8.4～	8.5～	9.0～	9.5～	10.0～	10.1～	10.5～	11.0～	11.5～	12.0～	合計	記載なし	総計	平均	標準偏差
2年未満	297	450	1,723	5,299	1,437	14,741	14,180	6,667	727	1,596	772	288	104	96	48,377	2,493	50,870	8.98	0.72
(%)	(0.6)	(0.9)	(3.6)	(11.0)	(3.0)	(30.5)	(29.3)	(13.8)	(1.5)	(3.3)	(1.6)	(0.6)	(0.2)	(0.2)	(100.0)				
2年～	220	289	1,232	4,165	1,261	14,506	17,573	9,951	1,121	2,634	1,201	371	129	101	54,754	2,101	56,855	9.13	0.70
(%)	(0.4)	(0.5)	(2.3)	(7.6)	(2.3)	(26.5)	(32.1)	(18.2)	(2.0)	(4.8)	(2.2)	(0.7)	(0.2)	(0.2)	(100.0)				
5年～	206	249	1,057	3,386	1,021	12,031	16,971	11,739	1,561	3,790	1,768	555	214	146	54,694	2,129	56,823	9.25	0.74
(%)	(0.4)	(0.5)	(1.9)	(6.2)	(1.9)	(22.0)	(31.0)	(21.5)	(2.9)	(6.9)	(3.2)	(1.0)	(0.4)	(0.3)	(100.0)				
10年～	127	116	528	1,547	443	5,465	7,962	6,395	908	2,363	1,250	417	154	96	27,771	1,122	28,893	9.33	0.79
(%)	(0.5)	(0.4)	(1.9)	(5.6)	(1.6)	(19.7)	(28.7)	(23.0)	(3.3)	(8.5)	(4.5)	(1.5)	(0.6)	(0.3)	(100.0)				
15年～	62	70	304	905	248	2,714	3,908	3,293	495	1,256	674	256	83	65	14,333	609	14,942	9.34	0.81
(%)	(0.4)	(0.5)	(2.1)	(6.3)	(1.7)	(18.9)	(27.3)	(23.0)	(3.5)	(8.8)	(4.7)	(1.8)	(0.6)	(0.5)	(100.0)				
20年～	49	55	184	481	146	1,470	1,962	1,729	220	651	328	126	48	24	7,473	305	7,778	9.30	0.82
(%)	(0.7)	(0.7)	(2.5)	(6.4)	(2.0)	(19.7)	(26.3)	(23.1)	(2.9)	(8.7)	(4.4)	(1.7)	(0.6)	(0.3)	(100.0)				
25年～	16	38	99	291	60	728	1,014	876	110	323	176	58	14	15	3,818	166	3,984	9.29	0.81
(%)	(0.4)	(1.0)	(2.6)	(7.6)	(1.6)	(19.1)	(26.6)	(22.9)	(2.9)	(8.5)	(4.6)	(1.5)	(0.4)	(0.4)	(100.0)				
30年～	11	14	60	154	38	363	583	446	52	157	94	25	10	4	2,011	103	2,114	9.27	0.80
(%)	(0.5)	(0.7)	(3.0)	(7.7)	(1.9)	(18.1)	(29.0)	(22.2)	(2.6)	(7.8)	(4.7)	(1.2)	(0.5)	(0.2)	(100.0)				
35年～	8	14	40	69	17	189	249	232	30	84	42	24	7	7	1,012	58	1,070	9.30	0.88
(%)	(0.8)	(1.4)	(4.0)	(6.8)	(1.7)	(18.7)	(24.6)	(22.9)	(3.0)	(8.3)	(4.2)	(2.4)	(0.7)	(0.7)	(100.0)				
40年～	2	2	9	16	1	55	80	56	7	11	21	8	1	1	270	15	285	9.32	0.86
(%)	(0.7)	(0.7)	(3.3)	(5.9)	(0.4)	(20.4)	(29.6)	(20.7)	(2.6)	(4.1)	(7.8)	(3.0)	(0.4)	(0.4)	(100.0)				
合計	998	1,297	5,236	16,313	4,672	52,262	64,482	41,384	5,231	12,865	6,326	2,128	764	555	214,513	9,101	223,614	9.18	0.75
(%)	(0.5)	(0.6)	(2.4)	(7.6)	(2.2)	(24.4)	(30.1)	(19.3)	(2.4)	(6.0)	(2.9)	(1.0)	(0.4)	(0.3)	(100.0)				
不明			3	9	1	25	39	22	7	9	5	2		1	123	114	237	9.30	0.74
(%)			(2.4)	(7.3)	(0.8)	(20.3)	(31.7)	(17.9)	(5.7)	(7.3)	(4.1)	(1.6)		(0.8)	(100.0)				
記載なし																1	1		
(%)																			
総計	998	1,297	5,239	16,322	4,673	52,287	64,521	41,406	5,238	12,874	6,331	2,130	764	556	214,636	9,216	223,852	9.18	0.75
(%)	(0.5)	(0.6)	(2.4)	(7.6)	(2.2)	(24.4)	(30.1)	(19.3)	(2.4)	(6.0)	(2.9)	(1.0)	(0.4)	(0.3)	(100.0)				
平均	6.85	6.50	6.29	5.92	5.78	5.95	6.69	7.95	8.43	8.94	9.39	9.65	9.45	8.84	6.97	6.80	6.96		
標準偏差	7.70	8.10	7.61	7.06	6.70	6.66	6.82	7.25	7.20	7.39	7.70	7.96	7.62	7.84	7.09	7.44	7.10		

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2015年調査