

表 2884 透析前補正カルシウム濃度(mg/dL) 原疾患別（施設血液透析患者 透析回数週3回）

原疾患	7.0未満	7.0～	7.5～	8.0～	8.4～	8.5～	9.0～	9.5～	10.0～	10.1～	10.5～	11.0～	11.5～	12.0～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	255	331	1,384	4,089	1,222	13,297	17,673	13,025	1,660	4,457	2,264	746	258	179	60,840	2,634	63,474	9.25	0.77
(%)	(0.4)	(0.5)	(2.3)	(6.7)	(2.0)	(21.9)	(29.0)	(21.4)	(2.7)	(7.3)	(3.7)	(1.2)	(0.4)	(0.3)	(100.0)				
慢性腎盂腎炎	13	17	43	130	27	428	603	408	61	123	83	20	8	7	1,971	69	2,040	9.24	0.78
(%)	(0.7)	(0.9)	(2.2)	(6.6)	(1.4)	(21.7)	(30.6)	(20.7)	(3.1)	(6.2)	(4.2)	(1.0)	(0.4)	(0.4)	(100.0)				
急速進行性糸球体腎炎	6	6	32	105	46	396	564	311	50	106	40	25	4	4	1,695	76	1,771	9.21	0.71
(%)	(0.4)	(0.4)	(1.9)	(6.2)	(2.7)	(23.4)	(33.3)	(18.3)	(2.9)	(6.3)	(2.4)	(1.5)	(0.2)	(0.2)	(100.0)				
妊娠腎/妊娠中毒症	5	6	23	56	15	187	245	239	31	87	31	10	4	2	941	40	981	9.31	0.76
(%)	(0.5)	(0.6)	(2.4)	(6.0)	(1.6)	(19.9)	(26.0)	(25.4)	(3.3)	(9.2)	(3.3)	(1.1)	(0.4)	(0.2)	(100.0)				
その他の分類不能の腎炎	6	7	22	65	26	195	252	190	26	57	38	9	2		895	34	929	9.21	0.78
(%)	(0.7)	(0.8)	(2.5)	(7.3)	(2.9)	(21.8)	(28.2)	(21.2)	(2.9)	(6.4)	(4.2)	(1.0)	(0.2)		(100.0)				
多発性嚢胞腎	30	39	150	490	143	1,712	2,275	1,725	221	556	246	94	37	16	7,734	282	8,016	9.26	0.74
(%)	(0.4)	(0.5)	(1.9)	(6.3)	(1.8)	(22.1)	(29.4)	(22.3)	(2.9)	(7.2)	(3.2)	(1.2)	(0.5)	(0.2)	(100.0)				
腎硬化症	102	104	447	1,640	479	5,359	6,577	3,953	563	1,125	534	168	67	54	21,172	835	22,007	9.16	0.73
(%)	(0.5)	(0.5)	(2.1)	(7.7)	(2.3)	(25.3)	(31.1)	(18.7)	(2.7)	(5.3)	(2.5)	(0.8)	(0.3)	(0.3)	(100.0)				
悪性高血圧	5	8	37	124	39	451	550	348	52	100	45	14	6	4	1,783	78	1,861	9.19	0.70
(%)	(0.3)	(0.4)	(2.1)	(7.0)	(2.2)	(25.3)	(30.8)	(19.5)	(2.9)	(5.6)	(2.5)	(0.8)	(0.3)	(0.2)	(100.0)				
糖尿病性腎症	413	575	2,310	7,299	2,009	22,669	26,175	15,063	1,783	4,305	2,119	698	250	172	85,840	3,586	89,426	9.12	0.73
(%)	(0.5)	(0.7)	(2.7)	(8.5)	(2.3)	(26.4)	(30.5)	(17.5)	(2.1)	(5.0)	(2.5)	(0.8)	(0.3)	(0.2)	(100.0)				
SLE腎炎	5	10	36	94	27	302	432	325	39	97	43	18	4	3	1,435	69	1,504	9.24	0.75
(%)	(0.3)	(0.7)	(2.5)	(6.6)	(1.9)	(21.0)	(30.1)	(22.6)	(2.7)	(6.8)	(3.0)	(1.3)	(0.3)	(0.2)	(100.0)				
アミロイド腎	2	1	11	17	7	64	96	65	7	23	3	2	1	1	300	14	314	9.18	0.75
(%)	(0.7)	(0.3)	(3.7)	(5.7)	(2.3)	(21.3)	(32.0)	(21.7)	(2.3)	(7.7)	(1.0)	(0.7)	(0.3)	(0.3)	(100.0)				
痛風腎	5	4	26	60	15	162	213	138	31	51	27	11	4	3	750	28	778	9.22	0.82
(%)	(0.7)	(0.5)	(3.5)	(8.0)	(2.0)	(21.6)	(28.4)	(18.4)	(4.1)	(6.8)	(3.6)	(1.5)	(0.5)	(0.4)	(100.0)				
先天性代謝異常による腎不全	2	1	8	12	3	32	46	44	3	9	5	3		1	169	4	173	9.21	0.87
(%)	(1.2)	(0.6)	(4.7)	(7.1)	(1.8)	(18.9)	(27.2)	(26.0)	(1.8)	(5.3)	(3.0)	(1.8)		(0.6)	(100.0)				
腎・尿路結核		1	2	10	2	24	44	33	1	9	7		1		134	7	141	9.28	0.69
(%)		(0.7)	(1.5)	(7.5)	(1.5)	(17.9)	(32.8)	(24.6)	(0.7)	(6.7)	(5.2)		(0.7)		(100.0)				
腎・尿路結石	3	2	7	24	11	101	124	76	7	32	13	7	2	2	411	10	421	9.22	0.77
(%)	(0.7)	(0.5)	(1.7)	(5.8)	(2.7)	(24.6)	(30.2)	(18.5)	(1.7)	(7.8)	(3.2)	(1.7)	(0.5)	(0.5)	(100.0)				
腎・尿路腫瘍	7	5	13	56	20	161	188	116	22	36	26	4	4		658	20	678	9.14	0.81
(%)	(1.1)	(0.8)	(2.0)	(8.5)	(3.0)	(24.5)	(28.6)	(17.6)	(3.3)	(5.5)	(4.0)	(0.6)	(0.6)		(100.0)				
閉塞性尿路障害	2	2	9	33	11	112	146	98	9	23	15	5	5	1	471	23	494	9.22	0.73
(%)	(0.4)	(0.4)	(1.9)	(7.0)	(2.3)	(23.8)	(31.0)	(20.8)	(1.9)	(4.9)	(3.2)	(1.1)	(1.1)	(0.2)	(100.0)				
骨髄腫	4		4	13	7	54	50	24	5	4	6	3	1	5	180	10	190	9.18	1.01
(%)	(2.2)		(2.2)	(7.2)	(3.9)	(30.0)	(27.8)	(13.3)	(2.8)	(2.2)	(3.3)	(1.7)	(0.6)	(2.8)	(100.0)				
腎形成不全	2	4	14	35	10	74	116	80	10	25	12	4		2	388	12	400	9.17	0.79
(%)	(0.5)	(1.0)	(3.6)	(9.0)	(2.6)	(19.1)	(29.9)	(20.6)	(2.6)	(6.4)	(3.1)	(1.0)		(0.5)	(100.0)				
不明	100	122	493	1,519	403	4,890	6,127	3,758	478	1,220	595	209	79	81	20,074	1,102	21,176	9.18	0.77
(%)	(0.5)	(0.6)	(2.5)	(7.6)	(2.0)	(24.4)	(30.5)	(18.7)	(2.4)	(6.1)	(3.0)	(1.0)	(0.4)	(0.4)	(100.0)				
再導入	5	11	42	106	42	292	384	246	37	85	38	19	5	3	1,315	41	1,356	9.18	0.77
(%)	(0.4)	(0.8)	(3.2)	(8.1)	(3.2)	(22.2)	(29.2)	(18.7)	(2.8)	(6.5)	(2.9)	(1.4)	(0.4)	(0.2)	(100.0)				
その他	26	41	126	345	109	1,325	1,637	1,141	142	343	141	61	21	16	5,474	241	5,715	9.20	0.75
(%)	(0.5)	(0.7)	(2.3)	(6.3)	(2.0)	(24.2)	(29.9)	(20.8)	(2.6)	(6.3)	(2.6)	(1.1)	(0.4)	(0.3)	(100.0)				
合計	998	1,297	5,239	16,322	4,673	52,287	64,517	41,406	5,238	12,873	6,331	2,130	763	556	214,630	9,215	223,845	9.18	0.75
(%)	(0.5)	(0.6)	(2.4)	(7.6)	(2.2)	(24.4)	(30.1)	(19.3)	(2.4)	(6.0)	(2.9)	(1.0)	(0.4)	(0.3)	(100.0)				
記載なし							4			1			1		6	1	7	9.72	0.97
(%)							(66.7)			(16.7)			(16.7)		(100.0)				
総計	998	1,297	5,239	16,322	4,673	52,287	64,521	41,406	5,238	12,874	6,331	2,130	764	556	214,636	9,216	223,852	9.18	0.75
(%)	(0.5)	(0.6)	(2.4)	(7.6)	(2.2)	(24.4)	(30.1)	(19.3)	(2.4)	(6.0)	(2.9)	(1.0)	(0.4)	(0.3)	(100.0)				

数値下のかっこ内は行方向の合計に対する％です。
(C)Japanese Society for Dialysis Therapy
2015年調査