

表 2896 透析前補正カルシウム濃度(mg/dL) 性別（血液透析濾過患者 透析回数週3回）

性別	7.0未満	7.0～	7.5～	8.0～	8.4～	8.5～	9.0～	9.5～	10.0～	10.1～	10.5～	11.0～	11.5～	12.0～	合計	記載なし	総計	平均	標準偏差
男性	120	192	819	2,495	628	7,776	9,187	5,917	664	1,716	814	252	81	61	30,722	908	31,630	9.15	0.73
(%)	(0.4)	(0.6)	(2.7)	(8.1)	(2.0)	(25.3)	(29.9)	(19.3)	(2.2)	(5.6)	(2.6)	(0.8)	(0.3)	(0.2)	(100.0)				
女性	61	86	336	1,076	325	3,776	5,280	3,818	478	1,261	611	204	78	53	17,443	508	17,951	9.27	0.74
(%)	(0.3)	(0.5)	(1.9)	(6.2)	(1.9)	(21.6)	(30.3)	(21.9)	(2.7)	(7.2)	(3.5)	(1.2)	(0.4)	(0.3)	(100.0)				
合計	181	278	1,155	3,571	953	11,552	14,467	9,735	1,142	2,977	1,425	456	159	114	48,165	1,416	49,581	9.19	0.74
(%)	(0.4)	(0.6)	(2.4)	(7.4)	(2.0)	(24.0)	(30.0)	(20.2)	(2.4)	(6.2)	(3.0)	(0.9)	(0.3)	(0.2)	(100.0)				
記載なし																			
(%)																			
総計	181	278	1,155	3,571	953	11,552	14,467	9,735	1,142	2,977	1,425	456	159	114	48,165	1,416	49,581	9.19	0.74
(%)	(0.4)	(0.6)	(2.4)	(7.4)	(2.0)	(24.0)	(30.0)	(20.2)	(2.4)	(6.2)	(3.0)	(0.9)	(0.3)	(0.2)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2015年調査