

表 2898 透析前補正カルシウム濃度(mg/dL) 透析歴(年)別（血液透析濾過患者 透析回数週3回）

透析歴	7.0未満	7.0～	7.5～	8.0～	8.4～	8.5～	9.0～	9.5～	10.0～	10.1～	10.5～	11.0～	11.5～	12.0～	合計	記載なし	総計	平均	標準偏差
2年未満	28	62	220	650	165	1,933	1,828	859	77	209	100	34	13	17	6,195	210	6,405	8.99	0.70
(%)	(0.5)	(1.0)	(3.6)	(10.5)	(2.7)	(31.2)	(29.5)	(13.9)	(1.2)	(3.4)	(1.6)	(0.5)	(0.2)	(0.3)	(100.0)				
2年～	39	47	254	834	219	2,872	3,425	2,050	217	506	220	67	26	19	10,795	274	11,069	9.13	0.68
(%)	(0.4)	(0.4)	(2.4)	(7.7)	(2.0)	(26.6)	(31.7)	(19.0)	(2.0)	(4.7)	(2.0)	(0.6)	(0.2)	(0.2)	(100.0)				
5年～	44	61	233	779	218	2,890	3,853	2,693	304	770	382	83	36	21	12,367	407	12,774	9.22	0.71
(%)	(0.4)	(0.5)	(1.9)	(6.3)	(1.8)	(23.4)	(31.2)	(21.8)	(2.5)	(6.2)	(3.1)	(0.7)	(0.3)	(0.2)	(100.0)				
10年～	24	35	160	515	123	1,467	2,115	1,668	201	603	293	108	32	24	7,368	195	7,563	9.29	0.77
(%)	(0.3)	(0.5)	(2.2)	(7.0)	(1.7)	(19.9)	(28.7)	(22.6)	(2.7)	(8.2)	(4.0)	(1.5)	(0.4)	(0.3)	(100.0)				
15年～	16	22	106	299	87	953	1,304	999	139	383	165	70	20	17	4,580	116	4,696	9.29	0.77
(%)	(0.3)	(0.5)	(2.3)	(6.5)	(1.9)	(20.8)	(28.5)	(21.8)	(3.0)	(8.4)	(3.6)	(1.5)	(0.4)	(0.4)	(100.0)				
20年～	13	24	76	205	58	615	853	654	86	213	108	44	15	8	2,972	86	3,058	9.25	0.79
(%)	(0.4)	(0.8)	(2.6)	(6.9)	(2.0)	(20.7)	(28.7)	(22.0)	(2.9)	(7.2)	(3.6)	(1.5)	(0.5)	(0.3)	(100.0)				
25年～	12	11	50	140	33	394	565	388	47	133	70	20	8	2	1,873	57	1,930	9.22	0.76
(%)	(0.6)	(0.6)	(2.7)	(7.5)	(1.8)	(21.0)	(30.2)	(20.7)	(2.5)	(7.1)	(3.7)	(1.1)	(0.4)	(0.1)	(100.0)				
30年～	4	12	31	78	28	249	303	239	36	95	49	17	6	3	1,150	47	1,197	9.26	0.80
(%)	(0.3)	(1.0)	(2.7)	(6.8)	(2.4)	(21.7)	(26.3)	(20.8)	(3.1)	(8.3)	(4.3)	(1.5)	(0.5)	(0.3)	(100.0)				
35年～	1	3	19	49	18	128	160	132	26	47	30	9	2	2	626	12	638	9.27	0.78
(%)	(0.2)	(0.5)	(3.0)	(7.8)	(2.9)	(20.4)	(25.6)	(21.1)	(4.2)	(7.5)	(4.8)	(1.4)	(0.3)	(0.3)	(100.0)				
40年～		1	6	18	4	37	46	46	8	16	8	4		1	195	2	197	9.30	0.79
(%)		(0.5)	(3.1)	(9.2)	(2.1)	(19.0)	(23.6)	(23.6)	(4.1)	(8.2)	(4.1)	(2.1)		(0.5)	(100.0)				
合計	181	278	1,155	3,567	953	11,538	14,452	9,728	1,141	2,975	1,425	456	158	114	48,121	1,406	49,527	9.19	0.74
(%)	(0.4)	(0.6)	(2.4)	(7.4)	(2.0)	(24.0)	(30.0)	(20.2)	(2.4)	(6.2)	(3.0)	(0.9)	(0.3)	(0.2)	(100.0)				
不明				4		14	15	7	1	2			1		44	10	54	9.17	0.63
(%)				(9.1)		(31.8)	(34.1)	(15.9)	(2.3)	(4.5)			(2.3)		(100.0)				
記載なし																			
(%)																			
総計	181	278	1,155	3,571	953	11,552	14,467	9,735	1,142	2,977	1,425	456	159	114	48,165	1,416	49,581	9.19	0.74
(%)	(0.4)	(0.6)	(2.4)	(7.4)	(2.0)	(24.0)	(30.0)	(20.2)	(2.4)	(6.2)	(3.0)	(0.9)	(0.3)	(0.2)	(100.0)				
平均	9.65	9.77	9.56	9.10	9.44	8.70	9.36	10.35	11.43	11.49	11.81	12.73	11.68	10.75	9.69	9.50	9.68		
標準偏差	8.73	9.44	9.37	8.95	9.24	8.51	8.48	8.62	9.14	8.82	9.03	9.13	8.67	8.90	8.71	8.72	8.71		

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2015年調査