

表 2899 透析前補正カルシウム濃度(mg/dL) 原疾患別（血液透析濾過患者 透析回数週3回）

原疾患	7.0未満	7.0～	7.5～	8.0～	8.4～	8.5～	9.0～	9.5～	10.0～	10.1～	10.5～	11.0～	11.5～	12.0～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	45	99	400	1,160	304	3,726	4,944	3,745	456	1,178	617	187	71	44	16,976	486	17,462	9.25	0.75
(%)	(0.3)	(0.6)	(2.4)	(6.8)	(1.8)	(21.9)	(29.1)	(22.1)	(2.7)	(6.9)	(3.6)	(1.1)	(0.4)	(0.3)	(100.0)				
慢性腎盂腎炎	1	2	9	44	6	95	135	108	15	40	14	7			476	8	484	9.24	0.72
(%)	(0.2)	(0.4)	(1.9)	(9.2)	(1.3)	(20.0)	(28.4)	(22.7)	(3.2)	(8.4)	(2.9)	(1.5)			(100.0)				
急速進行性糸球体腎炎	3	1	12	21	7	75	98	78	8	34	5	5	1	1	349	4	353	9.23	0.75
(%)	(0.9)	(0.3)	(3.4)	(6.0)	(2.0)	(21.5)	(28.1)	(22.3)	(2.3)	(9.7)	(1.4)	(1.4)	(0.3)	(0.3)	(100.0)				
妊娠腎/妊娠中毒症	1	1	10	27	6	96	103	86	12	22	11	6	1	1	383	18	401	9.23	0.73
(%)	(0.3)	(0.3)	(2.6)	(7.0)	(1.6)	(25.1)	(26.9)	(22.5)	(3.1)	(5.7)	(2.9)	(1.6)	(0.3)	(0.3)	(100.0)				
その他の分類不能の腎炎		2	4	18	7	61	75	53	6	19	7	1	1	2	256	5	261	9.22	0.73
(%)		(0.8)	(1.6)	(7.0)	(2.7)	(23.8)	(29.3)	(20.7)	(2.3)	(7.4)	(2.7)	(0.4)	(0.4)	(0.8)	(100.0)				
多発性嚢胞腎	10	9	27	116	32	399	593	389	49	117	66	22	5	1	1,835	45	1,880	9.24	0.74
(%)	(0.5)	(0.5)	(1.5)	(6.3)	(1.7)	(21.7)	(32.3)	(21.2)	(2.7)	(6.4)	(3.6)	(1.2)	(0.3)	(0.1)	(100.0)				
腎硬化症	8	22	89	277	65	923	1,174	754	77	206	90	17	7	11	3,720	100	3,820	9.17	0.69
(%)	(0.2)	(0.6)	(2.4)	(7.4)	(1.7)	(24.8)	(31.6)	(20.3)	(2.1)	(5.5)	(2.4)	(0.5)	(0.2)	(0.3)	(100.0)				
悪性高血圧	2	2	8	25	8	88	110	72	10	36	10	6		1	378	13	391	9.25	0.74
(%)	(0.5)	(0.5)	(2.1)	(6.6)	(2.1)	(23.3)	(29.1)	(19.0)	(2.6)	(9.5)	(2.6)	(1.6)		(0.3)	(100.0)				
糖尿病性腎症	71	97	422	1,372	378	4,342	5,024	3,014	318	858	400	126	42	28	16,492	468	16,960	9.12	0.72
(%)	(0.4)	(0.6)	(2.6)	(8.3)	(2.3)	(26.3)	(30.5)	(18.3)	(1.9)	(5.2)	(2.4)	(0.8)	(0.3)	(0.2)	(100.0)				
SLE腎炎	2		11	26	11	91	116	86	15	25	16	7	3	2	411	15	426	9.27	0.79
(%)	(0.5)		(2.7)	(6.3)	(2.7)	(22.1)	(28.2)	(20.9)	(3.6)	(6.1)	(3.9)	(1.7)	(0.7)	(0.5)	(100.0)				
アミロイド腎				3	2	17	26	12	1	8	1	1		1	72	2	74	9.34	0.68
(%)				(4.2)	(2.8)	(23.6)	(36.1)	(16.7)	(1.4)	(11.1)	(1.4)	(1.4)		(1.4)	(100.0)				
痛風腎	2	2	4	5	2	28	50	43	3	14	5	1	1		160	6	166	9.29	0.78
(%)	(1.3)	(1.3)	(2.5)	(3.1)	(1.3)	(17.5)	(31.3)	(26.9)	(1.9)	(8.8)	(3.1)	(0.6)	(0.6)		(100.0)				
先天性代謝異常による腎不全	1		2	3		19	12	9		1	1				48	1	49	8.97	0.70
(%)	(2.1)		(4.2)	(6.3)		(39.6)	(25.0)	(18.8)		(2.1)	(2.1)				(100.0)				
腎・尿路結核				1	2	5	12	7	1	1	1				30	1	31	9.24	0.55
(%)				(3.3)	(6.7)	(16.7)	(40.0)	(23.3)	(3.3)	(3.3)	(3.3)				(100.0)				
腎・尿路結石			1	2	2	19	22	15	4	7	1	3			76	3	79	9.33	0.69
(%)			(1.3)	(2.6)	(2.6)	(25.0)	(28.9)	(19.7)	(5.3)	(9.2)	(1.3)	(3.9)			(100.0)				
腎・尿路腫瘍	1	1	1	7	2	28	38	29		6	3	1			117	2	119	9.16	0.73
(%)	(0.9)	(0.9)	(0.9)	(6.0)	(1.7)	(23.9)	(32.5)	(24.8)		(5.1)	(2.6)	(0.9)			(100.0)				
閉塞性尿路障害	3	2	3	11		19	28	24	4	6	1	2	1		104		104	9.12	0.93
(%)	(2.9)	(1.9)	(2.9)	(10.6)		(18.3)	(26.9)	(23.1)	(3.8)	(5.8)	(1.0)	(1.9)	(1.0)		(100.0)				
骨髄腫			1	5		7	3	8		3	3	1			31	2	33	9.32	0.88
(%)			(3.2)	(16.1)		(22.6)	(9.7)	(25.8)		(9.7)	(9.7)	(3.2)			(100.0)				
腎形成不全	1	1	5	14	3	34	38	19	1	9	8				133	3	136	9.10	0.78
(%)	(0.8)	(0.8)	(3.8)	(10.5)	(2.3)	(25.6)	(28.6)	(14.3)	(0.8)	(6.8)	(6.0)				(100.0)				
不明	22	29	98	323	86	1,106	1,365	876	123	276	122	42	19	18	4,505	198	4,703	9.20	0.75
(%)	(0.5)	(0.6)	(2.2)	(7.2)	(1.9)	(24.6)	(30.3)	(19.4)	(2.7)	(6.1)	(2.7)	(0.9)	(0.4)	(0.4)	(100.0)				
再導入	4	4	16	49	9	96	150	76	13	32	12	7	2		470	13	483	9.14	0.78
(%)	(0.9)	(0.9)	(3.4)	(10.4)	(1.9)	(20.4)	(31.9)	(16.2)	(2.8)	(6.8)	(2.6)	(1.5)	(0.4)		(100.0)				
その他	4	4	32	62	21	278	350	229	25	79	30	14	5		1,137	23	1,160	9.22	0.74
(%)	(0.4)	(0.4)	(2.8)	(5.5)	(1.8)	(24.5)	(30.8)	(20.1)	(2.2)	(6.9)	(2.6)	(1.2)	(0.4)	(0.4)	(100.0)				
合計	181	278	1,155	3,571	953	11,552	14,466	9,732	1,141	2,977	1,424	456	159	114	48,159	1,416	49,575	9.19	0.74
(%)	(0.4)	(0.6)	(2.4)	(7.4)	(2.0)	(24.0)	(30.0)	(20.2)	(2.4)	(6.2)	(3.0)	(0.9)	(0.3)	(0.2)	(100.0)				
記載なし							1	3	1		1				6		6	9.80	0.42
(%)							(16.7)	(50.0)	(16.7)		(16.7)				(100.0)				
総計	181	278	1,155	3,571	953	11,552	14,467	9,735	1,142	2,977	1,425	456	159	114	48,165	1,416	49,581	9.19	0.74
(%)	(0.4)	(0.6)	(2.4)	(7.4)	(2.0)	(24.0)	(30.0)	(20.2)	(2.4)	(6.2)	(3.0)	(0.9)	(0.3)	(0.2)	(100.0)				

数値下のかっこ内は行方向の合計に対する％です。

(C)Japanese Society for Dialysis Therapy

2015年調査