

表 2914 補正カルシウム濃度(mg/dL) 原疾患別（腹膜透析患者）

原疾患	7.0未満	7.0～	7.5～	8.0～	8.4～	8.5～	9.0～	9.5～	10.0～	10.1～	10.5～	11.0～	11.5～	12.0～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	12	13	35	77	27	311	516	513	70	201	109	24	13	6	1,927	903	2,830	9.42	0.81
(%)	(0.6)	(0.7)	(1.8)	(4.0)	(1.4)	(16.1)	(26.8)	(26.6)	(3.6)	(10.4)	(5.7)	(1.2)	(0.7)	(0.3)	(100.0)				
慢性腎盂腎炎	1	1	1	3		8	22	20	3	9	1	2			71	26	97	9.34	0.96
(%)	(1.4)	(1.4)	(1.4)	(4.2)		(11.3)	(31.0)	(28.2)	(4.2)	(12.7)	(1.4)	(2.8)			(100.0)				
急速進行性糸球体腎炎			1	2	1	7	17	11	2	2	1				44	22	66	9.25	0.60
(%)			(2.3)	(4.5)	(2.3)	(15.9)	(38.6)	(25.0)	(4.5)	(4.5)	(2.3)				(100.0)				
妊娠腎/妊娠中毒症			1	2	1	3	7	4	2	1	3				24	3	27	9.30	0.79
(%)			(4.2)	(8.3)	(4.2)	(12.5)	(29.2)	(16.7)	(8.3)	(4.2)	(12.5)				(100.0)				
その他の分類不能の腎炎		1	1	6		8	13	12	3	3	2	1	1		51	27	78	9.30	0.84
(%)		(2.0)	(2.0)	(11.8)		(15.7)	(25.5)	(23.5)	(5.9)	(5.9)	(3.9)	(2.0)	(2.0)		(100.0)				
多発性嚢胞腎		2	4	4	4	32	50	45	6	9	6	3	2		167	96	263	9.34	0.75
(%)		(1.2)	(2.4)	(2.4)	(2.4)	(19.2)	(29.9)	(26.9)	(3.6)	(5.4)	(3.6)	(1.8)	(1.2)		(100.0)				
腎硬化症	4	4	11	36	11	131	199	191	22	66	35	14	5	2	731	364	1,095	9.38	0.76
(%)	(0.5)	(0.5)	(1.5)	(4.9)	(1.5)	(17.9)	(27.2)	(26.1)	(3.0)	(9.0)	(4.8)	(1.9)	(0.7)	(0.3)	(100.0)				
悪性高血圧		1		6	3	13	19	18	4	5	3				72	35	107	9.27	0.66
(%)		(1.4)		(8.3)	(4.2)	(18.1)	(26.4)	(25.0)	(5.6)	(6.9)	(4.2)				(100.0)				
糖尿病性腎症	4	12	51	117	25	381	523	421	57	147	75	24	14	5	1,856	943	2,799	9.29	0.78
(%)	(0.2)	(0.6)	(2.7)	(6.3)	(1.3)	(20.5)	(28.2)	(22.7)	(3.1)	(7.9)	(4.0)	(1.3)	(0.8)	(0.3)	(100.0)				
SLE腎炎			1	1	1	5	7	7		4	5				31	15	46	9.51	0.78
(%)			(3.2)	(3.2)	(3.2)	(16.1)	(22.6)	(22.6)		(12.9)	(16.1)				(100.0)				
アミロイド腎						1	2	7		1					11	4	15	9.53	0.39
(%)						(9.1)	(18.2)	(63.6)		(9.1)					(100.0)				
痛風腎		1		1		5	5	10	2	1	1				26	6	32	9.33	0.73
(%)		(3.8)		(3.8)		(19.2)	(19.2)	(38.5)	(7.7)	(3.8)	(3.8)				(100.0)				
先天性代謝異常による腎不全						1	3	5		3					12	5	17	9.65	0.49
(%)						(8.3)	(25.0)	(41.7)		(25.0)					(100.0)				
腎・尿路結核							1								1	1	2	9.30	
(%)							(100.0)								(100.0)				
腎・尿路結石						2	7	5		2	1				17	4	21	9.55	0.51
(%)						(11.8)	(41.2)	(29.4)		(11.8)	(5.9)				(100.0)				
腎・尿路腫瘍			1				4	6	1						12	2	14	9.47	0.51
(%)			(8.3)				(33.3)	(50.0)	(8.3)						(100.0)				
閉塞性尿路障害			1			3	1	6	2	1	1	1			16	5	21	9.64	0.77
(%)			(6.3)			(18.8)	(6.3)	(37.5)	(12.5)	(6.3)	(6.3)	(6.3)			(100.0)				
骨髄腫						1	2			1					4	3	7	9.40	0.57
(%)						(25.0)	(50.0)			(25.0)					(100.0)				
腎形成不全				3		4	9	11	1	6	7	4	2	2	49	23	72	9.94	1.01
(%)			(6.1)			(8.2)	(18.4)	(22.4)	(2.0)	(12.2)	(14.3)	(8.2)	(4.1)	(4.1)	(100.0)				
不明	2	3	13	41	7	89	162	130	21	49	27	11	3	2	560	282	842	9.36	0.79
(%)	(0.4)	(0.5)	(2.3)	(7.3)	(1.3)	(15.9)	(28.9)	(23.2)	(3.8)	(8.8)	(4.8)	(2.0)	(0.5)	(0.4)	(100.0)				
再導入			1	1		5	6	8	5	4	4	1			35	20	55	9.65	0.73
(%)			(2.9)	(2.9)		(14.3)	(17.1)	(22.9)	(14.3)	(11.4)	(11.4)	(2.9)			(100.0)				
その他	1	2	3	3	3	30	37	46	5	23	8	11	2	2	176	87	263	9.56	1.01
(%)	(0.6)	(1.1)	(1.7)	(1.7)	(1.7)	(17.0)	(21.0)	(26.1)	(2.8)	(13.1)	(4.5)	(6.3)	(1.1)	(1.1)	(100.0)				
合計	24	40	123	305	83	1,040	1,612	1,476	206	538	289	96	42	19	5,893	2,876	8,769	9.37	0.80
(%)	(0.4)	(0.7)	(2.1)	(5.2)	(1.4)	(17.6)	(27.4)	(25.0)	(3.5)	(9.1)	(4.9)	(1.6)	(0.7)	(0.3)	(100.0)				
記載なし																			
(%)																			
総計	24	40	123	305	83	1,040	1,612	1,476	206	538	289	96	42	19	5,893	2,876	8,769	9.37	0.80
(%)	(0.4)	(0.7)	(2.1)	(5.2)	(1.4)	(17.6)	(27.4)	(25.0)	(3.5)	(9.1)	(4.9)	(1.6)	(0.7)	(0.3)	(100.0)				

数値下のかっこ内は行方向の合計に対する％です。

(C)Japanese Society for Dialysis Therapy

2015年調査