

表 2927 透析前リン濃度(mg/dL) 治療方法(新分類)別 (透析患者全体)

治療方法 (新分類)	3未満	3.0～	3.5～	4.0～	4.5～	5.0～	5.5～	6.0～	6.1～	6.5～	7.0～	7.5～	8.0～	8.5～	9.0～	合計	記載なし	総計	平均	標準偏差
施設血液透析	9,300	11,122	18,912	27,936	33,168	34,279	29,753	5,006	17,220	15,050	9,401	5,783	3,479	2,127	3,399	225,935	22,790	248,725	5.22	1.45
(%)	(4.1)	(4.9)	(8.4)	(12.4)	(14.7)	(15.2)	(13.2)	(2.2)	(7.6)	(6.7)	(4.2)	(2.6)	(1.5)	(0.9)	(1.5)	(100.0)				
Off-line HDF	217	239	367	574	702	705	618	94	345	366	209	118	94	46	100	4,794	538	5,332	5.28	1.52
(%)	(4.5)	(5.0)	(7.7)	(12.0)	(14.6)	(14.7)	(12.9)	(2.0)	(7.2)	(7.6)	(4.4)	(2.5)	(2.0)	(1.0)	(2.1)	(100.0)				
On-line HDF	1,085	1,478	2,832	4,570	5,855	6,387	5,746	1,060	3,534	3,123	2,010	1,254	741	456	737	40,868	3,659	44,527	5.42	1.43
(%)	(2.7)	(3.6)	(6.9)	(11.2)	(14.3)	(15.6)	(14.1)	(2.6)	(8.6)	(7.6)	(4.9)	(3.1)	(1.8)	(1.1)	(1.8)	(100.0)				
Push/Pull HDF	4	3	13	15	11	14	11	4	8	9	4	3	3	1	4	107	3	110	5.40	1.76
(%)	(3.7)	(2.8)	(12.1)	(14.0)	(10.3)	(13.1)	(10.3)	(3.7)	(7.5)	(8.4)	(3.7)	(2.8)	(2.8)	(0.9)	(3.7)	(100.0)				
ハイオフィльтраーション	8	11	17	29	32	40	37	8	23	20	13	4	4	1	5	252	15	267	5.35	1.40
(%)	(3.2)	(4.4)	(6.7)	(11.5)	(12.7)	(15.9)	(14.7)	(3.2)	(9.1)	(7.9)	(5.2)	(1.6)	(1.6)	(0.4)	(2.0)	(100.0)				
IHDF	131	176	294	472	507	503	425	77	255	227	117	86	69	32	39	3,410	130	3,540	5.19	1.41
(%)	(3.8)	(5.2)	(8.6)	(13.8)	(14.9)	(14.8)	(12.5)	(2.3)	(7.5)	(6.7)	(3.4)	(2.5)	(2.0)	(0.9)	(1.1)	(100.0)				
血液濾過	2	2		2		3	1	1		1	1					13	3	16	4.60	1.71
(%)	(15.4)	(15.4)		(15.4)		(23.1)	(7.7)	(7.7)		(7.7)	(7.7)					(100.0)				
血液吸着透析 (リクセル等使用)	37	50	107	164	233	221	210	27	105	76	38	18	14	2	11	1,313	63	1,376	5.14	1.21
(%)	(2.8)	(3.8)	(8.1)	(12.5)	(17.7)	(16.8)	(16.0)	(2.1)	(8.0)	(5.8)	(2.9)	(1.4)	(1.1)	(0.2)	(0.8)	(100.0)				
在宅血液透析	33	39	49	80	65	52	41	6	27	18	14	7	3	2	6	442	113	555	4.78	1.42
(%)	(7.5)	(8.8)	(11.1)	(18.1)	(14.7)	(11.8)	(9.3)	(1.4)	(6.1)	(4.1)	(3.2)	(1.6)	(0.7)	(0.5)	(1.4)	(100.0)				
腹膜透析 (CAPD)	82	146	310	494	561	566	397	69	204	181	104	67	40	17	24	3,262	1,640	4,902	5.09	1.25
(%)	(2.5)	(4.5)	(9.5)	(15.1)	(17.2)	(17.4)	(12.2)	(2.1)	(6.3)	(5.5)	(3.2)	(2.1)	(1.2)	(0.5)	(0.7)	(100.0)				
腹膜透析 (APD)	49	72	148	252	289	295	252	48	193	143	85	47	31	9	27	1,940	835	2,775	5.32	1.36
(%)	(2.5)	(3.7)	(7.6)	(13.0)	(14.9)	(15.2)	(13.0)	(2.5)	(9.9)	(7.4)	(4.4)	(2.4)	(1.6)	(0.5)	(1.4)	(100.0)				
腹膜透析 (CCPD)	12	19	52	103	114	133	121	29	70	65	30	18	5	3	9	783	309	1,092	5.36	1.30
(%)	(1.5)	(2.4)	(6.6)	(13.2)	(14.6)	(17.0)	(15.5)	(3.7)	(8.9)	(8.3)	(3.8)	(2.3)	(0.6)	(0.4)	(1.1)	(100.0)				
合計	10,960	13,357	23,101	34,691	41,537	43,198	37,612	6,429	21,984	19,279	12,026	7,405	4,483	2,696	4,361	283,119	30,098	313,217	5.25	1.44
(%)	(3.9)	(4.7)	(8.2)	(12.3)	(14.7)	(15.3)	(13.3)	(2.3)	(7.8)	(6.8)	(4.2)	(2.6)	(1.6)	(1.0)	(1.5)	(100.0)				
記載なし (%)																				
総計	10,960	13,357	23,101	34,691	41,537	43,198	37,612	6,429	21,984	19,279	12,026	7,405	4,483	2,696	4,361	283,119	30,098	313,217	5.25	1.44
(%)	(3.9)	(4.7)	(8.2)	(12.3)	(14.7)	(15.3)	(13.3)	(2.3)	(7.8)	(6.8)	(4.2)	(2.6)	(1.6)	(1.0)	(1.5)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。
 *腹膜透析 (CCPD) は、APDと手動のバック交換の両方を行うもの
 (C)Japanese Society for Dialysis Therapy
 2015年調査