

表 2932 透析前リン濃度(mg/dL) 原疾患別 (透析患者全体)

原疾患	3未満	3.0～	3.5～	4.0～	4.5～	5.0～	5.5～	6.0～	6.1～	6.5～	7.0～	7.5～	8.0～	8.5～	9.0～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	2,797	3,559	6,445	10,165	12,474	13,346	11,805	1,980	6,823	5,944	3,649	2,239	1,330	760	1,295	84,611	8,736	93,347	5.30	1.41
(%)	(3.3)	(4.2)	(7.6)	(12.0)	(14.7)	(15.8)	(14.0)	(2.3)	(8.1)	(7.0)	(4.3)	(2.6)	(1.6)	(0.9)	(1.5)	(100.0)				
慢性腎盂腎炎	95	121	223	343	428	406	346	71	212	180	112	61	35	23	43	2,699	236	2,935	5.24	1.42
(%)	(3.5)	(4.5)	(8.3)	(12.7)	(15.9)	(15.0)	(12.8)	(2.6)	(7.9)	(6.7)	(4.1)	(2.3)	(1.3)	(0.9)	(1.6)	(100.0)				
急速進行性糸球体腎炎	124	142	221	242	340	308	284	50	153	128	78	46	37	24	35	2,212	266	2,478	5.11	1.51
(%)	(5.6)	(6.4)	(10.0)	(10.9)	(15.4)	(13.9)	(12.8)	(2.3)	(6.9)	(5.8)	(3.5)	(2.1)	(1.7)	(1.1)	(1.6)	(100.0)				
妊娠腎/妊娠中毒症	35	47	103	155	197	224	221	46	105	115	73	53	25	13	29	1,441	104	1,545	5.45	1.43
(%)	(2.4)	(3.3)	(7.1)	(10.8)	(13.7)	(15.5)	(15.3)	(3.2)	(7.3)	(8.0)	(5.1)	(3.7)	(1.7)	(0.9)	(2.0)	(100.0)				
その他の分類不能の腎炎	55	64	86	138	184	208	170	32	103	93	50	29	25	13	33	1,283	122	1,405	5.33	1.52
(%)	(4.3)	(5.0)	(6.7)	(10.8)	(14.3)	(16.2)	(13.3)	(2.5)	(8.0)	(7.2)	(3.9)	(2.3)	(1.9)	(1.0)	(2.6)	(100.0)				
多発性囊胞腎	251	341	703	1,114	1,543	1,709	1,575	262	892	772	467	273	158	92	130	10,282	974	11,256	5.37	1.35
(%)	(2.4)	(3.3)	(6.8)	(10.8)	(15.0)	(16.6)	(15.3)	(2.5)	(8.7)	(7.5)	(4.5)	(2.7)	(1.5)	(0.9)	(1.3)	(100.0)				
腎硬化症	1,155	1,492	2,465	3,590	4,121	4,023	3,501	541	1,917	1,659	984	623	343	203	357	26,974	2,831	29,805	5.13	1.42
(%)	(4.3)	(5.5)	(9.1)	(13.3)	(15.3)	(14.9)	(13.0)	(2.0)	(7.1)	(6.2)	(3.6)	(2.3)	(1.3)	(0.8)	(1.3)	(100.0)				
悪性高血圧	75	118	183	267	340	346	292	44	190	177	121	75	46	29	60	2,363	255	2,618	5.39	1.53
(%)	(3.2)	(5.0)	(7.7)	(11.3)	(14.4)	(14.6)	(12.4)	(1.9)	(8.0)	(7.5)	(5.1)	(3.2)	(1.9)	(1.2)	(2.5)	(100.0)				
糖尿病性腎症	4,389	5,190	9,062	13,396	15,824	16,398	13,970	2,442	8,457	7,437	4,709	2,908	1,793	1,128	1,652	108,755	11,523	120,278	5.25	1.45
(%)	(4.0)	(4.8)	(8.3)	(12.3)	(14.6)	(15.1)	(12.8)	(2.2)	(7.8)	(6.8)	(4.3)	(2.7)	(1.6)	(1.0)	(1.5)	(100.0)				
SLE腎炎	76	79	150	245	283	321	260	52	134	158	92	63	32	27	36	2,008	219	2,227	5.34	1.49
(%)	(3.8)	(3.9)	(7.5)	(12.2)	(14.1)	(16.0)	(12.9)	(2.6)	(6.7)	(7.9)	(4.6)	(3.1)	(1.6)	(1.3)	(1.8)	(100.0)				
アミロイド腎	20	27	18	50	58	70	57	8	34	28	11	7	7	3	6	404	58	462	5.18	1.44
(%)	(5.0)	(6.7)	(4.5)	(12.4)	(14.4)	(17.3)	(14.1)	(2.0)	(8.4)	(6.9)	(2.7)	(1.7)	(1.7)	(0.7)	(1.5)	(100.0)				
痛風腎	29	45	66	110	142	148	156	24	77	77	36	25	18	9	23	985	90	1,075	5.38	1.48
(%)	(2.9)	(4.6)	(6.7)	(11.2)	(14.4)	(15.0)	(15.8)	(2.4)	(7.8)	(7.8)	(3.7)	(2.5)	(1.8)	(0.9)	(2.3)	(100.0)				
先天性代謝異常による腎不全	9	9	12	22	28	36	34	6	21	20	8	9	8	3	13	238	36	274	5.65	1.68
(%)	(3.8)	(3.8)	(5.0)	(9.2)	(11.8)	(15.1)	(14.3)	(2.5)	(8.8)	(8.4)	(3.4)	(3.8)	(3.4)	(1.3)	(5.5)	(100.0)				
腎・尿路結核	5	8	13	34	26	24	22	4	14	11	6	3	3		3	176	19	195	5.16	1.36
(%)	(2.8)	(4.5)	(7.4)	(19.3)	(14.8)	(13.6)	(12.5)	(2.3)	(8.0)	(6.3)	(3.4)	(1.7)	(1.7)		(1.7)	(100.0)				
腎・尿路結石	17	23	52	73	68	70	74	10	46	33	22	20	7	8	7	530	51	581	5.26	1.49
(%)	(3.2)	(4.3)	(9.8)	(13.8)	(12.8)	(13.2)	(14.0)	(1.9)	(8.7)	(6.2)	(4.2)	(3.8)	(1.3)	(1.5)	(1.3)	(100.0)				
腎・尿路腫瘍	33	48	77	85	107	125	113	22	74	45	31	15	14	12	19	820	90	910	5.26	1.52
(%)	(4.0)	(5.9)	(9.4)	(10.4)	(13.0)	(15.2)	(13.8)	(2.7)	(9.0)	(5.5)	(3.8)	(1.8)	(1.7)	(1.5)	(2.3)	(100.0)				
閉塞性尿路障害	37	28	59	81	96	89	87	16	49	40	31	16	13	7	11	660	70	730	5.22	1.53
(%)	(5.6)	(4.2)	(8.9)	(12.3)	(14.5)	(13.5)	(13.2)	(2.4)	(7.4)	(6.1)	(4.7)	(2.4)	(2.0)	(1.1)	(1.7)	(100.0)				
骨髄腫	19	20	34	42	43	29	26	4	10	10	5	3	4	2	2	253	38	291	4.73	1.41
(%)	(7.5)	(7.9)	(13.4)	(16.6)	(17.0)	(11.5)	(10.3)	(1.6)	(4.0)	(4.0)	(2.0)	(1.2)	(1.6)	(0.8)	(0.8)	(100.0)				
腎形成不全	15	28	35	51	75	101	90	12	48	42	32	29	9	6	24	597	63	660	5.59	1.63
(%)	(2.5)	(4.7)	(5.9)	(8.5)	(12.6)	(16.9)	(15.1)	(2.0)	(8.0)	(7.0)	(5.4)	(4.9)	(1.5)	(1.0)	(4.0)	(100.0)				
不明	1,325	1,484	2,301	3,376	3,927	3,850	3,355	598	1,950	1,886	1,117	644	413	232	401	26,659	3,238	29,897	5.16	1.47
(%)	(5.0)	(5.6)	(8.6)	(12.7)	(14.7)	(14.4)	(12.6)	(2.2)	(7.3)	(6.3)	(4.2)	(2.4)	(1.5)	(0.9)	(1.5)	(100.0)				
再導入	46	73	118	204	257	313	290	40	165	164	88	67	48	31	43	1,947	236	2,183	5.52	1.52
(%)	(2.4)	(3.7)	(6.1)	(10.5)	(13.2)	(16.1)	(14.9)	(2.1)	(8.5)	(8.4)	(4.5)	(3.4)	(2.5)	(1.6)	(2.2)	(100.0)				
その他	353	409	675	908	976	1,048	882	165	509	458	302	197	115	71	139	7,207	837	8,044	5.19	1.52
(%)	(4.9)	(5.7)	(9.4)	(12.6)	(13.5)	(14.5)	(12.2)	(2.3)	(7.1)	(6.4)	(4.2)	(2.7)	(1.6)	(1.0)	(1.9)	(100.0)				
合計	10,960	13,355	23,101	34,691	41,537	43,192	37,610	6,429	21,983	19,277	12,024	7,405	4,483	2,696	4,361	283,104	30,092	313,196	5.25	1.44
(%)	(3.9)	(4.7)	(8.2)	(12.3)	(14.7)	(15.3)	(13.3)	(2.3)	(7.8)	(6.8)	(4.2)	(2.6)	(1.6)	(1.0)	(1.5)	(100.0)				
記載なし		2				6	2		1	2	2					15	6	21	5.53	1.25
(%)		(13.3)				(40.0)	(13.3)		(6.7)	(13.3)	(13.3)					(100.0)				
総計	10,960	13,357	23,101	34,691	41,537	43,198	37,612	6,429	21,984	19,279	12,026	7,405	4,483	2,696	4,361	283,119	30,098	313,217	5.25	1.44
(%)	(3.9)	(4.7)	(8.2)	(12.3)	(14.7)	(15.3)	(13.3)	(2.3)	(7.8)	(6.8)	(4.2)	(2.6)	(1.6)	(1.0)	(1.5)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2015年調査