

表 2946 透析前リン濃度(mg/dL) 原疾患別（施設血液透析患者 透析回数週3回）

原疾患	3未満	3.0～	3.5～	4.0～	4.5～	5.0～	5.5～	6.0～	6.1～	6.5～	7.0～	7.5～	8.0～	8.5～	9.0～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	2,197	2,744	4,786	7,496	9,078	9,735	8,582	1,417	4,923	4,275	2,620	1,615	977	541	939	61,925	1,549	63,474	5.28	1.42
(%)	(3.5)	(4.4)	(7.7)	(12.1)	(14.7)	(15.7)	(13.9)	(2.3)	(7.9)	(6.9)	(4.2)	(2.6)	(1.6)	(0.9)	(1.5)	(100.0)				
慢性腎盂腎炎	71	95	159	259	314	296	260	56	155	136	86	44	24	17	28	2,000	40	2,040	5.22	1.39
(%)	(3.6)	(4.8)	(8.0)	(13.0)	(15.7)	(14.8)	(13.0)	(2.8)	(7.8)	(6.8)	(4.3)	(2.2)	(1.2)	(0.9)	(1.4)	(100.0)				
急速進行性糸球体腎炎	101	106	172	197	258	249	209	37	119	99	60	37	28	18	27	1,717	54	1,771	5.09	1.51
(%)	(5.9)	(6.2)	(10.0)	(11.5)	(15.0)	(14.5)	(12.2)	(2.2)	(6.9)	(5.8)	(3.5)	(2.2)	(1.6)	(1.0)	(1.6)	(100.0)				
妊娠腎/妊娠中毒症	27	29	65	93	137	145	145	35	72	77	53	37	17	11	19	962	19	981	5.49	1.44
(%)	(2.8)	(3.0)	(6.8)	(9.7)	(14.2)	(15.1)	(15.1)	(3.6)	(7.5)	(8.0)	(5.5)	(3.8)	(1.8)	(1.1)	(2.0)	(100.0)				
その他の分類不能の腎炎	36	51	58	90	134	148	122	25	74	62	36	20	17	9	23	905	24	929	5.34	1.51
(%)	(4.0)	(5.6)	(6.4)	(9.9)	(14.8)	(16.4)	(13.5)	(2.8)	(8.2)	(6.9)	(4.0)	(2.2)	(1.9)	(1.0)	(2.5)	(100.0)				
多発性嚢胞腎	196	265	559	872	1,204	1,302	1,188	199	666	571	364	195	104	65	92	7,842	174	8,016	5.34	1.33
(%)	(2.5)	(3.4)	(7.1)	(11.1)	(15.4)	(16.6)	(15.1)	(2.5)	(8.5)	(7.3)	(4.6)	(2.5)	(1.3)	(0.8)	(1.2)	(100.0)				
腎硬化症	927	1,174	1,975	2,842	3,263	3,190	2,826	436	1,505	1,312	779	494	261	168	278	21,430	577	22,007	5.13	1.41
(%)	(4.3)	(5.5)	(9.2)	(13.3)	(15.2)	(14.9)	(13.2)	(2.0)	(7.0)	(6.1)	(3.6)	(2.3)	(1.2)	(0.8)	(1.3)	(100.0)				
悪性高血圧	68	93	138	202	270	275	212	34	146	130	92	55	35	26	42	1,818	43	1,861	5.36	1.54
(%)	(3.7)	(5.1)	(7.6)	(11.1)	(14.9)	(15.1)	(11.7)	(1.9)	(8.0)	(7.2)	(5.1)	(3.0)	(1.9)	(1.4)	(2.3)	(100.0)				
糖尿病性腎症	3,613	4,251	7,344	10,765	12,715	13,109	11,179	1,901	6,621	5,798	3,732	2,281	1,384	886	1,290	86,869	2,557	89,426	5.23	1.45
(%)	(4.2)	(4.9)	(8.5)	(12.4)	(14.6)	(15.1)	(12.9)	(2.2)	(7.6)	(6.7)	(4.3)	(2.6)	(1.6)	(1.0)	(1.5)	(100.0)				
SLE腎炎	52	64	110	180	206	234	186	34	94	124	59	42	22	20	24	1,451	53	1,504	5.31	1.46
(%)	(3.6)	(4.4)	(7.6)	(12.4)	(14.2)	(16.1)	(12.8)	(2.3)	(6.5)	(8.5)	(4.1)	(2.9)	(1.5)	(1.4)	(1.7)	(100.0)				
アミロイド腎	11	20	15	36	43	54	43	6	28	20	7	6	7	1	4	301	13	314	5.20	1.36
(%)	(3.7)	(6.6)	(5.0)	(12.0)	(14.3)	(17.9)	(14.3)	(2.0)	(9.3)	(6.6)	(2.3)	(2.0)	(2.3)	(0.3)	(1.3)	(100.0)				
痛風腎	21	35	59	83	118	108	128	15	59	57	26	17	14	6	17	763	15	778	5.34	1.43
(%)	(2.8)	(4.6)	(7.7)	(10.9)	(15.5)	(14.2)	(16.8)	(2.0)	(7.7)	(7.5)	(3.4)	(2.2)	(1.8)	(0.8)	(2.2)	(100.0)				
先天性代謝異常による腎不全	7	7	7	16	19	25	26	5	18	8	8	6	4	3	9	168	5	173	5.61	1.67
(%)	(4.2)	(4.2)	(4.2)	(9.5)	(11.3)	(14.9)	(15.5)	(3.0)	(10.7)	(4.8)	(4.8)	(3.6)	(2.4)	(1.8)	(5.4)	(100.0)				
腎・尿路結核	4	4	10	28	17	19	18	4	11	9	5	3	2		3	137	4	141	5.22	1.40
(%)	(2.9)	(2.9)	(7.3)	(20.4)	(12.4)	(13.9)	(13.1)	(2.9)	(8.0)	(6.6)	(3.6)	(2.2)	(1.5)		(2.2)	(100.0)				
腎・尿路結石	15	17	46	56	54	49	61	7	35	25	17	15	5	6	5	413	8	421	5.22	1.52
(%)	(3.6)	(4.1)	(11.1)	(13.6)	(13.1)	(11.9)	(14.8)	(1.7)	(8.5)	(6.1)	(4.1)	(3.6)	(1.2)	(1.5)	(1.2)	(100.0)				
腎・尿路腫瘍	27	39	62	69	89	98	87	14	59	36	29	12	13	10	17	661	17	678	5.27	1.56
(%)	(4.1)	(5.9)	(9.4)	(10.4)	(13.5)	(14.8)	(13.2)	(2.1)	(8.9)	(5.4)	(4.4)	(1.8)	(2.0)	(1.5)	(2.6)	(100.0)				
閉塞性尿路障害	28	21	41	57	76	69	60	14	36	28	21	8	8	6	8	481	13	494	5.18	1.49
(%)	(5.8)	(4.4)	(8.5)	(11.9)	(15.8)	(14.3)	(12.5)	(2.9)	(7.5)	(5.8)	(4.4)	(1.7)	(1.7)	(1.2)	(1.7)	(100.0)				
骨髄腫	13	14	21	27	30	21	22	3	8	9	5	2	3	2	2	182	8	190	4.86	1.48
(%)	(7.1)	(7.7)	(11.5)	(14.8)	(16.5)	(11.5)	(12.1)	(1.6)	(4.4)	(4.9)	(2.7)	(1.1)	(1.6)	(1.1)	(1.1)	(100.0)				
腎形成不全	12	17	30	29	43	65	61	8	35	26	20	22	5	5	15	393	7	400	5.57	1.63
(%)	(3.1)	(4.3)	(7.6)	(7.4)	(10.9)	(16.5)	(15.5)	(2.0)	(8.9)	(6.6)	(5.1)	(5.6)	(1.3)	(1.3)	(3.8)	(100.0)				
不明	1,064	1,136	1,775	2,561	3,031	2,941	2,546	456	1,460	1,275	825	470	303	170	302	20,315	861	21,176	5.14	1.47
(%)	(5.2)	(5.6)	(8.7)	(12.6)	(14.9)	(14.5)	(12.5)	(2.2)	(7.2)	(6.3)	(4.1)	(2.3)	(1.5)	(0.8)	(1.5)	(100.0)				
再導入	35	58	79	146	172	213	189	25	104	119	61	44	31	21	30	1,327	29	1,356	5.50	1.56
(%)	(2.6)	(4.4)	(6.0)	(11.0)	(13.0)	(16.1)	(14.2)	(1.9)	(7.8)	(9.0)	(4.6)	(3.3)	(2.3)	(1.6)	(2.3)	(100.0)				
その他	283	321	527	696	751	794	679	123	386	339	221	147	88	57	106	5,518	197	5,715	5.17	1.53
(%)	(5.1)	(5.8)	(9.6)	(12.6)	(13.6)	(14.4)	(12.3)	(2.2)	(7.0)	(6.1)	(4.0)	(2.7)	(1.6)	(1.0)	(1.9)	(100.0)				
合計	8,808	10,561	18,038	26,800	32,022	33,139	28,829	4,854	16,614	14,535	9,126	5,572	3,352	2,048	3,280	217,578	6,267	223,845	5.23	1.44
(%)	(4.0)	(4.9)	(8.3)	(12.3)	(14.7)	(15.2)	(13.2)	(2.2)	(7.6)	(6.7)	(4.2)	(2.6)	(1.5)	(0.9)	(1.5)	(100.0)				
記載なし		1				2	1		1	1						6	1	7	5.43	1.28
(%)		(16.7)				(33.3)	(16.7)		(16.7)	(16.7)						(100.0)				
総計	8,808	10,562	18,038	26,800	32,022	33,141	28,830	4,854	16,615	14,536	9,126	5,572	3,352	2,048	3,280	217,584	6,268	223,852	5.23	1.44
(%)	(4.0)	(4.9)	(8.3)	(12.3)	(14.7)	(15.2)	(13.3)	(2.2)	(7.6)	(6.7)	(4.2)	(2.6)	(1.5)	(0.9)	(1.5)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2015年調査