

表 2960 透析前リン濃度(mg/dL) 原疾患別（血液透析濾過患者 透析回数週3回）

原疾患	3未満	3.0～	3.5～	4.0～	4.5～	5.0～	5.5～	6.0～	6.1～	6.5～	7.0～	7.5～	8.0～	8.5～	9.0～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	418	562	1,157	1,889	2,532	2,717	2,457	438	1,480	1,314	825	487	286	186	292	17,040	422	17,462	5.42	1.39
(%)	(2.5)	(3.3)	(6.8)	(11.1)	(14.9)	(15.9)	(14.4)	(2.6)	(8.7)	(7.7)	(4.8)	(2.9)	(1.7)	(1.1)	(1.7)	(100.0)				
慢性腎盂腎炎	11	15	37	50	77	81	59	13	39	37	20	13	8	6	12	478	6	484	5.45	1.49
(%)	(2.3)	(3.1)	(7.7)	(10.5)	(16.1)	(16.9)	(12.3)	(2.7)	(8.2)	(7.7)	(4.2)	(2.7)	(1.7)	(1.3)	(2.5)	(100.0)				
急速進行性糸球体腎炎	12	19	28	24	66	42	55	11	26	26	15	9	6	4	7	350	3	353	5.35	1.48
(%)	(3.4)	(5.4)	(8.0)	(6.9)	(18.9)	(12.0)	(15.7)	(3.1)	(7.4)	(7.4)	(4.3)	(2.6)	(1.7)	(1.1)	(2.0)	(100.0)				
妊娠腎/妊娠中毒症	7	13	30	46	50	65	60	9	27	34	17	14	4	1	8	385	16	401	5.40	1.38
(%)	(1.8)	(3.4)	(7.8)	(11.9)	(13.0)	(16.9)	(15.6)	(2.3)	(7.0)	(8.8)	(4.4)	(3.6)	(1.0)	(0.3)	(2.1)	(100.0)				
その他の分類不能の腎炎	12	9	14	26	28	44	37	4	22	27	10	8	5	2	9	257	4	261	5.50	1.57
(%)	(4.7)	(3.5)	(5.4)	(10.1)	(10.9)	(17.1)	(14.4)	(1.6)	(8.6)	(10.5)	(3.9)	(3.1)	(1.9)	(0.8)	(3.5)	(100.0)				
多発性囊胞腎	33	53	105	171	262	302	296	49	175	155	80	59	48	19	30	1,837	43	1,880	5.53	1.37
(%)	(1.8)	(2.9)	(5.7)	(9.3)	(14.3)	(16.4)	(16.1)	(2.7)	(9.5)	(8.4)	(4.4)	(3.2)	(2.6)	(1.0)	(1.6)	(100.0)				
腎硬化症	137	184	283	490	544	572	498	80	276	245	159	102	63	29	59	3,721	99	3,820	5.25	1.43
(%)	(3.7)	(4.9)	(7.6)	(13.2)	(14.6)	(15.4)	(13.4)	(2.1)	(7.4)	(6.6)	(4.3)	(2.7)	(1.7)	(0.8)	(1.6)	(100.0)				
悪性高血圧	4	16	27	39	52	51	53	9	31	37	20	14	10	3	15	381	10	391	5.61	1.51
(%)	(1.0)	(4.2)	(7.1)	(10.2)	(13.6)	(13.4)	(13.9)	(2.4)	(8.1)	(9.7)	(5.2)	(3.7)	(2.6)	(0.8)	(3.9)	(100.0)				
糖尿病性腎症	533	655	1,223	1,928	2,324	2,477	2,156	424	1,419	1,284	785	502	331	191	285	16,517	443	16,960	5.38	1.46
(%)	(3.2)	(4.0)	(7.4)	(11.7)	(14.1)	(15.0)	(13.1)	(2.6)	(8.6)	(7.8)	(4.8)	(3.0)	(2.0)	(1.2)	(1.7)	(100.0)				
SLE腎炎	15	8	24	45	64	64	57	12	31	27	24	18	5	7	12	413	13	426	5.53	1.58
(%)	(3.6)	(1.9)	(5.8)	(10.9)	(15.5)	(15.5)	(13.8)	(2.9)	(7.5)	(6.5)	(5.8)	(4.4)	(1.2)	(1.7)	(2.9)	(100.0)				
アミロイド腎	5	7	3	6	11	10	11	2	3	8	3	1		1	1	72	2	74	5.19	1.65
(%)	(6.9)	(9.7)	(4.2)	(8.3)	(15.3)	(13.9)	(15.3)	(2.8)	(4.2)	(11.1)	(4.2)	(1.4)		(1.4)	(1.4)	(100.0)				
痛風腎	5	6	5	16	18	29	24	7	15	14	8	6	3	3	2	161	5	166	5.57	1.54
(%)	(3.1)	(3.7)	(3.1)	(9.9)	(11.2)	(18.0)	(14.9)	(4.3)	(9.3)	(8.7)	(5.0)	(3.7)	(1.9)	(1.9)	(1.2)	(100.0)				
先天性代謝異常 による腎不全	1	1	3	4	5	6	6	1	2	10		3	3		3	48	1	49	5.97	1.74
(%)	(2.1)	(2.1)	(6.3)	(8.3)	(10.4)	(12.5)	(12.5)	(2.1)	(4.2)	(20.8)		(6.3)	(6.3)		(6.3)	(100.0)				
腎・尿路結核	1	3		4	9	5	3		3			1		1		30	1	31	5.00	1.15
(%)	(3.3)	(10.0)		(13.3)	(30.0)	(16.7)	(10.0)		(10.0)		(3.3)		(3.3)			(100.0)				
腎・尿路結石	1	3	2	9	7	16	6	3	10	7	4	5	2	1	1	77	2	79	5.67	1.38
(%)	(1.3)	(3.9)	(2.6)	(11.7)	(9.1)	(20.8)	(7.8)	(3.9)	(13.0)	(9.1)	(5.2)	(6.5)	(2.6)	(1.3)	(1.3)	(100.0)				
腎・尿路腫瘍	3	7	10	11	12	20	21	8	11	6	2	2	1	2	1	117	2	119	5.27	1.29
(%)	(2.6)	(6.0)	(8.5)	(9.4)	(10.3)	(17.1)	(17.9)	(6.8)	(9.4)	(5.1)	(1.7)	(1.7)	(0.9)	(1.7)	(0.9)	(100.0)				
閉塞性尿路障害		2	12	11	12	14	17	1	9	9	5	7	1	1	3	104		104	5.63	1.57
(%)		(1.9)	(11.5)	(10.6)	(11.5)	(13.5)	(16.3)	(1.0)	(8.7)	(8.7)	(4.8)	(6.7)	(1.0)	(1.0)	(2.9)	(100.0)				
骨髄腫	3	2	3	4	7	3	3	1	1	1		1	1			30	3	33	4.74	1.38
(%)	(10.0)	(6.7)	(10.0)	(13.3)	(23.3)	(10.0)	(10.0)	(3.3)	(3.3)	(3.3)		(3.3)	(3.3)			(100.0)				
腎形成不全	1	9	4	17	19	20	18	2	8	9	10	7	3	1	6	134	2	136	5.67	1.73
(%)	(0.7)	(6.7)	(3.0)	(12.7)	(14.2)	(14.9)	(13.4)	(1.5)	(6.0)	(6.7)	(7.5)	(5.2)	(2.2)	(0.7)	(4.5)	(100.0)				
不明	158	214	349	558	638	686	601	104	354	304	230	130	83	48	78	4,535	168	4,703	5.31	1.46
(%)	(3.5)	(4.7)	(7.7)	(12.3)	(14.1)	(15.1)	(13.3)	(2.3)	(7.8)	(6.7)	(5.1)	(2.9)	(1.8)	(1.1)	(1.7)	(100.0)				
再導入	7	10	24	40	60	84	79	13	49	31	23	20	14	9	10	473	10	483	5.66	1.38
(%)	(1.5)	(2.1)	(5.1)	(8.5)	(12.7)	(17.8)	(16.7)	(2.7)	(10.4)	(6.6)	(4.9)	(4.2)	(3.0)	(1.9)	(2.1)	(100.0)				
その他	35	46	78	142	142	165	162	33	88	93	60	36	19	11	24	1,134	26	1,160	5.41	1.47
(%)	(3.1)	(4.1)	(6.9)	(12.5)	(12.5)	(14.6)	(14.3)	(2.9)	(7.8)	(8.2)	(5.3)	(3.2)	(1.7)	(1.0)	(2.1)	(100.0)				
合計	1,402	1,844	3,421	5,530	6,939	7,476	6,679	1,224	4,079	3,678	2,301	1,444	897	525	858	48,294	1,281	49,575	5.39	1.43
(%)	(2.9)	(3.8)	(7.1)	(11.5)	(14.4)	(15.5)	(13.8)	(2.5)	(8.4)	(7.6)	(4.8)	(3.0)	(1.9)	(1.1)	(1.8)	(100.0)				
記載なし						3	1				2					6		6	5.92	1.03
(%)						(50.0)	(16.7)				(33.3)					(100.0)				
総計	1,402	1,844	3,421	5,530	6,939	7,476	6,680	1,224	4,079	3,678	2,303	1,444	897	525	858	48,300	1,281	49,581	5.39	1.43
(%)	(2.9)	(3.8)	(7.1)	(11.4)	(14.4)	(15.5)	(13.8)	(2.5)	(8.4)	(7.6)	(4.8)	(3.0)	(1.9)	(1.1)	(1.8)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2015年調査