

表 2974 リン濃度(mg/dL) 原疾患別（腹膜透析患者）

原疾患	3未満	3.0～	3.5～	4.0～	4.5～	5.0～	5.5～	6.0～	6.1～	6.5～	7.0～	7.5～	8.0～	8.5～	9.0～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	36	58	146	263	321	327	285	60	142	139	94	53	19	7	22	1,972	858	2,830	5.30	1.28
(%)	(1.8)	(2.9)	(7.4)	(13.3)	(16.3)	(16.6)	(14.5)	(3.0)	(7.2)	(7.0)	(4.8)	(2.7)	(1.0)	(0.4)	(1.1)	(100.0)				
慢性腎盂腎炎	3	4	8	11	15	11	9		10	1	1	2			1	76	21	97	4.92	1.27
(%)	(3.9)	(5.3)	(10.5)	(14.5)	(19.7)	(14.5)	(11.8)		(13.2)	(1.3)	(1.3)	(2.6)			(1.3)	(100.0)				
急速進行性糸球体腎炎	4	1	7	6	4	6	6	1	4	3	2					44	22	66	4.82	1.39
(%)	(9.1)	(2.3)	(15.9)	(13.6)	(9.1)	(13.6)	(13.6)	(2.3)	(9.1)	(6.8)	(4.5)					(100.0)				
妊娠腎/妊娠中毒症			2	5	3	5	2	1	1	1	1		2		2	25	2	27	5.74	1.77
(%)			(8.0)	(20.0)	(12.0)	(20.0)	(8.0)	(4.0)	(4.0)	(4.0)	(4.0)		(8.0)		(8.0)	(100.0)				
その他の分類不能の腎炎	1	1	3	12	12	5	7	3	4	1		1	2	1	1	54	24	78	5.22	1.47
(%)	(1.9)	(1.9)	(5.6)	(22.2)	(22.2)	(9.3)	(13.0)	(5.6)	(7.4)	(1.9)		(1.9)	(3.7)	(1.9)	(1.9)	(100.0)				
多発性囊胞腎	2	5	11	16	21	38	22	8	15	12	7	7	3	1		168	95	263	5.42	1.17
(%)	(1.2)	(3.0)	(6.5)	(9.5)	(12.5)	(22.6)	(13.1)	(4.8)	(8.9)	(7.1)	(4.2)	(4.2)	(1.8)	(0.6)		(100.0)				
腎硬化症	18	42	93	100	149	110	75	14	62	40	15	10	8	1	5	742	353	1,095	4.97	1.23
(%)	(2.4)	(5.7)	(12.5)	(13.5)	(20.1)	(14.8)	(10.1)	(1.9)	(8.4)	(5.4)	(2.0)	(1.3)	(1.1)	(0.1)	(0.7)	(100.0)				
悪性高血圧	1	2	8	10	9	10	11	1	8	4	3	1	1		2	71	36	107	5.34	1.44
(%)	(1.4)	(2.8)	(11.3)	(14.1)	(12.7)	(14.1)	(15.5)	(1.4)	(11.3)	(5.6)	(4.2)	(1.4)	(1.4)		(2.8)	(100.0)				
糖尿病性腎症	55	77	145	263	284	327	246	41	149	125	62	39	25	13	19	1,870	929	2,799	5.20	1.31
(%)	(2.9)	(4.1)	(7.8)	(14.1)	(15.2)	(17.5)	(13.2)	(2.2)	(8.0)	(6.7)	(3.3)	(2.1)	(1.3)	(0.7)	(1.0)	(100.0)				
SLE腎炎	3	1	3	5	4	4	3		3		3		3			32	14	46	5.17	1.56
(%)	(9.4)	(3.1)	(9.4)	(15.6)	(12.5)	(12.5)	(9.4)		(9.4)		(9.4)		(9.4)			(100.0)				
アミロイド腎				5	1	2	2		1							11	4	15	4.84	0.78
(%)				(45.5)	(9.1)	(18.2)	(18.2)		(9.1)							(100.0)				
痛風腎	1	1	1	5	2	4	1	1	1	4	1	1	1		2	26	6	32	5.78	2.02
(%)	(3.8)	(3.8)	(3.8)	(19.2)	(7.7)	(15.4)	(3.8)	(3.8)	(3.8)	(15.4)	(3.8)	(3.8)	(3.8)		(7.7)	(100.0)				
先天性代謝異常 による腎不全	1		2	1	2	3	2		1	1						13	4	17	4.89	1.23
(%)	(7.7)		(15.4)	(7.7)	(15.4)	(23.1)	(15.4)		(7.7)	(7.7)						(100.0)				
腎・尿路結核				1												1	1	2	4.40	
(%)				(100.0)												(100.0)				
腎・尿路結石		2	3	2	3	3	4									17	4	21	4.62	0.91
(%)		(11.8)	(17.6)	(11.8)	(17.6)	(17.6)	(23.5)									(100.0)				
腎・尿路腫瘍		1	3		2	4			1	1						12	2	14	4.81	1.15
(%)		(8.3)	(25.0)		(16.7)	(33.3)			(8.3)	(8.3)						(100.0)				
閉塞性尿路障害	2	1	1	2	1	2	4		1		2					16	5	21	4.91	1.42
(%)	(12.5)	(6.3)	(6.3)	(12.5)	(6.3)	(12.5)	(25.0)		(6.3)		(12.5)					(100.0)				
骨髄腫		1		2		1										4	3	7	4.10	0.90
(%)		(25.0)		(50.0)		(25.0)										(100.0)				
腎形成不全	2	1	1	4	6	13	9	1	4	4	2				2	49	23	72	5.46	1.30
(%)	(4.1)	(2.0)	(2.0)	(8.2)	(12.2)	(26.5)	(18.4)	(2.0)	(8.2)	(8.2)	(4.1)				(4.1)	(100.0)				
不明	10	27	45	110	93	76	61	11	49	38	20	11	8	4	4	567	275	842	5.14	1.27
(%)	(1.8)	(4.8)	(7.9)	(19.4)	(16.4)	(13.4)	(10.8)	(1.9)	(8.6)	(6.7)	(3.5)	(1.9)	(1.4)	(0.7)	(0.7)	(100.0)				
再導入			5	4	4	8	5	1	3	3		1	1	1		36	19	55	5.37	1.21
(%)			(13.9)	(11.1)	(11.1)	(22.2)	(13.9)	(2.8)	(8.3)	(8.3)		(2.8)	(2.8)	(2.8)		(100.0)				
その他	4	12	23	22	28	35	16	3	8	12	6	6	3	1		179	84	263	5.03	1.31
(%)	(2.2)	(6.7)	(12.8)	(12.3)	(15.6)	(19.6)	(8.9)	(1.7)	(4.5)	(6.7)	(3.4)	(3.4)	(1.7)	(0.6)		(100.0)				
合計	143	237	510	849	964	994	770	146	467	389	219	132	76	29	60	5,985	2,784	8,769	5.20	1.30
(%)	(2.4)	(4.0)	(8.5)	(14.2)	(16.1)	(16.6)	(12.9)	(2.4)	(7.8)	(6.5)	(3.7)	(2.2)	(1.3)	(0.5)	(1.0)	(100.0)				
記載なし																				
(%)																				
総計	143	237	510	849	964	994	770	146	467	389	219	132	76	29	60	5,985	2,784	8,769	5.20	1.30
(%)	(2.4)	(4.0)	(8.5)	(14.2)	(16.1)	(16.6)	(12.9)	(2.4)	(7.8)	(6.5)	(3.7)	(2.2)	(1.3)	(0.5)	(1.0)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2015年調査