

表 2986 PTH値(pg/mL) 治療方法(新分類)別(透析患者全体) PTH測定法:intact-PTH

治療方法 (新分類)	20未満	20～	40～	60～	80～	100～	120～	140～	160～	180～	240～	241～	300～	360～	420～	480～	500～	540～	600～	800～	合計	記載なし	総計	平均	標準偏差
施設血液透析	8,425	13,015	14,096	15,027	15,510	15,167	14,651	13,649	12,427	28,432	362	16,683	9,760	5,623	3,346	824	1,275	1,416	2,391	1,862	193,941	4,839	198,780	177.14	168.85
(%)	(4.3)	(6.7)	(7.3)	(7.7)	(8.0)	(7.8)	(7.6)	(7.0)	(6.4)	(14.7)	(0.2)	(8.6)	(5.0)	(2.9)	(1.7)	(0.4)	(0.7)	(0.7)	(1.2)	(1.0)	(100.0)				
Off-line HDF	215	292	303	341	352	343	334	267	266	563	6	355	197	126	68	21	25	26	54	54	4,208	58	4,266	176.77	183.48
(%)	(5.1)	(6.9)	(7.2)	(8.1)	(8.4)	(8.2)	(7.9)	(6.3)	(6.3)	(13.4)	(0.1)	(8.4)	(4.7)	(3.0)	(1.6)	(0.5)	(0.6)	(0.6)	(1.3)	(1.3)	(100.0)				
On-line HDF	1,471	2,047	2,334	2,680	2,875	2,819	2,775	2,552	2,373	5,500	61	3,320	1,923	1,088	681	167	260	271	488	434	36,119	642	36,761	185.82	178.76
(%)	(4.1)	(5.7)	(6.5)	(7.4)	(8.0)	(7.8)	(7.7)	(7.1)	(6.6)	(15.2)	(0.2)	(9.2)	(5.3)	(3.0)	(1.9)	(0.5)	(0.7)	(0.8)	(1.4)	(1.2)	(100.0)				
Push/Pull HDF		10	4	7	9	3	5	7	5	19	2	9	7	5	3	1			1		97	3	100	184.90	120.51
(%)		(10.3)	(4.1)	(7.2)	(9.3)	(3.1)	(5.2)	(7.2)	(5.2)	(19.6)	(2.1)	(9.3)	(7.2)	(5.2)	(3.1)	(1.0)			(1.0)		(100.0)				
ハイオフィルトレーション	7	12	11	13	15	13	14	13	13	42		13	14	9	4			1	7	2	203		203	197.50	168.30
(%)	(3.4)	(5.9)	(5.4)	(6.4)	(7.4)	(6.4)	(6.9)	(6.4)	(6.4)	(20.7)		(6.4)	(6.9)	(4.4)	(2.0)			(0.5)	(3.4)	(1.0)	(100.0)				
IHDF	131	184	227	253	265	253	259	228	204	450	7	262	150	75	41	14	23	25	39	21	3,111	58	3,169	170.96	146.67
(%)	(4.2)	(5.9)	(7.3)	(8.1)	(8.5)	(8.1)	(8.3)	(7.3)	(6.6)	(14.5)	(0.2)	(8.4)	(4.8)	(2.4)	(1.3)	(0.5)	(0.7)	(0.8)	(1.3)	(0.7)	(100.0)				
血液濾過	2	1			2		1		1	2		1									10		10	127.43	95.69
(%)	(20.0)	(10.0)			(20.0)		(10.0)		(10.0)	(20.0)		(10.0)									(100.0)				
血液吸着透析 (リクセル等使用)	109	78	99	92	95	97	79	64	67	147	2	75	41	26	17	6	5	8	7	7	1,121	18	1,139	152.90	156.40
(%)	(9.7)	(7.0)	(8.8)	(8.2)	(8.5)	(8.7)	(7.0)	(5.7)	(6.0)	(13.1)	(0.2)	(6.7)	(3.7)	(2.3)	(1.5)	(0.5)	(0.4)	(0.7)	(0.6)	(0.6)	(100.0)				
在宅血液透析	14	16	17	31	34	23	26	33	25	46		37	22	11	9	5	3	4	12	9	377	5	382	213.53	205.55
(%)	(3.7)	(4.2)	(4.5)	(8.2)	(9.0)	(6.1)	(6.9)	(8.8)	(6.6)	(12.2)		(9.8)	(5.8)	(2.9)	(2.4)	(1.3)	(0.8)	(1.1)	(3.2)	(2.4)	(100.0)				
腹膜透析 (CAPD)	76	118	156	150	152	161	148	152	123	334	6	277	167	88	78	25	28	39	56	61	2,395	180	2,575	224.97	210.41
(%)	(3.2)	(4.9)	(6.5)	(6.3)	(6.3)	(6.7)	(6.2)	(6.3)	(5.1)	(13.9)	(0.3)	(11.6)	(7.0)	(3.7)	(3.3)	(1.0)	(1.2)	(1.6)	(2.3)	(2.5)	(100.0)				
腹膜透析 (APD)	27	77	79	85	82	99	81	83	74	209	2	169	98	61	37	14	20	20	47	44	1,408	125	1,533	242.77	247.28
(%)	(1.9)	(5.5)	(5.6)	(6.0)	(5.8)	(7.0)	(5.8)	(5.9)	(5.3)	(14.8)	(0.1)	(12.0)	(7.0)	(4.3)	(2.6)	(1.0)	(1.4)	(1.4)	(3.3)	(3.1)	(100.0)				
腹膜透析 (CCPD)	12	35	32	51	37	44	40	33	38	97	1	57	43	28	20	2	4	9	16	13	612	23	635	226.21	252.55
(%)	(2.0)	(5.7)	(5.2)	(8.3)	(6.0)	(7.2)	(6.5)	(5.4)	(6.2)	(15.8)	(0.2)	(9.3)	(7.0)	(4.6)	(3.3)	(0.3)	(0.7)	(1.5)	(2.6)	(2.1)	(100.0)				
合計 (%)	10,489 (4.3)	15,885 (6.5)	17,358 (7.1)	18,730 (7.7)	19,428 (8.0)	19,022 (7.8)	18,413 (7.6)	17,081 (7.0)	15,616 (6.4)	35,841 (14.7)	449 (0.2)	21,258 (8.7)	12,422 (5.1)	7,140 (2.9)	4,304 (1.8)	1,079 (0.4)	1,643 (0.7)	1,819 (0.7)	3,118 (1.3)	2,507 (1.0)	243,602 (100.0)	5,951	249,553	179.28	171.80
記載なし (%)																									
総計 (%)	10,489 (4.3)	15,885 (6.5)	17,358 (7.1)	18,730 (7.7)	19,428 (8.0)	19,022 (7.8)	18,413 (7.6)	17,081 (7.0)	15,616 (6.4)	35,841 (14.7)	449 (0.2)	21,258 (8.7)	12,422 (5.1)	7,140 (2.9)	4,304 (1.8)	1,079 (0.4)	1,643 (0.7)	1,819 (0.7)	3,118 (1.3)	2,507 (1.0)	243,602 (100.0)	5,951	249,553	179.28	171.80

数値下のかっこ内は行方向の合計に対する%です。
*腹膜透析 (CCPD) は、APDと手動のバック交換の両方を行うもの
(C)Japanese Society for Dialysis Therapy
2015年調査