

表 3004 PTH値(pg/mL) 原疾患別（施設血液透析患者 透析回数週3回）PTH測定法intact-PTH

原疾患	20未満	20～	40～	60～	80～	100～	120～	140～	160～	180～	240～	241～	300～	360～	420～	480～	500～	540～	600～	800～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	2,588	3,351	3,658	4,033	4,063	4,099	3,969	3,713	3,352	7,755	96	4,638	2,867	1,737	1,028	284	418	441	820	752	53,662	981	54,643	187.52	188.80
(%)	(4.8)	(6.2)	(6.8)	(7.5)	(7.6)	(7.6)	(7.4)	(6.9)	(6.2)	(14.5)	(0.2)	(8.6)	(5.3)	(3.2)	(1.9)	(0.5)	(0.8)	(0.8)	(1.5)	(1.4)	(100.0)				
慢性腎盂腎炎	91	87	108	134	134	134	123	107	103	259	1	154	80	55	25	10	13	16	23	21	1,678	37	1,715	184.17	173.10
(%)	(5.4)	(5.2)	(6.4)	(8.0)	(8.0)	(8.0)	(7.3)	(6.4)	(6.1)	(15.4)	(0.1)	(9.2)	(4.8)	(3.3)	(1.5)	(0.6)	(0.8)	(1.0)	(1.4)	(1.3)	(100.0)				
急速進行性糸球体腎炎	55	92	121	117	129	112	110	89	101	218	2	109	62	28	20	5	9	7	12	9	1,407	38	1,445	162.33	137.13
(%)	(3.9)	(6.5)	(8.6)	(8.3)	(9.2)	(8.0)	(7.8)	(6.3)	(7.2)	(15.5)	(0.1)	(7.7)	(4.4)	(2.0)	(1.4)	(0.4)	(0.6)	(0.5)	(0.9)	(0.6)	(100.0)				
妊娠腎/妊娠中毒症	44	57	70	77	58	52	56	63	60	109		70	55	16	9	4	12	12	14	9	847	9	856	182.96	193.16
(%)	(5.2)	(6.7)	(8.3)	(9.1)	(6.8)	(6.1)	(6.6)	(7.4)	(7.1)	(12.9)		(8.3)	(6.5)	(1.9)	(1.1)	(0.5)	(1.4)	(1.4)	(1.7)	(1.1)	(100.0)				
その他の分類不能の腎炎	37	52	56	48	63	63	69	54	52	95	1	81	46	23	16	6	4	3	6	17	792	8	800	193.35	225.36
(%)	(4.7)	(6.6)	(7.1)	(6.1)	(8.0)	(8.0)	(8.7)	(6.8)	(6.6)	(12.0)	(0.1)	(10.2)	(5.8)	(2.9)	(2.0)	(0.8)	(0.5)	(0.4)	(0.8)	(2.1)	(100.0)				
多発性囊胞腎	255	408	422	491	552	529	505	482	448	998	15	636	340	202	158	23	45	56	93	71	6,729	118	6,847	185.38	176.29
(%)	(3.8)	(6.1)	(6.3)	(7.3)	(8.2)	(7.9)	(7.5)	(7.2)	(6.7)	(14.8)	(0.2)	(9.5)	(5.1)	(3.0)	(2.3)	(0.3)	(0.7)	(0.8)	(1.4)	(1.1)	(100.0)				
腎硬化症	620	1,182	1,339	1,427	1,441	1,401	1,485	1,326	1,132	2,803	43	1,558	941	545	332	82	132	131	214	156	18,290	447	18,737	178.41	165.06
(%)	(3.4)	(6.5)	(7.3)	(7.8)	(7.9)	(7.7)	(8.1)	(7.2)	(6.2)	(15.3)	(0.2)	(8.5)	(5.1)	(3.0)	(1.8)	(0.4)	(0.7)	(0.7)	(1.2)	(0.9)	(100.0)				
悪性高血圧	53	82	110	101	141	102	100	108	101	233	1	122	82	62	39	10	8	13	31	20	1,519	27	1,546	194.05	179.63
(%)	(3.5)	(5.4)	(7.2)	(6.6)	(9.3)	(6.7)	(6.6)	(7.1)	(6.6)	(15.3)	(0.1)	(8.0)	(5.4)	(4.1)	(2.6)	(0.7)	(0.5)	(0.9)	(2.0)	(1.3)	(100.0)				
糖尿病性腎症	3,222	5,352	5,665	5,832	6,121	5,902	5,755	5,415	4,840	10,951	141	6,276	3,555	1,988	1,072	239	402	466	738	414	74,346	2,053	76,399	166.92	145.21
(%)	(4.3)	(7.2)	(7.6)	(7.8)	(8.2)	(7.9)	(7.7)	(7.3)	(6.5)	(14.7)	(0.2)	(8.4)	(4.8)	(2.7)	(1.4)	(0.3)	(0.5)	(0.6)	(1.0)	(0.6)	(100.0)				
SLE腎炎	67	82	80	106	108	82	83	87	77	180		110	57	40	15	7	5	15	5	17	1,223	37	1,260	175.20	168.98
(%)	(5.5)	(6.7)	(6.5)	(8.7)	(8.8)	(6.7)	(6.8)	(7.1)	(6.3)	(14.7)		(9.0)	(4.7)	(3.3)	(1.2)	(0.6)	(0.4)	(1.2)	(0.4)	(1.4)	(100.0)				
アミロイド腎	7	25	22	28	29	18	12	15	22	27		11	10	7	5	1	2	2	2		245	7	252	150.38	123.24
(%)	(2.9)	(10.2)	(9.0)	(11.4)	(11.8)	(7.3)	(4.9)	(6.1)	(9.0)	(11.0)		(4.5)	(4.1)	(2.9)	(2.0)	(0.4)	(0.8)	(0.8)	(0.8)		(100.0)				
痛風腎	25	36	37	46	60	49	42	38	51	97	1	63	24	27	14	3	4	7	16	9	649	15	664	198.02	199.59
(%)	(3.9)	(5.5)	(5.7)	(7.1)	(9.2)	(7.6)	(6.5)	(5.9)	(7.9)	(14.9)	(0.2)	(9.7)	(3.7)	(4.2)	(2.2)	(0.5)	(0.6)	(1.1)	(2.5)	(1.4)	(100.0)				
先天性代謝異常による腎不全	9	13	7	16	11	9	10	7	7	16	1	14	8	5	3	1	1	1	3		142	4	146	172.15	144.27
(%)	(6.3)	(9.2)	(4.9)	(11.3)	(7.7)	(6.3)	(7.0)	(4.9)	(4.9)	(11.3)	(0.7)	(9.9)	(5.6)	(3.5)	(2.1)	(0.7)	(0.7)	(0.7)	(2.1)		(100.0)				
腎・尿路結核	10	9	3	9	12	12	6	8	8	17		7	7	2	2		2				114	2	116	152.38	112.75
(%)	(8.8)	(7.9)	(2.6)	(7.9)	(10.5)	(10.5)	(5.3)	(7.0)	(7.0)	(14.9)		(6.1)	(6.1)	(1.8)	(1.8)		(1.8)				(100.0)				
腎・尿路結石	16	19	28	31	33	25	31	24	22	49		40	14	10	7		1	1	2	6	361	9	370	195.46	366.30
(%)	(4.4)	(5.3)	(7.8)	(8.6)	(9.1)	(6.9)	(8.6)	(6.6)	(6.1)	(13.6)		(11.1)	(3.9)	(2.8)	(1.9)	(0.6)	(0.3)	(0.3)	(0.6)	(1.7)	(100.0)				
腎・尿路腫瘍	18	37	40	55	45	34	53	37	34	78	1	51	26	9	13	1	10	5	7	3	557	13	570	175.84	155.93
(%)	(3.2)	(6.6)	(7.2)	(9.9)	(8.1)	(6.1)	(9.5)	(6.6)	(6.1)	(14.0)	(0.2)	(9.2)	(4.7)	(1.6)	(2.3)	(0.2)	(1.8)	(0.9)	(1.3)	(0.5)	(100.0)				
閉塞性尿路障害	16	34	29	25	26	24	26	23	23	72		37	21	10	9		5	5	16	9	410	8	418	221.35	319.81
(%)	(3.9)	(8.3)	(7.1)	(6.1)	(6.3)	(5.9)	(6.3)	(5.6)	(5.6)	(17.6)		(9.0)	(5.1)	(2.4)	(2.2)		(1.2)	(1.2)	(3.9)	(2.2)	(100.0)				
骨髄腫	8	12	15	8	12	12	11	10	15	28		5	6	5	3			1	1		152	4	156	152.81	116.51
(%)	(5.3)	(7.9)	(9.9)	(5.3)	(7.9)	(7.9)	(7.2)	(6.6)	(9.9)	(18.4)		(3.3)	(3.9)	(3.3)	(2.0)			(0.7)	(0.7)		(100.0)				
腎形成不全	19	20	25	22	15	14	20	27	27	49		39	15	18	8	1	1	5	10	6	341	3	344	209.00	195.53
(%)	(5.6)	(5.9)	(7.3)	(6.5)	(4.4)	(4.1)	(5.9)	(7.9)	(7.9)	(14.4)		(11.4)	(4.4)	(5.3)	(2.3)	(0.3)	(0.3)	(1.5)	(2.9)	(1.8)	(100.0)				
不明	719	1,178	1,284	1,411	1,403	1,444	1,293	1,143	1,153	2,553	33	1,471	890	458	332	86	111	124	219	195	17,500	447	17,947	178.65	178.79
(%)	(4.1)	(6.7)	(7.3)	(8.1)	(8.0)	(8.3)	(7.4)	(6.5)	(6.6)	(14.6)	(0.2)	(8.4)	(5.1)	(2.6)	(1.9)	(0.5)	(0.6)	(0.7)	(1.3)	(1.1)	(100.0)				
再導入	65	66	82	76	78	102	71	82	67	151	2	106	55	31	28	3	10	11	24	24	1,134	19	1,153	199.06	226.10
(%)	(5.7)	(5.8)	(7.2)	(6.7)	(6.9)	(9.0)	(6.3)	(7.2)	(5.9)	(13.3)	(0.2)	(9.3)	(4.9)	(2.7)	(2.5)	(0.3)	(0.9)	(1.0)	(2.1)	(2.1)	(100.0)				
その他	190	316	373	365	399	408	329	319	300	681	6	437	242	127	73	16	36	45	57	53	4,772	115	4,887	178.06	165.28
(%)	(4.0)	(6.6)	(7.8)	(7.6)	(8.4)	(8.5)	(6.9)	(6.7)	(6.3)	(14.3)	(0.1)	(9.2)	(5.1)	(2.7)	(1.5)	(0.3)	(0.8)	(0.9)	(1.2)	(1.1)	(100.0)				
合計	8,134	12,510	13,574	14,458	14,933	14,627	14,159	13,177	11,995	27,419	344	16,035	9,403	5,405	3,211	784	1,231	1,367	2,313	1,791	186,870	4,401	191,271	177.13	169.01
(%)	(4.4)	(6.7)	(7.3)	(7.7)	(8.0)	(7.8)	(7.6)	(7.1)	(6.4)	(14.7)	(0.2)	(8.6)	(5.0)	(2.9)	(1.7)	(0.4)	(0.7)	(0.7)	(1.2)	(1.0)	(100.0)				
記載なし				1				1		1		1									4	1	5	176.50	80.86
(%)				(25.0)				(25.0)		(25.0)		(25.0)									(100.0)				
総計	8,134	12,510	13,574	14,459	14,933	14,627	14,159	13,178	11,995	27,420	344	16,036	9,403	5,405	3,211	784	1,231	1,367	2,313	1,791	186,874	4,402	191,276	177.13	169.01
(%)	(4.4)	(6.7)	(7.3)	(7.7)	(8.0)	(7.8)	(7.6)	(7.1)	(6.4)	(14.7)	(0.2)	(8.6)	(5.0)	(2.9)	(1.7)	(0.4)	(0.7)	(0.7)	(1.2)	(1.0)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy
2015年調査